


Please cite the Published Version

Andreyo, Evan, Unverzagt, Casey, Dos'Santos, Thomas  and Dawes, J. Jay (2024) Clinical Utility of Qualitative Change of Direction Movement Assessment in ACL Injury Risk Evaluation. *International Journal of Sports Physical Therapy*, 19 (10). pp. 1263-1278. ISSN 2159-2896

DOI: <https://doi.org/10.26603/001c.123483>

Publisher: North American Sports Medicine Institute

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/636107/>

Usage rights:  [Creative Commons: Attribution-Noncommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

Additional Information: This is an open access article published in *International Journal of Sports Physical Therapy*, by North American Sports Medicine Institute.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Clinical Commentary/Current Concept Review

Clinical Utility of Qualitative Change of Direction Movement Assessment in ACL Injury Risk Evaluation

Evan Andreyo^{1a}, Casey Unverzagt², Thomas Dos'Santos³, J. Jay Dawes⁴¹ Health Sciences, Rocky Mountain University of Health Professions, ² Physical Therapy, Baylor University, ³ Exercise and Sports Science, Manchester Metropolitan University, ⁴ School of Kinesiology, Oklahoma State University

Keywords: anterior cruciate ligament (ACL), injury prevention, return to sport, change of direction, movement assessment

<https://doi.org/10.26603/001c.123483>

International Journal of Sports Physical Therapy

Vol. 19, Issue 10, 2024

Anterior cruciate ligament (ACL) injuries are complex and influenced by numerous internal and external risk factors that should be considered to effectively mitigate injury and facilitate informed return to sport decision-making. Among these risk factors, movement quality exhibited during sport-specific tasks has been identified as a significant predictor of injury occurrence. Particularly, change of direction (COD) movements, when performed with sub-optimal movement quality, such as knee valgus and lateral trunk flexion, are prominent mechanisms of ACL injury in multidirectional sports. Unfortunately, the formal and objective assessment of COD movement quality is underutilized in clinical and sports practice, with existing methods often confined to expensive, sophisticated laboratory settings impractical for everyday clinicians. The purpose of this clinical commentary is to demonstrate the necessity of integrating COD movement assessments to screen for potential ACL injury risk, particularly among higher-risk populations. The authors will review cost-effective and clinic-friendly objective tests used to qualitatively screen COD movements, such as the Cutting Movement Assessment Score and The Expanded Cutting Alignment Tool. Additionally, this commentary will discuss key considerations when assessing COD movement.

Level of Evidence

5

ANTERIOR CRUCIATE LIGAMENT INJURIES

Anterior cruciate ligament (ACL) injuries are prevalent in athletic populations, with approximately 80,000 to 250,000 ACL injuries occurring annually in the United States alone.¹ Approximately 100,000 individuals undergo ACL reconstructions (ACLR) annually, making it the sixth most common orthopedic procedure in the United States.¹ The incidence of ACL injuries and ACLR have increased over the last two decades, particularly among young female athletes.²⁻⁴ The consequences of an ACL injury are substantial, with most injuries being season-ending and resulting in the most missed games compared to other injuries across competition levels.^{5,6} Additionally, only 69-81% of athletes return to sport, with only 55% returning to a competitive level.⁷⁻⁹ Moreover, there is a particularly high rate of ACL re-injury, reported as high as one out of five, with a six

times greater likelihood of ACL injury compared to healthy controls.^{10,11} An ACL injury can also result in a significant economic burden, psychological consequences, and an increased likelihood of developing knee osteoarthritis.¹²⁻¹⁶ Consequently, further investigation regarding the factors that may predispose an individual to this injury and available strategies to screen and mitigate injury risk are warranted.

ACL INJURY RISK FACTORS

Up to 75% of sports ACL injuries occur without physical contact (i.e., non-contact) with another player or object, most often during deceleration and a change of direction (COD) through a planted lower limb.^{1,17,18} These injuries are most prevalent in multidirectional sports that require frequent COD movements, such as soccer, basketball,

^a Corresponding Author:

Evan Andreyo, PT, DPT, PhD

Department of Health Sciences, Rocky Mountain University of Health Professions, Provo, UT, 84606, USA

Email Address: evan.andreyo@augie.edu

lacrosse, and football.¹⁹⁻²² Females participating in school-aged sports have a significantly greater risk of experiencing a non-contact ACL injury when compared to males at the same age and sport.¹⁹⁻²⁴ Female athletes have been reported to have an ACL injury incidence rate (IR) of 1.88 per 10,000 athletic exposures (AEs) compared to 0.87 per 10,000 AEs among male athletes across sports such as basketball, lacrosse, and soccer.²² The overall injury incidence rate ratio (IRR) for female athletes compared to males is reported to be 1.40 to 3.00.^{20,22,25,26} IRR is the ratio of injuries per AEs in females compared to males. Considering these findings, clinicians should evaluate why certain populations are at a greater risk of ACL injury than others and what appropriate and actionable mitigation strategies may be.

When considering what factors may be involved in this sex disparity in injury incidence, one must acknowledge that injury risk is multifactorial in nature. A review by Ruddy et al.²⁷ highlights the complexity of team sport injuries and the need to account for multiple variables when assessing injury risk rather than a reductionist approach. This complexity is particularly true for ACL injuries, with intrinsic (non-modifiable) and extrinsic (modifiable) factors influencing injury risk. There are various intrinsic characteristics more common in females that may promote a greater predisposition to an ACL injury, including hormonal influences, intercondylar femoral notch width, anterior knee laxity, and tibial plateau slope.^{28,29} However, these non-modifiable risk factors offer limited opportunities for strategizing and implementing ACL injury mitigation compared to their modifiable counterparts.

Poor neuromuscular control during landing and COD movements are common modifiable contributors to ACL injury risk.^{1,17,30-32} Neuromuscular control is defined by Herrington et al.³³ as “the ability to deliver a movement in an ‘optimal’ manner which minimizes loading stresses or maximizes the distribution of loading stresses on the tissues.” Suboptimal trunk and lower extremity movement quality during COD or landing tasks have been identified as visible characteristics of non-contact ACL injury and are determinants of potentially hazardous knee joint loads that increase ACL loading.³⁴⁻³⁹ Specifically, a dynamic knee valgus movement during deceleration or COD can pose a multiplanar load to the ACL and, when it exceeds its load capacity, result in injury.^{17,34,40,41} A knee abduction moment (KAM) describes the kinetic forces applied to the knee during knee valgus.^{32,42} Several authors have identified an increased KAM, increased hip adduction, decreased hip and knee flexion, and increased hip internal rotation in females compared to their male counterparts in both cutting and landing tasks common in sports participation, which may partially account for the greater relative ACL injury risk observed in females.^{35,43-49}

Several studies using video analysis have identified knee valgus as a prominent movement pattern during sport-related ACL injury, with some authors reporting knee valgus during 81-88% of non-contact ACL injuries.^{34,36,37,50} Additionally, frontal and sagittal plane trunk positioning, lateral foot width placement, knee flexion angle, and penultimate

foot contact braking have been demonstrated to significantly influence knee joint loading during COD tasks.⁵¹ These findings have also been recognized for second ACL injuries, with Paterno et al.^{30,31} reporting increased lower extremity neuromuscular control deficits and knee valgus during postural control and landing tasks among those going on to experience a second ACL injury compared to those who did not. Collectively, these findings suggest that specific movement characteristics and quality deficits, particularly with increased KAM, can potentially lead to increased strain on the ACL.

ASSESSING MOVEMENT QUALITY

The assessment of tasks associated with ACL injury (i.e., jump landing and changing direction) is crucial for mitigating first and subsequent ACL injuries, especially in female athletes participating in multidirectional sports. While a review by Arundale et al.⁵² highlights that no test can solely predict ACL injury, the utilization of movement screening may offer insight into the presence or absence of specific movement patterns associated with ACL injury, which, in theory, increases the relative risk of injury. Several strategies have been investigated and recommended to identify modifiable movement patterns associated with ACL injury risk.^{36,44,50,53} With better insight into movement quality, clinicians can implement targeted exercise-based knee and ACL injury prevention strategies to promote better movement control and injury risk reduction, an approach outlined by Dos’Santos et al.⁴² supported by strong evidence in the Exercise-Based Knee and ACL Injury Prevention Clinical Practice Guidelines.⁵⁴ Furthermore, an assessment of movement quality can also serve as an important component of a testing battery when making return to sport decisions. While the use of return to sport testing has inconsistent predictive value in determining reinjury after ACLR, this finding should only bolster the need to ensure the use of comprehensive and individualized testing.^{55,56} Several review publications recommend assessing movement quality as part of a larger testing battery to quantify an athlete’s readiness for return to sport after ACLR.⁵⁷⁻⁶¹

MOVEMENT ASSESSMENT METHODS

Researchers have demonstrated the use of 3-dimensional (3D) movement analysis coupled with force plate technology to assess kinetic and kinematic variables, serving as a reference standard for other measures of movement quality.^{53,62-65} A 3D movement analysis can allow for a thorough evaluation of an athlete’s neuromuscular control during sport-related movements, with KAM serving as a significant predictor and surrogate measure for ACL injury.³² While certainly the gold standard, utilizing 3D movement analysis requires extensive time, equipment, trained staff, and costs that are not always practical or realistic for those who are within a clinical setting working with a greater number of athletes, fewer staff, and fewer resources.³² As an alternative to 3D movement analysis, 2-dimensional (2D) video analysis has been proposed as

a means of assessing movement quality. McLean et al.⁶⁶ compared 2D and 3D movement assessments to screen for movements associated with ACL injury risk. Their study demonstrated strong correlations between 2D and 3D assessment of knee valgus during side-step and side-jump maneuvers. Several other authors have proposed clinic-friendly qualitative 2D video or real-time assessment tools to identify movement patterns associated with ACL injuries, including different variations of a single leg squat,^{67, 68} the Drop Vertical Jump (DVJ),^{32,69} the Landing Error Scoring System (LESS),^{62,70} the Tuck Jump Assessment (TJA),^{71,72} and the Vail Sports Test.^{73,74}

TASK SPECIFICITY

While the various 2D video and real-time assessment tools presented offer clinical utility, they are predominantly limited to assessing vertical landing or squatting mechanics. Indeed, these tasks are important and have application, particularly for screening athletes who participate in jump-landing sports such as basketball, volleyball, and netball. However, none of the included assessments focus on COD movements. This is pertinent as changing direction is biomechanically different from jump-landing, is a common movement in multidirectional sports, and directional changes are a prominent mechanism for ACL injury.^{17,34, 37,40,41,75} Movement assessment may be task-dependent, with several studies demonstrating differences in movement quality between drop landing and COD tasks.⁷⁶⁻⁷⁸ An individual who displays sub-optimal mechanics or higher knee joint loads during landing may not necessarily display poor mechanics and higher knee joint loads during COD and vice versa.⁷⁶⁻⁷⁸ Cowley et al.⁷⁹ identified that basketball and soccer players had increased knee valgus with a 45-degree COD compared to the drop vertical jump. Additionally, King et al.⁸⁰ reported that biomechanical differences during sidestep maneuvers were identified nine months after ACLR in a group of 156 subjects. These findings suggest that a qualitative assessment of movement during a COD task is warranted to screen for potential first and second ACL injury risk among athletes who must replicate these movements in their sport.

Despite the evidence favoring the utilization of a qualitative 2D assessment of COD, two literature reviews reported that this is rarely assessed in clinical practice after ACLR.^{58,81} Burgi et al.⁵⁸ suggest that assessing movement quality or other performance-based tests may be less commonly used than impairment-based measures (like strength and knee laxity) because of the potential need for equipment, large amounts of space, and lack of testing standardization. However, the authors point out that impairment-based measures may not relate as strongly to sports participation as performance-based measures. The specificity of sport-specific movement testing requires consideration, given the frequency of COD actions in multidirectional sports and the propensity to generate hazardous mechanical loads with poor movement quality.

ASSESSING CHANGE OF DIRECTION MOVEMENT QUALITY

Authors commonly use terms such as COD, agility, quickness, and cutting, often with no consistency in definition and context.⁸² The *Essentials of Strength Training and Conditioning* textbook by Haff and Triplett⁸³ defines COD as a situation when an individual must utilize “skills and abilities needed to change movement direction, velocity, or modes.” Dos’Santos et al.⁸⁴ identified several COD strategies, including a sidestep, crossover cut, split step, pivot, and shuffle step. There are also multiple means of formally assessing COD. As previously mentioned, while often the gold standard, 3D movement analysis and force plates are not always realistic in clinical settings.³² By contrast, Welling et al.⁸⁵ propose various on-field tests to consider after ACLR. However, of the tests identified, only one of the proposed tests formally assesses athletes for movements that may predispose them to an ACL injury.^{63,64} Furthermore, work conducted by Nimphius et al.⁸⁶ provides an overview of over 40 tests of non-reactive and reactive COD that can be used to assess an athlete’s performance, typically through time to completion. However, while the authors advocate using qualitative movement assessment in conjunction with quantitative performance, none of the included tests offer a formalized or objective means of doing so.

Several methods of objectively assessing 2D movement quality during a COD task have been proposed, with varying methodologies and populations. The Cutting Movement Assessment Score (CMAS),^{42,63,64} the Expanded Cutting Alignment Scoring Tool (E-CAST),^{87,88} and tests proposed by Della Villa et al.,⁶⁵ Di Paolo et al.,⁵³ and Weir et al.⁸⁹ all utilize 2D video analysis to assess COD movements.⁸⁹ Each assessment entails subjects performing between a 45 and 90-degree side step maneuver or a transition to a backward sprint, with movement quality later assessed using frontal and sagittal plane 2D video.

Among these 2D COD assessments, the CMAS and the E-CAST have demonstrated the most promising concurrent validity compared to 3D movement analysis, as well as inter-rater and intra-rater reliability.^{63,64,87,88,90-92} Additionally, these assessments can be set up and performed in a way that is feasible for clinical practice, requiring minimal equipment. As for the remaining tests by Weir et al.,⁸⁹ Della Villa et al.,⁶⁵ and Di Paolo et al.,⁵³ some considerations may make their clinical utility challenging. For example, the 2D movement analysis by Weir et al.⁸⁹ has fewer studies investigating its reliability and requires the use of manual measurements and digitization, which can be time-consuming. Tools proposed by Della Villa et al.⁹³ and Di Paolo et al.⁵³ both require more complex use of vectors through force platforms in addition to measuring 2D angles during video analysis. Thus, for the purpose of this commentary, the authors will focus on qualitative movement screening tools that require subjective evaluation of techniques and postures.




Figure 1. Setup for the Cutting Movement Assessment Score (CMAS) as demonstrated in Dos'Santos et al.⁴²

The green arrow and cameras are used for a right foot cut, and the orange arrow and cameras are used for a left foot cut. The blue arrow and camera are used for both.

THE CUTTING MOVEMENT ASSESSMENT SCORE (CMAS)

The CMAS is a validated screening tool that utilizes a 9-item scoring rubric (Appendix 1) to assess an athlete's movement quality during a 45-90° COD maneuver.^{42,63,64} A higher score in this assessment represents the presence of a greater number of movements associated with ACL injury. Each of the nine items included in the CMAS, operationally defined by Dos'Santos et al.,⁴² is based on technical determinants of KAM during COD movement and has been demonstrated as having a significant influence on knee joint loading associated with ACL injury.^{51,63,64}

The CMAS can be performed with two high-speed cameras, cones, and the CMAS grading rubric, making it a more clinically practical tool than 3D movement analysis.⁴² In the testing procedure, subjects sprint 5 meters "as fast as possible" and perform a 45-90° side step cut at a designated area under video recording (Figure 1).⁴² Video sampling should be recorded using a minimum of two cameras capable of recording at least 100 frames per second (fps), a feature available for most smartphones and tablets.⁴² Dos'Santos et al.⁴² recommend using an additional camera 20° from the frontal plane view to reduce parallax error in subjects who pre-rotate. However, in a more recent study, Jones et al.⁹⁴ identified excellent reliability for total CMAS scores with only two camera angles, as long as one included the sagittal plane view. Practitioners should evaluate 2-3 trials per limb for each athlete, with video footage ideally viewed in software that enables videos to be played at different speeds and be viewed frame-by-frame.⁴²

Several studies have reported the concurrent validity of the CMAS in identifying movements associated with ACL injury compared to 3D movement analysis.^{42,63,64} The overall inter-rater and intra-rater reliability of the CMAS has been moderate to excellent, with only one study by Aparicio-Sarmiento et al.⁹⁵ identifying poor reliability of the CMAS. Their findings may have been confounded by insufficient training regarding the use of the CMAS.⁹⁵ A study by Jones et al.⁹⁴ also assessed the reliability of the CMAS using sports scientists and medical practitioners, including physical therapists. They identified good to excellent inter-rater agreements across all practitioners for total scores ($K = 0.63-0.84$), offering promising implications for using the CMAS across disciplines. A summary of the CMAS validity and reliability is provided in Table 1. Dos'Santos et al.⁴² further proposed a stratification of injury risk based on a subject's total score: low risk is a CMAS ≤ 3 ; moderate risk is a CMAS 4-6; and high risk is a CMAS ≥ 7 , with athletes displaying scores ≥ 7 displaying greater knee joint loads and postures associated with ACL loading compared to athletes with lower scores.

THE EXPANDED CUTTING ALIGNMENT SCORING TOOL (E-CAST)

The E-CAST is a validated screening tool that uses a 6-item dichotomous scoring rubric (Appendix 2) to assess movement quality during a 45° COD maneuver, with a higher score representing a greater number of movements associated with ACL injury.^{88,91} A score of "1" is awarded for each item when the movement fault is present and the total score is determined.^{51,87} The E-CAST is an expansion from

Table 1. A comparison between the CMAS and the E-CAST

	CMAS ^{42,63,64}	E-CAST ^{87,91,92,96}
Number of Items	9	6
Scoring	Items 2 and 6 scored 0, 1, or 2 All remaining items scored 0 or 1	All items scored 0 or 1
Number of Cameras	2 or 3	2
Proposed recording quality	≥100fps	60fps; 1080p
Cutting Angle	45 to 90°	45°
Instructions for subject forward sprint effort	“as fast as possible”	80% maximum effort
Approach distance	5 meters	8.8 meters
Concurrent Validity compared to 3D movement analysis	<p>Dos’Santos et al.⁶⁴ Spearman’s correlation revealed a significant and very large association between CMAS and peak KAMs. ($\rho = 0.796$, 95% CI = 0.647-0.887, $p < 0.001$) Significantly greater cutting multiplanar knee joint loads were demonstrated by subjects with higher CMASs compared to lower, with moderate to very large effect sizes.</p> <p>Jones et al.⁶³ Spearman’s correlation revealed a significant large association between CMAS and KAMs ($\rho = 0.633$; $p < 0.001$).</p>	<p>Butler et al.⁹¹ Each item demonstrated sensitivity and specificity ranging from 70-85% and 55-89%, respectively. Across items, the area under the curve ranged from 0.67 to 0.91.</p>
Reliability	<p>Jones et al.⁹⁴ Good to excellent inter-rater reliability for total score ($k=0.63-0.84$)</p> <p>Olivares-Jabalera et al.⁹⁷ Excellent intra-rater reliability for total score (ICC = 0.70) Moderate inter-rater reliability for total scores (ICC = 0.58)</p> <p>Needham and Herrington⁹⁰ Excellent intra-rater reliability for total score (ICC = 0.98)</p> <p>Aparicio-Sarmiento et al.⁹⁵ Poor inter-rater reliability for total score (ICC = 0.11-0.45) Moderate intra-rater reliability for total score (ICC = 0.71)</p> <p>Dos’Santos et al.⁶⁴ Excellent intra-rater reliability for total score (ICC = 0.95) Moderate inter-rater reliability for total scores (ICC = 0.69)</p> <p>Jones et al.⁶³ Excellent intra-rater reliability for total scores (ICC = 0.922) Excellent inter-rater reliability for total scores (ICC = 0.913)</p>	<p>Butler et al.⁸⁷ Good intra-rater reliability for total score (ICC=0.78) Moderate inter-rater reliability for total score (ICC=0.71)</p> <p>Butler et al.⁹² Good intra-rater reliability for total scores (ICC=0.821) Good inter-rater reliability (ICC=0.752)</p>

the 4-item Cutting Alignment Scoring Tool (CAST) that originally only included a frontal plane assessment of trunk lean, cut width, and knee valgus.⁸⁷ The expanded version now includes sagittal plane assessments of knee flexion angle and ankle plantar flexion.⁸⁸ Butler et al.⁹² compared the use of the E-CAST qualitative scoring rubric against a quantitative version requiring 2D kinematic measurements and found no significant differences between the two, suggesting that either can serve as a reliable means of assessing movement quality.

The E-CAST can be performed in a means similar to that of the CMAS using two high-speed cameras, cones, and the E-CAST grading rubric, making it more practical than 3D movement analysis.⁹² Subjects sprint 8.8 meters at 80% of their maximum speed towards an “opponent cone” placed

just beyond the pivoting area (Figure 2). A side step cut is performed at a 45° angle under video recording.⁸⁷ Previous investigations of the E-CAST recorded video at 60fps with 1080p quality that was later slowed down by 50% for video analysis.⁸⁷ Subjects completed three trials per direction, with one randomly selected for analysis.^{87,91}

Butler et al.^{88,92} have demonstrated the concurrent validity of the E-CAST with 3D movement analysis as well as good inter- and intra-rater reliability (ICC=0.752 and ICC=0.821).^{88,91} The original 4-item CAST demonstrated good inter-rater and intra-rater reliability (ICC=0.808 and ICC=0.753) among medical doctors, physical therapists, and athletic trainers.⁸⁷ However, the utility of the E-CAST has only been assessed with physical therapists as the adminis-




Figure 2. Setup for the Expanded Cutting Alignment Scoring Tool as demonstrated in Butler et al.⁹⁶

trators and raters.^{88,91,92} A summary of the E-CAST validity and reliability is provided in [Table 1](#).

The CMAS and the E-CAST can both potentially serve as a practical and insightful standardized assessment of ACL injury risk with COD movements. Given their clinic-friendly utility and objective information about COD movement quality, implementing one of these assessment tools is merited to screen for both first ACL injuries and after injury as part of a return to sport testing battery. While The CMAS and the E-CAST have similar qualities, they also carry noteworthy differences. For example, they each have a different number of items (9 vs. 6), with the CMAS including an assessment of penultimate foot contact and foot rotation, each correlated with knee joint loads.^{51,63,64,88} Further examples of the differences between the CMAS and the E-CAST are outlined in [Table 1](#). Each clinician must assess the unique needs and movement characteristics of the individual athlete they are working with, their respective sport, and the set up most conducive to the clinician's setting.

LIMITATIONS

A significant limitation of the presented 2D COD movement analysis tools is that there is currently no prospective evidence showing they can predict injury. While the evidence strongly supports the concurrent validity and reliability of the presented measures, a further direction of research should be determining their predictive validity, particularly for ACL injury. Despite mixed validity for the utilization of return to sport testing as a whole, it is still advisable to screen for movements associated with ACL injury in sport-specific movement patterns, as opposed to limiting screening to time from injury and basic clinical examination measures.^{58,98-102} Some investigations have suggested that

COD biomechanical issues may still be present longer than strength deficits post-op ACLR and that improved movement quality during return to sport testing is associated with a decreased risk of second ACL injury.^{80,101}

Additional limitations of the provided movement analysis tools are the potential barriers related to space, time, and cost. The CMAS requires an open space that is roughly 8x10 meters, with the E-CAST requiring roughly 13x7 meters. Many clinicians may not have access to an open area this size if they are limited to a small office already filled with treatment tables and equipment. Also, performing all the required practice trials and valid trials does take time, an indispensable resource in clinical practice. However, Dos'Santos et al.⁴² have reported completing six CMAS trials in groups of athletes as large as twelve in less than 15 minutes. This suggests that, with experience, testing can be completed in a reasonable time. Lastly, clinicians must consider equipment costs. Utilizing timing gates and high-end cameras comes at a significant economic cost that may not be attainable for practitioners. Fortunately, the test setup can be completed with cones and with the use of any recording device capable of recording at a higher frame rate, a feature available on all iPad and iPhone devices. When considering these limitations, not all clinics may be capable of conducting these tests. Patients may need to be referred to specific testing sites with the appropriate space and equipment, in line with recommendations by Unverzagt et al.⁵⁹

CONSIDERING ADDITIONAL VARIABLES

Marques et al.¹⁰³ offer insight into the application of COD movement assessment after ACLR. In their review, they do not emphasize a specific tool such as the CMAS or E-CAST,

but they do identify key variables to consider when utilizing field and laboratory-based COD testing, such as planned vs. unplanned, approach velocity, and cutting angle.

PLANNED VS. UNPLANNED MOVEMENT

An additional consideration for the assessment of COD movement quality should be whether the COD is planned or unplanned. Sports participation often requires an athlete to process multiple sources of information and stimuli when executing sporting actions. In the context of COD, athletes often need to respond to an external stimulus, such as an opponent or a ball, that requires externally directed attention.¹⁰⁴ Several researchers have suggested that this external focus on unplanned stimuli increases neurocognitive loads, presenting a particular after ACL injury given changes in neuroplasticity and neurocognitive compensation.¹⁰⁴⁻¹⁰⁸ Sheppard et al.⁸² propose that “In order to be considered ‘reactive’ agility, the movement should not only involve a change in speed and direction, but must also be an open skilled task, involving a reaction to a stimulus.”

The performance of COD tasks under reactive conditions has been proposed as a uniquely different task compared to pre-planned COD, entailing different cognitive challenges in response to spatial and temporal stimuli.^{82,109,110} Unplanned COD may be more representative of sports participation, given the requirements in team sports to respond to an external stimulus, such as an opponent or ball, without pre-planning. Without a reactive element, movement assessment tools only replicate a portion of the conditions and scenarios common in sports and ACL injuries. While most ACL injuries are non-contact, the majority occur with externally directed attention and increased neurocognitive loading, typically with an opponent or ball.^{36,50} Several authors have described that knee movement mechanics and timed performance differ in unplanned and planned movements.¹¹¹⁻¹¹⁵ Specifically, increased KAM and internal rotation moments have been identified at the knee under unplanned conditions compared to planned conditions, both associated with increased risk of ACL injury.^{111,113,116} Moreover, the 2016 Consensus on Return to Sport and additional authors have suggested that utilizing unplanned movements may be superior to incorporate over planned tasks alone.⁵⁷⁻⁶⁰ Separate reviews by Grooms et al.¹⁰⁸ and Wilk et al.⁶⁰ recommend utilizing neurocognitive and reactive testing to assess return to sport readiness after ACLR. The combined recommendations of these authors suggest that the formal assessment of movement quality under unplanned movements is warranted. However, while unplanned COD may hold many benefits, there are challenges with producing a valid, standardized sport-specific stimulus that has good reliability.

A study by Needham and Herrington⁹⁰ investigated differences in CMASs between planned and unplanned side-step maneuvers for female soccer players. They identified significantly greater total CMASs for unplanned COD maneuvers (5.53 ± 0.71) compared to planned maneuvers (3.55 ± 0.85 , $p < 0.012$). The findings of this study match previous results that unplanned movements may reveal a higher number of movement patterns associated with ACL

injury risk.^{111,113} This study suggests that a 2D COD movement assessment tool can offer utility in assessing both planned and unplanned conditions and that only assessing planned conditions may not wholly represent an athlete’s injury-risk profile.

There are noteworthy challenges with the implementation of unplanned movement assessment. There is often a tradeoff between ecological validity and standardization when selecting a stimulus for unplanned movement. In the study by Needham and Herrington,⁹⁰ a soccer ball pass was used to cue the subjects in their cutting direction. Their approach may promote greater ecological validity compared to light or sound-based stimuli. However, as acknowledged by Needham and Herrington,⁹⁰ this approach does carry limitations in its consistency of timing and trajectory, thus potentially influencing subsequent reaction times and movement patterns. By contrast, the utilization of light stimuli is easier to regulate but is lacking in sport specificity. Additionally, caution and a stepwise progression must be utilized when integrating unplanned COD maneuvers given the potential for increased knee joint loads, particularly in the context of return to sport testing after injury.

APPROACH VELOCITY

One additional variable that is pertinent in the assessment of COD is the velocity at which the individual enters into the COD task (i.e., approach velocity). Reviews by Marques et al.¹⁰⁵ and Dos’Santos et al.¹¹⁷ highlight that approach velocity significantly influences COD speed and biomechanics. Kristianslund et al.¹¹⁸ demonstrated that increased KAMs are seen with higher approach velocities compared to lower velocities among female athletes during COD maneuvers. Further studies have found that when subjects performed COD maneuvers with higher approach velocities, they exhibited greater knee joint stiffness and peak knee valgus.¹¹⁹⁻¹²¹ An investigation on injury mechanisms for ACL injuries in handball by Olsen et al.³⁴ also identified that non-contact ACL injuries commonly occurred at high approach velocities during COD tasks. When considering this in practical application, approach velocity should be considered when assessing movement quality, as those moving at faster speeds may be more likely to exhibit more potentially hazardous movement mechanics.^{103,117} As a potential means of mitigating injury and improving standardization in movement assessment, Vanrenterghem et al.¹²¹ suggest that velocity should be standardized to 4 meters per second.

CUTTING ANGLE

Both Dos’Santos et al.¹¹⁷ and Marques et al.¹⁰⁵ highlight that cutting angle is also a pertinent consideration when assessing COD movement quality. These authors identify that sharper COD angles (90 vs 45°) result in increases in knee joint loading and require lower approach velocities for optimal execution (i.e., COM angle deflection). Dos’Santos et al.¹¹⁷ identifies this as the “angle-velocity trade-off,” whereby faster approach velocities can compromise

the COD execution. Given the established correlation between approach velocity and COD movement quality, it must be accounted for as a variable for execution performance and should be considered when screening COD movement quality.

OPPORTUNITIES FOR INTERVENTION

The clinical assessment of COD movement quality to determine ACL injury risk would be limited in value if there were no opportunities for measurable improvement. Fortunately, COD movement quality has been shown to be modifiable. Clinicians should first utilize a cutting assessment tool to identify movement patterns associated with increased injury risk. While this may have standalone value in making return to sport decisions, clinicians should also use their findings to formulate impairment-based neuromuscular interventions to address underlying movement patterns. For example, testing results may indicate movement deficits related to poor trunk control, increased knee valgus, or a lack of knee flexion. Interventions aimed at improving muscle strength and neuromuscular control can be prescribed to mitigate the identified movement patterns.

Neuromuscular training has been demonstrated to improve proprioception and decrease biomechanical deficits.^{122,123} These strategies can vary considerably but generally include unilateral and bilateral lower limb and core exercises with visual, verbal, and/or tactile feedback to improve trunk and knee control.^{122,123} This approach can be integrated through foundational exercises, plyometrics, and cutting maneuvers. A review by Buckthorpe¹²⁴ provides recommendations for a movement retraining progression after ACLR, beginning with foundational movement training, then progressing to high-load 'sport-type' movement retraining, and lastly, integrating sport-specific movement retraining.

Studies by Olivares-Jabalera et al.¹²⁵ and Dos'Santos et al.¹²⁶ investigated the effectiveness of separate six-week technique modification programs to improve cutting and jump-landing movement quality among soccer players. Both studies identified significant pre-to-post improvements in CMAS scores after six weeks of COD movement training with individualized feedback from a practitioner. Additionally, a study by Nijmeijer et al.¹²⁷ identified improvements in CMAS scores when subjects were provided video and verbal feedback, particularly when they were given autonomy to select the timing in which they received it. These findings indicate that COD movement quality can be improved with task-specific training. Additionally, movement assessment can serve as a valuable test-re-test measure to track modifiable risk factors associated with ACL injury risk. Dos'Santos et al.¹²⁸ describe this process for movement assessment, treatment, and reassessment, depicted in [Figure 3](#). This process can be completed multiple times over the course of an individual's episode of care




Figure 3. A process for screening movement quality and applying interventions.

or season to track patient improvement and reduce injury risk.

SUMMARY

ACL injuries are formidable and consequential injuries in athletics that are influenced by considerable intrinsic and extrinsic risk factors.^{5,6,12-16} Of these risk factors, an individual's movement quality has been demonstrated as a significant variable associated with ACL injury risk, particularly among female athletes in multidirectional sports.^{1,17,19-23,26,30-32} While there are various means of assessing movement quality, an assessment during a COD maneuver provides task-specific insight that is a prevalent mechanism of ACL injury.⁷⁶⁻⁷⁸ The CMAS and the E-CAST both provide valid, reliable, and objective means of assessing movement quality during COD tasks worth considering when screening for ACL injury and re-injury risk.^{42,63,64,88,91,92,94} Additionally, variables such as planned and unplanned movement, approach velocity, and cutting angle should all be considered as pertinent influences on task performance and injury risk and must be considered when screening and profiling movement quality.^{82,103-110,117}

CONFLICTS OF INTEREST

The authors report no conflicts of interest.

Submitted: April 17, 2024 CDT, Accepted: August 28, 2024 CDT
© The Author(s)



This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CCBY-NC-4.0). View this license's legal deed at <https://creativecommons.org/licenses/by-nc/4.0> and legal code at <https://creativecommons.org/licenses/by-nc/4.0/legalcode> for more information.

REFERENCES

1. Logerstedt DS, Scalzitti D, Risberg MA, et al. Knee stability and movement coordination impairments: knee ligament sprain revision 2017: clinical practice guidelines linked to the international classification of functioning, disability and health from the orthopaedic section of the American Physical Therapy Association. *J Orthop Sports Phys Ther.* 2017;47(11):A1-A47. [doi:10.2519/jospt.2017.0303](https://doi.org/10.2519/jospt.2017.0303)
2. Zbrojkiewicz D, Vertullo C, Grayson JE. Increasing rates of anterior cruciate ligament reconstruction in young Australians, 2000–2015. *Med J Australia.* 2018;208(8):354-358. [doi:10.5694/mja17.00974](https://doi.org/10.5694/mja17.00974)
3. Sanders TL, Maradit Kremers H, Bryan AJ, et al. Incidence of anterior cruciate ligament tears and reconstruction: a 21-year population-based study. *Am J Sports Med.* 2016;44(6):1502-1507. [doi:10.1177/0363546516629944](https://doi.org/10.1177/0363546516629944)
4. Maniar N, Verhagen E, Bryant AL, Opar DA. Trends in Australian knee injury rates: An epidemiological analysis of 228,344 knee injuries over 20 years. *The Lancet Regional Health - Western Pacific.* 2022;21:100409. [doi:10.1016/j.lanwpc.2022.100409](https://doi.org/10.1016/j.lanwpc.2022.100409)
5. Agel J, Rockwood T, Klossner D. Collegiate ACL injury rates across 15 Sports: National Collegiate Athletic Association injury surveillance system data update (2004-2005 through 2012-2013). *Clin J Sport Med.* 2016;26(6):6. [doi:10.1097/ISM.0000000000000290](https://doi.org/10.1097/ISM.0000000000000290)
6. Drakos MC, Domb B, Starkey C, Callahan L, Allen AA. Injury in the National Basketball Association: A 17-Year overview. *Sports Health.* 2010;2(4):284-290. [doi:10.1177/1941738109357303](https://doi.org/10.1177/1941738109357303)
7. Ardern CL, Taylor NF, Feller JA, Webster KE. Fifty-five per cent return to competitive sport following anterior cruciate ligament reconstruction surgery: an updated systematic review and meta-analysis including aspects of physical functioning and contextual factors. *Br J Sports Med.* 2014;48(21):1543-1552. [doi:10.1136/bjsports-2013-093398](https://doi.org/10.1136/bjsports-2013-093398)
8. Randsborg PH, Cepeda N, Adamec D, Rodeo SA, Ranawat A, Pearle AD. Patient-reported outcome, return to sport, and revision rates 7-9 years after anterior cruciate ligament reconstruction: results from a cohort of 2042 patients. *Am J Sports Med.* 2022;50(2):423-432. [doi:10.1177/03635465211060333](https://doi.org/10.1177/03635465211060333)
9. Fones L, Kostyun RO, Cohen AD, Pace JL. Patient-reported outcomes, return-to-sport status, and reinjury rates after anterior cruciate ligament reconstruction in adolescent athletes: minimum 2-year follow-up. *Orthop J Sports Med.* 2020;8(11):232596712096447. [doi:10.1177/2325967120964471](https://doi.org/10.1177/2325967120964471)
10. Paterno MV, Rauh MJ, Schmitt LC, Ford KR, Hewett TE. Incidence of second ACL injuries 2 years after primary ACL reconstruction and return to sport. *Am J Sports Med.* 2014;42(7):1567-1573. [doi:10.1177/0363546514530088](https://doi.org/10.1177/0363546514530088)
11. Barber-Westin S, Noyes FR. One in 5 athletes sustain reinjury upon return to high-risk sports after ACL reconstruction: a systematic review in 1239 athletes younger than 20 years. *Sports Health.* 2020;12(6):587-597. [doi:10.1177/1941738120912846](https://doi.org/10.1177/1941738120912846)
12. Hewett TE. Preventive biomechanics: A paradigm shift with a translational approach to biomechanics. *J Sci Med Sport.* 2017;45(11):2654-2664. [doi:10.1016/j.jsams.2017.01.002](https://doi.org/10.1016/j.jsams.2017.01.002)
13. Baez SE, Hoch MC, Hoch JM. Psychological factors are associated with return to pre-injury levels of sport and physical activity after ACL reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2020;28(2):495-501. [doi:10.1007/s00167-019-05696-9](https://doi.org/10.1007/s00167-019-05696-9)
14. McPherson AL, Feller JA, Hewett TE, Webster KE. Psychological readiness to return to sport is associated with second anterior cruciate ligament injuries. *Am J Sports Med.* 2019;47(4):857-862. [doi:10.1177/0363546518825258](https://doi.org/10.1177/0363546518825258)
15. Poulsen E, Goncalves GH, Bricca A, Roos EM, Thorlund JB, Juhl CB. Knee osteoarthritis risk is increased 4-6 fold after knee injury – a systematic review and meta-analysis. *Br J Sports Med.* 2019;53(23):1454-1463. [doi:10.1136/bjsports-2018-100022](https://doi.org/10.1136/bjsports-2018-100022)
16. Lohmander LS, Englund PM, Dahl LL, Roos EM. The long-term consequence of anterior cruciate ligament and meniscus injuries: osteoarthritis. *Am J Sports Med.* 2007;35(10):1756-1769. [doi:10.1177/0363546507307396](https://doi.org/10.1177/0363546507307396)
17. Wetters N, Weber AE, Wuerz TH, Schub DL, Mandelbaum BR. Mechanism of injury and risk factors for anterior cruciate ligament injury. *Op Tech Sports Med.* 2016;24(1):2-6. [doi:10.1053/j.otsm.2015.09.001](https://doi.org/10.1053/j.otsm.2015.09.001)

18. Bates NA, Schilaty ND, Nagelli CV, Krych AJ, Hewett TE. Multiplanar loading of the knee and its influence on anterior cruciate ligament and medial collateral ligament strain during simulated landings and noncontact tears. *Am J Sports Med.* 2019;47(8):1844-1853. doi:10.1177/0363546519850165
19. Joseph AM, Collins CL, Henke NM, Yard EE, Fields SK, Comstock RD. A multisport epidemiologic comparison of anterior cruciate ligament injuries in high school athletics. *J Athl Train.* 2013;48(6):810-817. doi:10.4085/1062-6050-48.6.03
20. Gornitzky AL, Lott A, Yellin JL, Fabricant PD, Lawrence JT, Ganley TJ. Sport-specific yearly risk and incidence of anterior cruciate ligament tears in high school athletes: a systematic review and meta-analysis. *Am J Sports Med.* 2016;44(10):2716-2723. doi:10.1177/0363546515617742
21. Beynnon BD, Vacek PM, Newell MK, et al. The effects of level of competition, sport, and sex on the incidence of first-time noncontact anterior cruciate ligament injury. *Am J Sports Med.* 2014;42(8):1806-1812. doi:10.1177/0363546514540862
22. Montalvo AM, Schneider DK, Webster KE, et al. Anterior cruciate ligament injury risk in sport: a systematic review and meta-analysis of injury incidence by sex and sport classification. *J Athl Train.* 2019;54(5):472-482. doi:10.4085/1062-6050-407-16
23. Montalvo AM, Schneider DK, Yut L, et al. "What's my risk of sustaining an ACL injury while playing sports?" A systematic review with meta-analysis. *Br J Sports Med.* 2019;53(16):1003-1012. doi:10.1136/bjsports-2016-096274
24. Bram JT, Magee LC, Mehta NN, Patel NM, Ganley TJ. Anterior cruciate ligament injury incidence in adolescent athletes: a systematic review and meta-analysis. *Am J Sports Med.* 2021;49(7):1962-1972. doi:10.1177/0363546520959619
25. Zech A, Hollander K, Junge A, et al. Sex differences in injury rates in team-sport athletes: A systematic review and meta-regression analysis. *J Sport Health Sci.* 2022;11(1):104-114. doi:10.1016/j.jshs.2021.04.003
26. Bram JT, Magee LC, Mehta NN, Patel NM, Ganley TJ. Anterior cruciate ligament injury incidence in adolescent athletes: a systematic review and meta-analysis. *Am J Sports Med.* 2021;49(7):1962-1972. doi:10.1177/0363546520959619
27. Ruddy JD, Cormack SJ, Whiteley R, Williams MD, Timmins RG, Opar DA. Modeling the risk of team sport injuries: A narrative review of different statistical approaches. *Front Physiol.* 2019;10:829. doi:10.3389/fphys.2019.00829
28. Smith HC, Vacek P, Johnson RJ, et al. Risk factors for anterior cruciate ligament injury: a review of the literature—part 2: hormonal, genetic, cognitive function, previous injury, and extrinsic risk factors. *Sports Health.* 2012;4(2):155-161. doi:10.1177/1941738111428282
29. Smith HC, Vacek P, Johnson RJ, et al. Risk factors for anterior cruciate ligament injury: a review of the literature — part 1: neuromuscular and anatomic risk. *Sports Health.* 2012;4(1):69-78. doi:10.1177/1941738111428281
30. Paterno MV, Schmitt LC, Ford KR, et al. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med.* 2010;38(10):1968-1978. doi:10.1177/0363546510376053
31. Paterno MV, Kiefer AW, Bonnette S, et al. Prospectively identified deficits in sagittal plane hip-ankle coordination in female athletes who sustain a second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Clin Biomech.* 2015;30(10):1094-1101. doi:10.1016/j.clinbiomech.2015.08.019
32. Hewett TE, Myer GD, Ford KR, et al. Biomechanical measures of neuromuscular control and valgus loading of the knee predict anterior cruciate ligament injury risk in female athletes: a prospective study. *Am J Sports Med.* 2005;33(4):492-501. doi:10.1177/0363546504269591
33. Herrington L, Munro A, Jones P. Assessment of factors associated with injury risk. In: *Performance Assessment in Strength and Conditioning.* 1st ed. Routledge; 2018:53-95. doi:10.4324/9781315222813-7
34. Olsen OE, Myklebust G, Engebretsen L, Bahr R. Injury mechanisms for anterior cruciate ligament injuries in team handball: a systematic video analysis. *Am J Sports Med.* 2004;32(4):1002-1012. doi:10.1177/0363546503261724
35. Salci Y, Kentel BB, Heycan C, Akin S, Korkusuz F. Comparison of landing maneuvers between male and female college volleyball players. *Clin Biomech.* 2004;19(6):622-628. doi:10.1016/j.clinbiomech.2004.03.006

36. Della Villa F, Buckthorpe M, Grassi A, et al. Systematic video analysis of ACL injuries in professional male football (soccer): injury mechanisms, situational patterns and biomechanics study on 134 consecutive cases. *Br J Sports Med.* 2020;54(23):1423-1432. [doi:10.1136/bjsports-2019-101247](https://doi.org/10.1136/bjsports-2019-101247)
37. Johnston JT, Mandelbaum BR, Schub D, et al. Video analysis of anterior cruciate ligament tears in professional american football athletes. *Am J Sports Med.* 2018;46(4):862-868. [doi:10.1177/0363546518756328](https://doi.org/10.1177/0363546518756328)
38. Larwa J, Stoy C, Chafetz RS, Boniello M, Franklin C. Stiff Landings, Core Stability, and Dynamic Knee Valgus: A Systematic Review on Documented Anterior Cruciate Ligament Ruptures in Male and Female Athletes. *Int J Environ Res Pub Health.* 2021;18(7):3826. [doi:10.3390/ijerph18073826](https://doi.org/10.3390/ijerph18073826)
39. Saki F, Tahayori B, Bakhtiari Khou S. Female athletes with ligament dominance exhibiting altered hip and ankle muscle co-contraction patterns compared to healthy individuals during single-leg landing. *Gait Posture.* 2022;93:225-229. [doi:10.1016/j.gaitpost.2022.02.011](https://doi.org/10.1016/j.gaitpost.2022.02.011)
40. Hewett TE, Myer GD, Ford KR, Paterno MV, Quatman CE. Mechanisms, prediction, and prevention of ACL injuries: Cut risk with three sharpened and validated tools: ACL injury prevention. *J Orthop Res.* 2016;34(11):1843-1855. [doi:10.1002/jor.23414](https://doi.org/10.1002/jor.23414)
41. Krosshaug T, Nakamae A, Boden BP, et al. Mechanisms of anterior cruciate ligament injury in basketball: video analysis of 39 cases. *Am J Sports Med.* 2007;35(3):359-367. [doi:10.1177/0363546506293899](https://doi.org/10.1177/0363546506293899)
42. Dos'Santos T, Thomas C, McBurnie A, Donelon T, Herrington L, Jones PA. The cutting movement assessment score (CMAS) qualitative screening tool: application to mitigate anterior cruciate ligament injury risk during cutting. *Biomech.* 2021;1(1):83-101. [doi:10.3390/biomechanics1010007](https://doi.org/10.3390/biomechanics1010007)
43. Pollard CD, Davis IM, Hamill J. Influence of gender on hip and knee mechanics during a randomly cued cutting maneuver. *Clin Biomech.* 2004;19(10):1022-1031. [doi:10.1016/j.clinbiomech.2004.07.007](https://doi.org/10.1016/j.clinbiomech.2004.07.007)
44. Weir G, Stillman M, van Emmerik R, Wyatt H, Jewell C, Hamill J. Differences in Kinetics, Kinematics and Muscle Activation Strategies in Male and Female Team Sport Athletes During Unanticipated Sidestepping. *J Sci Sport Exerc.* 2019;1(2):159-167. [doi:10.1007/s42978-019-0019-2](https://doi.org/10.1007/s42978-019-0019-2)
45. Mclean SG, Lipfert SW, Van Den Bogert AJ. Effect of gender and defensive opponent on the biomechanics of sidestep cutting. *Med Sci Sports Exerc.* 2004;36(6):1008-1016. [doi:10.1249/01.MSS.0000128180.51443.83](https://doi.org/10.1249/01.MSS.0000128180.51443.83)
46. Malinzak RA, Colby SM, Kirkendall DT, Yu B, Garrett WE. A comparison of knee joint motion patterns between men and women in selected athletic tasks. *Clin Biomech.* 2001;16(5):438-445. [doi:10.1016/S0268-0033\(01\)00019-5](https://doi.org/10.1016/S0268-0033(01)00019-5)
47. Ogasawara I, Miyakawa S, Wakitani S. Gender difference in neuromuscular hip and knee joint control during single-leg landing. *MJHES.* 2014;4(1):1-11.
48. Ford KR, Myer GD, Hewett TE. Valgus Knee Motion during Landing in High School Female and Male Basketball Players. *Med Sci Sports Exerc.* 2003;35(10):1745-1750. [doi:10.1249/01.MSS.0000089346.85744.D9](https://doi.org/10.1249/01.MSS.0000089346.85744.D9)
49. Russell KA, Palmieri RM, Zinder SM, Ingersoll CD. Sex differences in valgus knee angle during a single-leg drop jump. *J Athl Train.* 2006;41(2):166-171.
50. Lucarno S, Zago M, Buckthorpe M, et al. Systematic video analysis of anterior cruciate ligament injuries in professional female soccer players. *Am J Sports Med.* 2021;49(7):1794-1802. [doi:10.1177/03635465211008169](https://doi.org/10.1177/03635465211008169)
51. Donelon TA, Dos'Santos T, Pitchers G, Brown M, Jones PA. Biomechanical determinants of knee joint loads associated with increased anterior cruciate ligament loading during cutting: a systematic review and technical framework. *Sports Med - Open.* 2020;6(1):53. [doi:10.1186/s40798-020-00276-5](https://doi.org/10.1186/s40798-020-00276-5)
52. Arundale AJH, Silvers-Granelli HJ, Myklebust G. ACL injury prevention: Where have we come from and where are we going? *J Orthop Res.* 2022;40(1):43-54. [doi:10.1002/jor.25058](https://doi.org/10.1002/jor.25058)
53. Di Paolo S, Zaffagnini S, Tosarelli F, et al. A 2D qualitative movement assessment of a deceleration task detects football players with high knee joint loading. *Knee Surg Sports Traumatol Arthrosc.* 2021;29(12):4032-4040. [doi:10.1007/s00167-021-06709-2](https://doi.org/10.1007/s00167-021-06709-2)
54. Arundale AJH, Bizzini M, Dix C, et al. Exercise-based knee and anterior cruciate ligament injury prevention: Clinical Practice Guidelines linked to the International Classification of Functioning, Disability and Health From the Academy of Orthopaedic Physical Therapy and the American Academy of Sports Physical Therapy. *J Orthop Sports Phys Ther.* 2023;53(1):CPG1-CPG34. [doi:10.2519/jospt.2023.0301](https://doi.org/10.2519/jospt.2023.0301)

55. Gill VS, Tummala SV, Sullivan G, et al. Functional return-to-sport testing demonstrates inconsistency in predicting short-term outcomes following anterior cruciate ligament reconstruction: A systematic review. *Arthroscopy*. Published online January 11, 2024. doi:10.1016/j.arthro.2023.12.032
56. Hurley ET, Mojica ES, Haskel JD, et al. Return to play testing following anterior cruciate reconstruction – A systematic review & meta-analysis. *The Knee*. 2022;34:134-140. doi:10.1016/j.knee.2021.11.010
57. Ardern CL, Glasgow P, Schneiders A, et al. 2016 consensus statement on return to sport from the first world congress in sports physical therapy, Bern. *Br J Sports Med*. 2016;50(14):853-864. doi:10.1136/bjsports-2016-096278
58. Burgi CR, Peters S, Ardern CL, et al. Which criteria are used to clear patients to return to sport after primary ACL reconstruction? A scoping review. *Br J Sports Med*. 2019;53(18):1154-1161. doi:10.1136/bjsports-2018-099982
59. Unverzagt C, Andreyo E, Tompkins J. ACL return to sport testing: it's time to step up our game. *Int J Sports Phys Ther*. 2021;16(4):1169-1177. doi:10.26603/001c.25463
60. Wilk K, Thomas ZM, Arrigo CA, Davies GJ. The need to change return to play testing in athletes following ACL injury: A theoretical model. *Int J Sports Phys Ther*. 2023;18(1). doi:10.26603/001c.67988
61. Meredith SJ, Rauer T, Chmielewski TL, et al. Return to sport after anterior cruciate ligament injury: Panther Symposium ACL Injury Return to Sport Consensus Group. *Journal of ISAKOS*. 2021;6(3):138-146. doi:10.1136/jisakos-2020-000495
62. Hanzlíková I, Hébert-Losier K. Is the landing error scoring system reliable and valid? A systematic review. *Sports Health*. 2020;12(2):181-188. doi:10.1177/1941738119886593
63. Jones P, Donelon T, Dos'Santos T. A preliminary investigation into a qualitative assessment tool to identify athletes with high knee abduction moments during cutting: Cutting Movement Assessment Score (CMAS). *Prof Strength Condit*. Published online 2017:37-42.
64. Dos'Santos T, McBurnie A, Donelon T, Thomas C, Comfort P, Jones PA. A qualitative screening tool to identify athletes with 'high-risk' movement mechanics during cutting: The cutting movement assessment score (CMAS). *Phys Ther Sport*. 2019;38:152-161. doi:10.1016/j.ptsp.2019.05.004
65. Della Villa F, Di Paolo S, Santagati D, et al. A 2D video-analysis scoring system of 90° change of direction technique identifies football players with high knee abduction moment. *Knee Surg Sports Traumatol Arthrosc*. 2021;30(11):3616-3625. doi:10.1007/s00167-021-06571-2
66. McLean SG. Evaluation of a two dimensional analysis method as a screening and evaluation tool for anterior cruciate ligament injury. *Br J Sports Med*. 2005;39(6):355-362. doi:10.1136/bjsm.2005.018598
67. Ressman J, Grooten WJA, Rasmussen Barr E. Visual assessment of movement quality in the single leg squat test: a review and meta-analysis of inter-rater and intrarater reliability. *BMJ Open Sport Exerc Med*. 2019;5(1):e000541. doi:10.1136/bmjsem-2019-000541
68. Almangoush A, Herrington L, Jones R. A preliminary reliability study of a qualitative scoring system of limb alignment during single leg squat. *Phys Ther Rehabil*. 2014;1(1):2. doi:10.7243/2055-2386-1-2
69. Numata H, Nakase J, Kitaoka K, et al. Two-dimensional motion analysis of dynamic knee valgus identifies female high school athletes at risk of non-contact anterior cruciate ligament injury. *Knee Surg Sports Traumatol Arthrosc*. 2018;26(2):442-447. doi:10.1007/s00167-017-4681-9
70. Padua DA, DiStefano LJ, Beutler AI, de la Motte SJ, DiStefano MJ, Marshall SW. The landing error scoring system as a screening tool for an anterior cruciate ligament injury-prevention program in elite-youth soccer athletes. *J Athl Train*. 2015;50(6):589-595. doi:10.4085/1062-6050-50.1.10
71. Fort-Vanmeerhaeghe A, Montalvo AM, Lloyd RS, Read P, Myer GD. Intra- and inter-rater reliability of the modified tuck jump assessment. *J Sports Sci Med*. 2017;16:117-124.
72. Myer GD, Ford KR, Hewett TE. Tuck jump assessment for reducing anterior cruciate ligament injury risk. Hubbard TJ, ed. *Athl Ther Today*. 2008;13(5):39-44. doi:10.1123/att.13.5.39
73. Hannon J, Wang-Price S, Swank C, et al. The validity and reliability of the Vail Sport Test™ as a measure of performance following anterior cruciate ligament reconstruction. *Phys Ther Sport*. 2019;38:162-169. doi:10.1016/j.ptsp.2019.05.001
74. Garrison JC, Shanley E, Thigpen C, Geary R, Osler M, DelGiorno J. The reliability of the vail sport test as a measure of physical performance following anterior cruciate ligament reconstruction. *Int J Sports Phys Ther*. 2012;7(1):11.

75. Kimura Y, Ishibashi Y, Tsuda E, Yamamoto Y, Tsukada H, Toh S. Mechanisms for anterior cruciate ligament injuries in badminton. *Br J Sports Med.* 2010;44(15):1124-1127. doi:10.1136/bjism.2010.074153
76. Chinnasee C, Weir G, Sasimontongkul S, Alderson J, Donnelly C. A biomechanical comparison of single-leg landing and unplanned sidestepping. *Int J Sports Med.* 2018;39. doi:10.1055/a-0592-7422
77. Jones PA, Herrington LC, Munro AG, Graham-Smith P. Is there a relationship between landing, cutting, and pivoting tasks in terms of the characteristics of dynamic valgus? *Am J Sports Med.* 2014;42(9):2095-2102. doi:10.1177/0363546514539446
78. Kristianslund E, Krosshaug T. Comparison of drop jumps and sport-specific sidestep cutting: implications for anterior cruciate ligament injury risk screening. *Am J Sports Med.* 2013;41(3):684-688. doi:10.1177/0363546512472043
79. Cowley HR, Ford KR, Myer GD, Kernozek TW, Hewett TE. Differences in neuromuscular strategies between landing and cutting tasks in female basketball and soccer athletes. *J Athl Train.* 2006;41(1):7.
80. King E, Richter C, Franklyn-Miller A, et al. Biomechanical but not timed performance asymmetries persist between limbs 9 months after ACL reconstruction during planned and unplanned change of direction. *J Biomech.* 2018;81:93-103. doi:10.1016/j.jbiomech.2018.09.021
81. Abrams GD, Harris JD, Gupta AK, et al. Functional performance testing after anterior cruciate ligament reconstruction: A systematic review. *Orthop J Sports Med.* 2014;2(1):232596711351830. doi:10.1177/2325967113518305
82. Sheppard J, Dawes J, Jeffreys I, Spiteri T, Nimphius S. Broadening the view of agility: a scientific review of the literature. *J Australian Strength Condit.* 2014;22(3):6-25.
83. Haff G, Triplett T. *Essentials of Strength Training and Conditioning.* 4th Edition. National Strength and Conditioning Association; 2016.
84. Dos'Santos T, McBurnie A, Thomas C, Jones PA, Harper D. Attacking agility actions: match play contextual applications with coaching and technique guidelines. *Strength Condit J.* 2022;44(5):102-118. doi:10.1519/SSC.0000000000000697
85. Welling W, Frik L. On-field tests for patients after anterior cruciate ligament reconstruction: a scoping review. *Orthop J Sports Med.* 2022;10(1):232596712110554. doi:10.1177/23259671211055481
86. Nimphius S, Callaghan SJ, Bezodis NE, Lockie RG. Change of direction and agility tests: challenging our current measures of performance. *Strength Condit J.* 2017;40(1):26-38. doi:10.1519/SSC.0000000000000309
87. Butler LS, Milian EK, DeVerna A, et al. Reliability of the cutting alignment scoring tool (CAST) to assess trunk and limb alignment during a 45-degree side-step cut. *Int J Sports Phys Ther.* Published online April 1, 2021. doi:10.26603/001c.21419
88. Butler LS, Martinez AR, Sugimoto D, et al. Reliability of the Expanded Cutting Alignment Scoring Tool (E-CAST) to assess trunk and limb alignment during a 45-degree side-step cut. *Int J Sports Phys Ther.* 2022;17(3). doi:10.26603/001c.33045
89. Weir G, Alderson J, Smailes N, Elliott B, Donnelly C. A reliable video-based ACL injury screening tool for female team sport athletes. *Int J Sports Med.* 2019;40(03):191-199. doi:10.1055/a-0756-9659
90. Needham C, Herrington L. Cutting movement assessment scores during anticipated and unanticipated 90-degree sidestep cutting manoeuvres within female professional footballers. *Sports.* 2022;10(9):128. doi:10.3390/sports10090128
91. Butler L, Martinez A, Erdman A, et al. Concurrent validity of the expanded cutting alignment scoring tool (E-CAST). *Int J Sports Phys Ther.* 2023;18(5). doi:10.26603/001c.87633
92. Butler L, Wyatt C, Martinez A, et al. No difference in two-dimensional kinematic assessment of a 45-degree sidestep cut compared to qualitative assessment. *Int J Sports Phys Ther.* 2023;18(3). doi:10.26603/001c.74366
93. Della Villa F, Di Paolo S, Santagati D, et al. A 2D video-analysis scoring system of 90° change of direction technique identifies football players with high knee abduction moment. *Knee Surg Sports Traumatol Arthrosc.* 2021;30(11):3616-3625. doi:10.1007/s00167-021-06571-2
94. Jones PA, Rai A, Dos'Santos T, Herrington LC. Inter-professional and methodological agreement in using the cutting movement assessment score (CMAS). *Biomech.* 2023;3(2):181-192. doi:10.3390/biomechanics3020016

95. Aparicio-Sarmiento A, Hernández-García R, Cejudo A, Palao JM, Sainz de Baranda P. Reliability of a qualitative instrument to assess high-risk mechanisms during a 90° change of direction in female football players. *IJERPH*. 2022;19(7):4143. doi:10.3390/ijerph19074143
96. Butler LS, Martinez AR, Sugimoto D, et al. Reliability of the expanded cutting alignment scoring tool (E-CAST) to assess trunk and limb alignment during a 45-degree side-step cut. *Intl J Sports Phys Ther*. 2022;17(3). doi:10.26603/001c.33045
97. Olivares-Jabalera J, Fíler-Ruger A, Dos'Santos T, et al. Is there association between cutting and jump-landing movement quality in semi-professional football players? Implications for ACL injury risk screening. *Phys Ther Sport*. 2022;56:15-23. doi:10.1016/j.ptsp.2022.05.015
98. Webster KE, Hewett TE. What is the Evidence for and Validity of Return-to-Sport Testing after Anterior Cruciate Ligament Reconstruction Surgery? A Systematic Review and Meta-Analysis. *Sports Medicine*. 2019;49(6):917-929. doi:10.1007/s40279-019-01093-x
99. Manoj Rajakaruna P, Dempsey AR, Learmonth YC. Reliability, validity, and predictability of screening methods for noncontact anterior cruciate ligament injury- A systematic review. *J Sci Med Sport*. 2023;26:S110-S111. doi:10.1016/j.jsams.2023.08.136
100. Fältström A, Hägglund M, Hedevik H, Kvist J. Poor validity of functional performance tests to predict knee injury in female soccer players with or without anterior cruciate ligament reconstruction. *Am J Sports Med*. 2021;49(6):1441-1450. doi:10.1177/03635465211002541
101. Van Melick N, Pronk Y, Nijhuis-van Der Sanden M, Rutten S, Van Tienen T, Hoogeboom T. Meeting movement quantity or quality return to sport criteria is associated with reduced second ACL injury rate. *J Orthop Res*. 2022;40(1):117-128. doi:10.1002/jor.25017
102. Capin JJ, Snyder-Mackler L, Risberg MA, Grindem H. Keep calm and carry on testing: A substantive reanalysis and critique of 'what is the evidence for and validity of return-to-sport testing after anterior cruciate ligament reconstruction surgery? A systematic review and meta-analysis. *Br J Sports Med*. 2019;53(23):1444-1446. doi:10.1136/bjsports-2019-100906
103. Marques JB, Paul DJ, Graham-Smith P, Read PJ. Change of direction assessment following anterior cruciate ligament reconstruction: a review of current practice and considerations to enhance practical application. *Sports Med*. 2020;50(1):55-72. doi:10.1007/s40279-019-01189-4
104. Piskin D, Benjaminse A, Dimitrakis P, Gokeler A. Neurocognitive and neurophysiological functions related to ACL injury: A framework for neurocognitive approaches in rehabilitation and return-to-sports tests. *Sports Health*. Published online July 8, 2021:194173812110292. doi:10.1177/19417381211029265
105. Grooms DR, Chaudhari A, Page SJ, Nichols-Larsen DS, Onate JA. Visual-motor control of drop landing after anterior cruciate ligament reconstruction. *J Athl Train*. 2018;53(5):486-496. doi:10.4085/1062-6050-178-16
106. Gokeler A, Neuhaus D, Benjaminse A, Grooms DR, Baumeister J. Principles of motor learning to support neuroplasticity after ACL injury: implications for optimizing performance and reducing risk of second ACL injury. *Sports Med*. 2019;49(6):853-865. doi:10.1007/s40279-019-01058-0
107. Faltus J, Criss CR, Grooms DR. Shifting focus: A clinician's guide to understanding neuroplasticity for anterior cruciate ligament rehabilitation. *American College of Sports Medicine*. 2020;19(2):8. doi:10.1249/ISR.0000000000000688
108. Grooms DR, Chaput M, Simon JE, Criss CR, Myer GD, Diekfuss JA. Combining neurocognitive and functional tests to improve return to sport decisions following ACL reconstruction. *J Orthop Sports Phys Ther*. Published online May 15, 2023:1-14. doi:10.2519/jospt.2023.11489
109. Serpell BG, Young WB, Ford M. Are the perceptual and decision-making components of agility trainable? A preliminary investigation. *J Strength Condit Res*. 2011;25(5):1240-1248. doi:10.1519/JSC.0b013e3181d682e6
110. Safaric A, Bird S. Agility drills for basketball: Review and practical applications. *J Australian Strength Condit*. 2011;19:24-32.
111. Brown SR, Brughelli M, Hume PA. Knee mechanics during planned and unplanned sidestepping: a systematic review and meta-analysis. *Sports Med*. 2014;44(11):1573-1588. doi:10.1007/s40279-014-0225-3
112. Simonek J, Horicka P, Hianik J. Differences in pre-planned agility and reactive agility performance in sport games. *Acta Gymnica*. 2016;46(2):68-73. doi:10.5507/ag.2016.006
113. Besier TF, Lloyd DG, Ackland TR, Cochrane JL. Anticipatory effects on knee joint loading during running and cutting maneuvers. *Med Sci Sports Exerc*. Published online July 2001:1176-1181. doi:10.1097/00005768-200107000-00015

114. Fuerst P, Gollhofer A, Gehring D. Preparation time influences ankle and knee joint control during dynamic change of direction movements. *J Sports Sci.* 2017;35(8):762-768. doi:10.1080/02640414.2016.1189084
115. Mornieux G, Gehring D, Fürst P, Gollhofer A. Anticipatory postural adjustments during cutting manoeuvres in football and their consequences for knee injury risk. *J Sports Sci.* 2014;32(13):1255-1262. doi:10.1080/02640414.2013.876508
116. Hughes G, Dai B. The influence of decision making and divided attention on lower limb biomechanics associated with anterior cruciate ligament injury: a narrative review. *Sports Biomech.* 2023;22(1):30-45. doi:10.1080/14763141.2021.1898671
117. Dos'Santos T, Thomas C, Comfort P, Jones PA. The effect of angle and velocity on change of direction biomechanics: an angle-velocity trade-off. *Sports Med.* 2018;48(10):2235-2253. doi:10.1007/s40279-018-0968-3
118. Kristianslund E, Faul O, Bahr R, Myklebust G, Krosshaug T. Sidestep cutting technique and knee abduction loading: implications for ACL prevention exercises. *Br J Sports Med.* 2014;48(9):779-783. doi:10.1136/bjsports-2012-091370
119. Kimura K, Sakurai S. A sidestep cut preparation strategy decreases the external load applied to the knee joint. *Int J Sport Health Sci.* 2013;11(0):109-117. doi:10.5432/ijshs.201309
120. Nedergaard NJ, Kersting U, Lake M. Using accelerometry to quantify deceleration during a high-intensity soccer turning manoeuvre. *J Sports Sci.* 2014;32(20):1897-1905. doi:10.1080/02640414.2014.965190
121. Vanrenterghem J, Venables E, Pataky T, Robinson MA. The effect of running speed on knee mechanical loading in females during side cutting. *J Biomech.* 2012;45(14):2444-2449. doi:10.1016/j.jbiomech.2012.06.029
122. Nagelli CV, Wordeman SC, Di Stasi S, Hoffman J, Marulli T, Hewett TE. Neuromuscular Training Improves Biomechanical Deficits at the Knee in Anterior Cruciate Ligament-Reconstructed Athletes. *Clin J Sport Med.* Published online 2019. doi:10.1097/ISM.0000000000000723
123. Arumugam A, Björklund M, Mikko S, Häger CK. Effects of neuromuscular training on knee proprioception in individuals with anterior cruciate ligament injury: a systematic review and GRADE evidence synthesis. *BMJ Open.* 2021;11(5):e049226. doi:10.1136/bmjopen-2021-049226
124. Buckthorpe M. Recommendations for movement re-training after ACL reconstruction. *Sports Med.* 2021;51(8):1601-1618. doi:10.1007/s40279-021-01454-5
125. Olivares-Jabalera J, Fíler A, Dos'Santos T, Ortega-Domínguez J, Soto Hermoso VM, Requena B. The safe landing warm up technique modification programme: an effective anterior cruciate ligament injury mitigation strategy to improve cutting and jump-movement quality in soccer players. *J Sports Sci.* 2022;40(24):2784-2794. doi:10.1080/02640414.2023.2193451
126. Dos'Santos T, McBurnie A, Comfort P, Jones PA. The effects of six-weeks change of direction speed and technique modification training on cutting performance and movement quality in male youth soccer players. *Sports.* 2019;7(9):205. doi:10.3390/sports7090205
127. Nijmeijer EM, Elferink-Gemser MT, McCrory S, Cortes N, Benjaminse A. How to improve movement execution in sidestep cutting? Involve me and I will learn. *Hum Mov Sci.* 2023;90:103115. doi:10.1016/j.humov.2023.103115
128. Dos'Santos T, Thomas C, McBurnie A, Donelon T, Herrington L, Jones PA. The Cutting Movement Assessment Score (CMAS) Qualitative Screening Tool: Application to mitigate anterior cruciate ligament injury risk during cutting. *Biomech.* 2021;1(1):83-101. doi:10.3390/biomechanics1010007

SUPPLEMENTARY MATERIALS

Appendix 1

Download: <https://ijspt.scholasticahq.com/article/123483-clinical-utility-of-qualitative-change-of-direction-movement-assessment-in-acl-injury-risk-evaluation/attachment/245157.docx>

Appendix 2

Download: <https://ijspt.scholasticahq.com/article/123483-clinical-utility-of-qualitative-change-of-direction-movement-assessment-in-acl-injury-risk-evaluation/attachment/245156.docx>
