




**Please cite the Published Version**

Keightley, Anna T , Field, Chris D , Rowson, James G and Caporn, Simon J M  (2024) Comparative Photosynthetic Capacity, Respiration Rates, and Nutrient Content of Micropropagated and Wild-Sourced Sphagnum. *International Journal of Plant Biology*, 15 (4). pp. 959-978. ISSN 2037-0164

**DOI:** <https://doi.org/10.3390/ijpb15040068>

**Publisher:** MDPI

**Version:** Published Version

**Downloaded from:** <https://e-space.mmu.ac.uk/636120/>

**Usage rights:**  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

**Additional Information:** This is an open access article which first appeared in *International Journal of Plant Biology*, published by MDPI

**Data Access Statement:** The original data presented in the study are openly available in Mendeley Data at DOI:10.17632/dfvmfmb3vz.1. <https://data.mendeley.com/datasets/dfvmfmb3vz/1> (accessed on 24 April 2024).

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

Article

# Comparative Photosynthetic Capacity, Respiration Rates, and Nutrient Content of Micropropagated and Wild-Sourced *Sphagnum*

Anna T. Keightley <sup>1,\*</sup> , Chris D. Field <sup>1</sup>, James G. Rowson <sup>2</sup> and Simon J. M. Caporn <sup>1</sup> 

<sup>1</sup> Department of Natural Sciences, Manchester Metropolitan University, Ormond Building, Lower Ormond Street, Manchester M15 6BX, UK; c.field@mmu.ac.uk (C.D.F.); s.j.m.caporn@mmu.ac.uk (S.J.M.C.)

<sup>2</sup> Jacobs UK Limited, 5 First Street, Manchester M15 4GU, UK

\* Correspondence: anna.keightley@mmu.ac.uk

**Abstract:** The rapid, effective restoration of degraded peatlands is urgently needed to reduce their current high levels of carbon loss. The re-introduction of *Sphagnum* moss, along with re-wetting, is key to returning carbon sequestration and retention capabilities to northern degraded bogs. Micropropagated *Sphagnum* has already been applied in large quantities, and more is planned, for restoration projects in Britain and parts of Europe. A comparison with wild-sourced *Sphagnum* material is therefore pertinent to demonstrate its safety and suitability for wide-scale application. Six *Sphagnum* species of both micropropagated and wild-sourced origin were assessed for photosynthetic capacity, nutrient content, form parity, chlorocyst size, and chloroplast numbers. Micropropagated *Sphagnum* had significantly higher light-saturated photosynthesis ( $P_{max}$ ) rates, little color expression, an open growth habit, greater chloroplast numbers, and more numerous, smaller shoot apices than wild-sourced *Sphagnum*. Higher  $P_{max}$  rates were associated with a lower bulk density and higher tissue nutrient concentrations. Potentially, greater chloroplast numbers in micropropagated *Sphagnum* facilitate higher photosynthesis rates, driving rapid growth in early-stage plants, particularly in optimum moisture conditions. Micropropagated *Sphagnum* can be used confidently, propagated in large quantities, and will likely establish well on application to sites where re-wetting has already occurred, therefore making it highly beneficial for the restoration of degraded bogs.

**Keywords:** *Sphagnum*; peatlands; restoration; carbon; physiology; chloroplasts



**Citation:** Keightley, A.T.; Field, C.D.; Rowson, J.G.; Caporn, S.J.M. Comparative Photosynthetic Capacity, Respiration Rates, and Nutrient Content of Micropropagated and Wild-Sourced *Sphagnum*. *Int. J. Plant Biol.* **2024**, *15*, 959–978. <https://doi.org/10.3390/ijpb15040068>

Academic Editor: Dennis H. Greer

Received: 11 September 2024

Revised: 25 September 2024

Accepted: 30 September 2024

Published: 2 October 2024



**Copyright:** © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

## 1. Introduction

*Sphagnum* is instrumental in its role as an ecosystem engineer in creating the cool, wet, acidic, low-nutrient conditions leading to the formation of peat in northern peatbogs, through chemical processes and recalcitrant plant tissues [1–3]. The uptake by *Sphagnum* of nutrients and water from rain and cloud water are well established in raised and blanket bogs [2]. *Sphagnum* absorbs aerially deposited nutrients rapidly and directly into plant tissue [4,5] and stores them efficiently, reducing their availability to the roots of vascular plants [6,7]. Moreover, the accumulation of carbohydrates occurs in environments that restrict plant growth and are stored as carbon, and those in *Sphagnum*, especially uronic acids (sphagnan), also form polymers in cell walls, responsible for ionic exchange, and are reckoned by Clymo [8] to make up 10–30% of fresh *Sphagnum* dry weight. Cation exchange between  $H^+$  and  $Ca^{2+}$  and  $Mg^{2+}$  especially, but also  $K^+$  and  $NH_4^+$ , traps nutrients for *Sphagnum* growth and creates an acid, nutrient-poor environment, most efficiently in the upper acrotelm [1,2]. Different species have also developed a range of adaptive or phylogenetic traits to specific ecological conditions of light, shade, and moisture [9–12], allowing them to outcompete vascular plants in the resulting hostile bog environment [2,7].

Intact peatlands store more atmospheric carbon per hectare than other terrestrial habitats [13,14]. However, many peatbogs are degraded through drainage and subsequent

land-use change. Greenhouse gas (GHG) emissions from degraded peatlands are substantial; for example, in the UK they were estimated at approximately 23.1 Mt CO<sub>2</sub>e yr<sup>-1</sup> by Evans et al. [15], which was 5% of the total GHG emissions for the UK in 2017 [16]. Restoration is vital to reduce this contribution to anthropogenic climate change [17,18]. *Sphagnum* is a key species for bog restoration [19], but the quantities available for re-introduction are insufficient from wild sources in many regions, including the UK, and sites are often protected [20]. Micropropagated *Sphagnum* (described below) has been developed to fill that gap in resources and has already proved successful on wide-scale introduction to blanket bogs in the north of England [21,22]. There is an urgent requirement to examine and understand the performance of micropropagated *Sphagnum* because of the scale of current and future peatland restoration needs.

The *Sphagnum* genus occupies a wide range of peatland ecological niches, and *Sphagnum* species vary widely in their photosynthetic rates, yet the plant and environmental factors driving production and carbon accumulation remain unclear [23]. Loisel et al. [24], in a wide-ranging review of global measurements, found *Sphagnum* stem-length growth and photosynthetically active radiation (PAR) were strongly correlated, with PAR a more important growth indicator than moisture levels, albeit for only two species assessed, *S. magellanicum* (probably *S. medium* or *S. divinum* [25]) and *S. fuscum*. However, studying the same two species, Bengtsson et al. [23] found that PAR had only a low annual effect on length increment, and that a range of other factors were more important, although the effects were species-specific. These included moisture levels, temperature, nitrogen deposition, and vascular plant cover, with the latter no doubt influencing the PAR levels available to *Sphagnum*. Photosynthesis is constrained by moisture levels in bryophytes as the plants are poikilohydric: too much moisture limits CO<sub>2</sub> diffusion and reduces carboxylation, and too low moisture damages photosynthetic apparatus [26]. Low levels of nutrients, particularly nitrogen, support photosynthesis in mosses, although higher levels can be toxic and promote shading from vascular plants [27,28]. *Sphagnum* utilize the same nutrient elements that all plants use for photosynthesis, respiration, and growth, but absorb them directly into cells [4] as they have no vascular transportation system for uptake from the soil, and allocate them differently, as nutrient resources are limited in an ombrotrophic bog system [9,29].

Photosynthesis rates generally decline following the ‘successional gradient’ of species in bog development towards ombrotrophic conditions [12]. Species with metabolic strategies, such as high bulk density and carotenoid concentration to tolerate drier, unshaded conditions, tend to have reduced rates of growth and photosynthesis [9], and shade-adapted species tend to have high photosynthesis rates [30]. Moreover, Hájek et al. [10], in laboratory conditions and under a range of light intensities, found a clear ranking of CO<sub>2</sub> uptake between species, with those sourced from shaded habitats tending to rank higher than those sourced from open habitats. The authors surmised that species from open habitats suffered persistent photodamage, which reduced photosynthetic capacity, despite photoprotective pigments such as sphagnorubin. Additionally, greater numbers of chloroplasts in plants is known to be associated with a greater photosynthetic capacity [31].

Micropropagated *Sphagnum* has been produced in bulk quantities in the UK from a few shoots of wild-sourced material sourced from bogs in the north of England, primarily the Peak District and Cumbria. Production involves standard techniques of plant division, whereby the *Sphagnum* is surface-sterilized and tissue-cultured under aseptic conditions [32]. It is then grown on under greenhouse conditions with minimal, bespoke nutrient application [20]. Studying micropropagated *Sphagnum* is an opportunity for novel comparisons of species that have been cultured and grown under optimum light, moisture, and nutrient levels and are at the same stage of development.

Studies into the photosynthetic capacity of different species of *Sphagnum* moss from contrasting sources are pertinent to understanding production and, therefore, carbon sequestration in bogs [24]. Respiration is part of the net CO<sub>2</sub> exchange, and so is also of study interest. The aims of this study were to make a comparison of photosynthesis and

respiration rates and examine any differences in chlorocyst (cell containing chloroplasts) size (by observation, size appeared to vary between the same species of micropropagated and wild-sourced *Sphagnum*), number of chloroplasts and nutrient content of six *Sphagnum* species of both tissue-cultured (micropropagated) and wild-sourced plants. A greater understanding of the potential carbon sequestration of each micropropagated species will help direct both product development and restoration efforts where these products are used.

The objectives were, firstly, to measure the CO<sub>2</sub> uptake (photosynthesis) and emission (respiration) of samples of *S. capillifolium*, *S. fallax*, *S. medium/divinum*, *S. palustre*, *S. papillosum*, and *S. squarrosum* in controlled conditions over a range of light intensities. These represent species from a broad environmental range and were readily available. Wild-sourced samples were taken from established, naturally occurring colonies in a range of peatland environments. Secondly, micropropagated and wild-sourced samples of the same six species used for photosynthesis rate studies were examined under a microscope and measurements made of chlorocyst size and number of chloroplasts to test for any differences that may influence capacity for photosynthesis. Thirdly, the nutritional content of samples used for photosynthesis measurements were analyzed to examine whether the levels of nutrients within tissues of micropropagated *Sphagnum*, which is a horticultural product, and wild-sourced *Sphagnum* had a bearing on their photosynthetic capacity.

Our research questions, which we succeeded in addressing, were:

1. Which *Sphagnum* species, either grown under the same conditions (micropropagated) or grown in the wild, show the greatest rates of photosynthesis and respiration?
2. Is there a difference in photosynthetic capacity and respiration rate between micropropagated and wild-sourced *Sphagnum* species?
3. Are there differences in chlorocyst size, chloroplast number, and nutrient content between micropropagated and wild-sourced *Sphagnum* species that may explain differences in their photosynthetic capacity and respiration rate?

## 2. Materials and Methods

### 2.1. *Sphagnum* Photosynthesis and Respiration

Approximately 2 liters of each wild-sourced *Sphagnum* species were collected in August 2017 from ombrotrophic mires and heaths in the north of England and Wales: *S. capillifolium*, *S. fallax*, and *S. palustre* from Chat Moss Remnants Site of Biological Importance (SBI) (now owned by Lancashire Wildlife Trust, and called 'Rindle Moss') (53°27'53.0" N, 2°26'56.4" W), *S. medium/divinum* from Borth Bog (Cors Fochno) (52°30'18.7" N, 4°00'43.5" W), and *S. papillosum* from Ruabon Moor (53°00'11.5" N, 3°08'21.6" W). *S. squarrosum* was collected from Alderley Edge (53°17'52.1" N, 2°12'18.0" W), a wooded basin mire.

Micropropagated samples of the same species were also collected in June–July 2017. Micropropagated *Sphagnum* material is cultivated in glasshouses, where shade is applied in summer and humidity and temperature conditions are controlled, with minimal, bespoke (commercially sensitive information) nutrient application. The samples used were grown from micropropagated *Sphagnum* suspended in a hydrocolloidal gel and applied directly onto the growing-medium surface. The species selected were typical of the micropropagated *Sphagnum* grown for restoration work.


All *Sphagnum* samples were acclimatized in a Fitotron growth cabinet (Weiss Technik, Lindenstruth-Reiskirchen) for a minimum of 5 days, set to typical summer-time environmental conditions in the local area: 20 °C during the day (0600–2200 h), 12 °C during the night, day-time light intensity of 750–800 µmol (photons) m<sup>-2</sup> s<sup>-1</sup> (values calculated from the nearby Astley Moss Weather Station mean data, 2012–2015), and 85% humidity. The *Sphagnum* was misted with rainwater as necessary to keep it hydrated.

A literature review was used to determine the range of light levels needed to capture the photosynthetic response of diverse *Sphagnum* species. Haraguchi and Yamada [33] found that optimum light levels for photosynthesis for a range of *Sphagnum* species were between 300 and 500 µmol (photons) m<sup>-2</sup> s<sup>-1</sup> of PPFD (Photosynthetic Photon Flux Den-



sity). However, Rice et al. [9] found that photoinhibition (to prevent high-light damage) in *Sphagnum* occurs at  $800 \mu\text{mol (photons) m}^{-2} \text{s}^{-1}$  and Loisel et al. [24] reported an optimum from 500 to  $900 \mu\text{mol (photons) m}^{-2} \text{s}^{-1}$ . Hájek et al. [10] found that the light saturation point for all *Sphagnum* they studied was similar at an average of  $2124 \pm 86$  (SE)  $\mu\text{mol (photons) m}^{-2} \text{s}^{-1}$ , much higher than those of other studies. In this study,  $800 \mu\text{mol (photons) m}^{-2} \text{s}^{-1}$  was the highest light level used.

A clear acrylic cylinder measuring 5 cm diameter  $\times$  3 cm high, fitted with a mesh base to allow air circulation through the plants (Figure 1), was used to hold each *Sphagnum* sample. Five replicate samples of each species from each type (micropropagated and wild-sourced) were cut to 3 cm lengths from the 2 L bulk amounts of *Sphagnum* at growing shoot density and placed in the cylinder (i.e., not a standard number of shoots, or compressed to fit) with the top surface of the sample level with the top of the cylinder. The number of shoots per sample varied by species and type and ranged from 19 to 96 (micropropagated) and from 13 to 73 (wild-sourced). An LGR™, Ultraportable Greenhouse Gas Analyzer (UGGA), Model 915-0011 (Los Gatos, Research, Palo Alto, CA, USA) (LGR), was fitted to a 500 mL sealable clear-glass chamber via tubing through air-tight ports in the lid, and *Sphagnum* samples placed in the chamber for analysis. Change in  $\text{CO}_2$  concentration within the chamber was measured over 2 min and a light response curve determined for each species starting from light intensities of  $800 \mu\text{mol (photons) m}^{-2} \text{s}^{-1}$  to zero in increments of  $50 \mu\text{mol (photons) m}^{-2} \text{s}^{-1}$  for each sample. A Skye PAR quantum sensor, integral to the growth cabinet, was placed on a level with the sample being assessed to set an accurate light intensity. The reduction in light transmission through the clear-glass chamber (8%) was accounted for by increasing the light intensity in the cabinet accordingly for each light intensity measurement. Samples were lightly misted with rainwater after each measurement and acclimated to each change in light level between measurements, after the light intensity had stabilized (similar to methods used by Kangas et al. [34]), for approximately 5 min.



**Figure 1.** Examples of *Sphagnum* species samples for the analysis of the photosynthesis rate. Micropropagated (top row of pair) and wild-sourced *Sphagnum* species samples show typical visual differences in pigmentation and form.

The samples were photographed prior to measurements (Figure 1) and the number of capitula per sample counted. Photography also allowed for the qualitative comparison of types, e.g., color expression. Samples were weighed to check for water loss between measurements, and the chamber was closed between each light level measurement to reduce drying. The LGR flow rate was  $0.8 \text{ L min}^{-1}$  with space between gas inlet and outlet points; air was released into the chamber along a small pipe with holes along the length to encourage mixing (LGR low flow rate threshold for analysis is  $\sim 0.35 \text{ L min}^{-1}$  [personal communication, Lewis John, LGR™ Sales Representative]).

Net photosynthesis or respiration rate was calculated through Microsoft Excel from the rate of  $\text{CO}_2$  depletion or increase using linear regression over a two-minute period, and further expressed by the surface area of the plant chamber ( $A_s$ ) and subsequently by the total plant dry weight (DW). Values are expressed using the leaf gas exchange sign

convention, whereby plant uptake of CO<sub>2</sub> from the atmosphere is expressed as positive and loss to the atmosphere is expressed as negative. The calculation to determine the photosynthesis (CO<sub>2</sub> uptake) or respiration (CO<sub>2</sub> emission) rate (adapted from Dossa et al. [35]) is:

$$\text{Photosynthesis or Respiration} = \frac{\Delta\text{CO}_2}{t} \times \frac{PV}{RT} \times \frac{1}{A_s} \times \left( \frac{44 \times 60 \times 60}{1000} \right) \quad (1)$$

(P (atm) = atmospheric pressure; V (m<sup>3</sup>) = chamber volume; R (L atm mol<sup>-1</sup> K) = universal gas constant; T (K) = gas temperature in Kelvin; A<sub>s</sub> (m<sup>2</sup>) = sample surface area; 44 g mol<sup>-1</sup> = molecular weight of CO<sub>2</sub>); Photosynthesis or Respiration = g CO<sub>2</sub> m<sup>-2</sup> h<sup>-1</sup>.

## 2.2. *Sphagnum* Samples' Bulk Density

High shoot density and bulk density are physiological aspects of *Sphagnum*'s protective adaptation to open conditions and may lead to reduced photosynthesis rates; so, bulk density measurements were made to quantify this. After photosynthesis measurements, samples were dried overnight at 105 °C (temperature used by other researchers, e.g., Limpens and Berendse [36], and McNeil and Waddington [37]) to obtain dry weight for calculations and for nutrient analysis. Dry weight bulk density was calculated from the dry weight divided by the known sample volume.

## 2.3. *Sphagnum* Samples' Nutrient Analysis

The nitrogen content of the samples was obtained using a LECO FP628 elemental analyzer. A minimum of 0.05 g of dry *Sphagnum* was used per sample. Each sample measured for photosynthesis (*n* = 60) was analyzed, plus 4 replicates of one species from each type (micropropagated and wild-sourced) to determine the experimental error (total *n* = 68).


For other elements, samples (as above) were prepared for Inductively Coupled Plasma Optical Emission spectroscopy (ICP-OES) through acid and microwave digest, using HNO<sub>3</sub> S.G. 1.42 (>68%) PrimerPlus-Trace analysis grade in a CEM Mars Xpress 5 Microwave, and the diluted solution was analyzed through ICP-OES (Thermo Fisher Scientific, Waltham, MA, USA).

## 2.4. *Sphagnum* Cell Measurements and Analysis


In addition to *Sphagnum* measured for photosynthesis and nutrient analysis, small samples (several plant shoots) of *Sphagnum* for microscopic analysis were collected in September 2019 from unheated greenhouses (micropropagated—daytime temperature ~20 °C) and from natural sources and assessed over the following week. Wild-sourced *Sphagnum* was from sites in the north of England: *S. fallax*, *S. medium/divinum*, and *S. palustre* from Cadishead Moss (53°27'07.9" N, 2°27'18.9" W); *S. capillifolium* and *S. papillosum* from Astley Moss (53°28'32.2" N, 2°27'15.5" W); and *S. squarrosum* from Windy Bank Wood (53°28'15.3" N, 2°28'59.6" W). Leaves on divergent branches just below the capitula are generally used for the microscopic observation of *Sphagnum* [38,39]. The cell structure changes across the leaf from proximal to distal ends and from edge to center, and between convex and concave aspects [39,40]. For the standardization of the results in this study, leaves from branches just below and immediately surrounding the capitulum (as indicated in Figure 2) were observed and measurements made centrally, on the concave aspect.

Leaves from three branches from below three capitula of each sample (i.e., 9 leaves) were removed onto a slide and photographed using a Brunel Eyecam Plus attached to a compound microscope at 1000× magnification; cell dimensions were measured after calibration at the same magnification. There is a collection of five or six chlorocysts (cells containing chloroplasts) surrounding a hyalocyst (a dead, thin-walled, and hollow cell with a water storage function). The width (rather than length, which has a very wide variation) of all segments around one hyalocyst per leaf was measured centrally (Figure 3), and the number of chloroplasts counted in each. A mean value was calculated from each set of measurements for each leaf, and thus, there were 9 values for chlorocyst width and for

number of chloroplasts for each species. Only chloroplasts within chlorocysts immediately surrounding each single hyalocyst assessed were included in the measurements.



**Figure 2.** Examples of capitula used for microscopic study; arrows indicate typical branch selection.



**Figure 3.** Examples of chlorocyst width measurement locations (double-ended arrows) and chloroplasts (single-arrowed) (A) *S. palustre* and (B) *S. squarrosum*.

### 2.5. Data Analysis

Data were analyzed using IBM SPSS Statistics for Windows (Version 25.0. Armonk, NY, USA: IBM Corp.) and PAST: Paleontological Statistics Software Package for Education and Data Analysis [41] where indicated. Data were tested for normality using Shapiro–Wilk tests. Data for maximum photosynthesis ( $P_{\max}$ ), respiration rates, number of capitula per sample, fresh (FW) and dry weight (DW) bulk density, and also data for microscopic measurements of chlorocyst width and number of chloroplasts were found to be normally distributed, and so, parametric *t*-tests were used to test differences between *Sphagnum* types (micropropagated and wild-sourced), as well as one- and two-way ANOVAs with post-hoc




Tukey's HSD to test differences between species within types and between types for each species. Dependency of respiration rates on  $P_{\max}$  rates was tested using linear regression. Data for nutrients were not normally distributed, and so, non-parametric independent variable tests (Mann–Whitney U) were used to test differences in distribution between *Sphagnum* types (micropropagated and wild-sourced). Associations between nutrient levels,  $P_{\max}$  and respiration rates, and *Sphagnum* type and species were examined with a correlation matrix through a principal component analysis using the PAST software [41]. Statistical significance was determined with a  $p$ -value  $< 0.05$ .

### 3. Results

#### 3.1. *Sphagnum* Photosynthesis and Respiration

The response of net photosynthesis rate ( $P_n$ ) to changing light levels showed a similar pattern across all species in both micropropagated and wild-sourced samples (Figure 4), reaching a maximum  $P_n$  ( $P_{\max}$ ) between 400 and 650, and 400 and 750  $\mu\text{mol}$  (photons)  $\text{m}^{-2} \text{s}^{-1}$ , respectively (Table 1), either levelling or reducing thereafter. Significant inter-species differences were seen within both micropropagated and wild-sourced *Sphagnum* types for  $P_{\max}$  ( $F = 11.29$  and  $115.9$ , respectively;  $p < 0.001$ ,  $df = 5$  for both) and respiration rates ( $F = 12.0$  and  $39.15$ , respectively;  $p < 0.001$ ,  $df = 5$  for both). Significant differences between species are indicated on Figure 5.  $P_{\max}$  was significantly higher in micropropagated than in wild-sourced samples overall ( $t = 8.647$ ,  $p < 0.001$ ,  $df = 58$ ). The  $P_{\max}$  of each species was higher in micropropagated than wild-sourced types (significant differences between species from two-way ANOVA Tukey's post hoc tests are indicated in Table 1). The  $P_{\max}$  rates across micropropagated samples were less variable than wild-sourced samples (coefficient of variation of 32.2% and 74.7%, respectively). Respiration rates were significantly greater in micropropagated than in wild-sourced samples overall ( $t = 5.816$ ,  $p < 0.001$ ,  $df = 58$ ) and by species (not significantly for *S. fallax* or *S. palustre*).




**Figure 4.** Comparison of micropropagated and wild-sourced *Sphagnum* species net photosynthesis ( $P_n$ ) response to light. A light intensity range from 0 to 800  $\mu\text{mol}$  (photons)  $\text{m}^{-2} \text{s}^{-1}$  was used to test each sample. Results are shown on a dry weight basis ( $n = 5$ ). Crosses indicate the mean, lines indicate the median, and interquartile range is exclusive with maximum and minimum values that are not outliers indicated by the whiskers.



**Table 1.** Comparison of light-saturated photosynthesis (with associated PAR level) and respiration rates of *Sphagnum* species within each type.

<i>Sphagnum</i> Species	Micropropagated			Wild-Sourced		
	PAR	P <sub>max</sub>	Respiration	PAR	P <sub>max</sub>	Respiration
<i>S. squarrosum</i>	500	46.94 ± 4.74 **	−8.15 ± 1.77 *	400	25.32 ± 3.75	−6.05 ± 1.11
<i>S. palustre</i>	400	37.69 ± 11.58 **	−5.88 ± 0.66	450	15.23 ± 1.00	−4.15 ± 0.60
<i>S. papillosum</i>	550	36.28 ± 3.65 **	−7.03 ± 0.47 **	750	3.55 ± 1.27	−1.91 ± 0.49
<i>S. fallax</i>	450	32.54 ± 6.17 **	−5.42 ± 0.82	450	16.14 ± 1.25	−4.37 ± 0.72
<i>S. capillifolium</i>	650	24.12 ± 2.53 **	−4.97 ± 0.55 *	500	6.97 ± 1.64	−2.54 ± 0.31
<i>S. medium/divinum</i>	450	20.57 ± 5.51 **	−4.07 ± 0.78 **	550	1.59 ± 0.42	−1.17 ± 0.29

Light-saturated photosynthesis (P<sub>max</sub>) with associated photosynthetically active radiation (PAR μmol (photons) m<sup>−2</sup> s<sup>−1</sup>) level and respiration rates of samples, expressed by dry weight (nmol CO<sub>2</sub> g<sup>−1</sup> s<sup>−1</sup>) ordered from the highest to the lowest P<sub>max</sub> by micropropagated species, paired with wild-sourced equivalents. Leaf gas exchange sign convention used (i.e., CO<sub>2</sub> uptake positive and CO<sub>2</sub> emission negative). Values are mean (n = 5) ± SD. Significant differences (two-way ANOVA Tukey’s post hoc tests) between each pair are indicated: \* p < 0.05; \*\* p < 0.001.



**Figure 5.** Inter-species differences in light-saturated photosynthesis (P<sub>max</sub>) and respiration (Resp) rates within *Sphagnum* types. Statistically significant differences between species on each graph are indicated by shared letters. Crosses indicate the mean, lines indicate the median, and interquartile range is exclusive with maximum and minimum values that are not outliers indicated by the whiskers.

Micropropagated species are ranked from the highest to lowest P<sub>max</sub> in Table 1, but the ranking for wild-sourced species was ordered differently for both P<sub>max</sub> and respiration rates. However, the ranking of micropropagated and wild-sourced species with the highest (*S. squarrosum*) and lowest (*S. medium/divinum*) P<sub>max</sub> and respiration rates were the same.

The ratio of P<sub>max</sub> to respiration on a DW basis was consistently higher in micropropagated samples (5.58) than wild-sourced samples (3.41). However, there was a strong positive linear regression between P<sub>max</sub> and respiration rates in wild-sourced samples (R<sup>2</sup> = 0.90, p < 0.001), which was not as evident in micropropagated samples (R<sup>2</sup> = 0.48, p < 0.001).

The weight (moisture) loss from samples during assessment through the range of light intensities was  $6.9 \pm 2.4\%$  and  $7.4 \pm 2.9\%$  for micropropagated and wild-sourced, respectively; minimum and maximum values were 4.4% (*S. medium/divinum*) and 10.5% (*S. squarrosum*) in micropropagated samples and 4.6% (*S. papillosum*) and 9.7% (*S. fallax*) in wild-sourced samples, respectively. The moisture content of samples at  $P_{\max}$  ([Sample  $P_{\max}$  fresh weight—sample dry weight]/sample dry weight  $\times 100$ ) was  $2335 \pm 420\%$  (CV = 18%) (micropropagated) and  $1551 \pm 320\%$  (CV = 21%) (wild-sourced).

There were significantly more capitula per sample in micropropagated than in wild-sourced *Sphagnum* overall (mean  $\pm$  SD =  $41.8 \pm 24.2$  and  $27.0 \pm 18.7$ , respectively;  $t = 2.635$ ,  $p = 0.011$ ,  $df = 58$ ) (Examples in Figure 1). As noted in the Section 2, samples were placed carefully in a cylinder for analysis at growing shoot density. Moreover, there were more capitula in micropropagated than in wild-sourced samples of each species, although (according to two-way ANOVA post hoc Tukey's HSD) this was not significant for *S. fallax*, *S. medium/divinum*, and *S. squarrosum* (*S. capillifolium* and *S. papillosum*,  $p < 0.001$ ; *S. palustre*,  $p < 0.01$ ,  $n = 10$ ).

### 3.2. *Sphagnum* Samples' Bulk Density

The FW and DW bulk density were greater in wild-sourced than in micropropagated samples (not *S. fallax* or *S. palustre* by FW) ( $F = 22.3$  [FW],  $F = 73.6$  [DW];  $p < 0.001$ ,  $df = 11$  for both) (Figure 6). Differences between types (micropropagated and wild-sourced) were significant (ANOVA post hoc Tukey's HSD) for *S. capillifolium* (FW  $p = 0.019$ , DW  $p < 0.001$ ) *S. medium/divinum* (FW  $p = 0.003$ , DW  $p < 0.001$ ), and *S. papillosum* (FW and DW  $p < 0.001$ ) ( $n = 10$  throughout). Within types, the bulk density across micropropagated samples was less variable than wild-sourced samples (coefficient of variation of 34.7% and 45.3% by FW and 25.0% and 44.5% by DW, respectively), with a greater bulk density in *S. capillifolium*, *S. medium/divinum*, and *S. papillosum* than other species in wild-sourced samples. There was a statistically significant difference between species within both micropropagated and wild-sourced samples as determined by one-way ANOVA (FW:  $F = 4.71$ ,  $p = 0.004$ ,  $F = 44.65$ ,  $p < 0.001$  and DW:  $F = 3.99$ ,  $p = 0.009$ ,  $F = 70.37$ ,  $p < 0.001$ , respectively). One-way ANOVA post hoc Tukey's HSD statistically significant differences are shown on Figure 6.

The linear regression revealed significant negative relationships between DW bulk density (mass/volume) and both  $P_{\max}$  (DW) and respiration (DW) ( $P_{\max}$  and respiration rates decreased as the bulk density increased) of micropropagated and particularly wild-sourced samples ( $P_{\max}$ :  $R^2 = 0.573$  and  $0.827$ , respectively; respiration:  $R^2 = 0.503$  and  $0.789$  respectively;  $p < 0.001$ ,  $df = 29$  throughout).





Figure 6. Cont.



**Figure 6.** Comparison of micropropagated and wild-sourced *Sphagnum* density by fresh weight (FW) and dry weight (DW). Shared letters within types (micropropagated and wild-sourced) indicate significant differences in density between species (one-way ANOVA post hoc Tukey's HSD). Crosses indicate the mean, lines indicate the median, and interquartile range is exclusive with maximum and minimum values that are not outliers indicated by the whiskers.

### 3.3. *Sphagnum* Samples' Nutrient Content

Macronutrient (Ca, K, Mg, N, P, S) concentrations were significantly higher in micropropagated than in wild-sourced samples overall (Mann–Whitney U test: S:  $p = 0.001$ , all other elements:  $p < 0.001$ ;  $n = 60$  throughout) (Table 2). N made up the largest proportion of the macronutrient content in all species throughout, and there was a high proportion of K in all micropropagated samples and in most of the wild-sourced samples. The principal component analysis (Figure 7) showed a noticeable clustering of wild-sourced samples away from micropropagated *Sphagnum*, and a disassociation from increasing macronutrients (Ca, Mg, P, N, K, S) in wild-sourced samples (Figure 7A), apart from *S. squarrosum* (a minerotrophic species), which was more closely associated with micropropagated *Sphagnum*. In contrast, micropropagated *Sphagnum* was associated with macronutrients, although there were differences between macronutrients and species. N, P, and K were closely associated, and there was an association between these macronutrients and the micropropagated *S. palustre*, *S. papillosum*, and *S. squarrosum*.

**Table 2.** Macronutrient content of *Sphagnum* samples.

<b>Sphagnum Type/sp.</b>	<b>Ca</b>	<b>K</b>	<b>Mg</b>	<b>N</b>	<b>P</b>	<b>S</b>
microprop						
<i>S. cap</i>	4.30 ± 0.38 **	11.58 ± 0.52 **	1.51 ± 0.08 **	21.08 ± 0.72 **	2.90 ± 0.16 **	0.68 ± 0.03
wild						
<i>S. cap</i>	1.74 ± 0.27	3.32 ± 0.56	0.65 ± 0.13	12.63 ± 1.43	0.44 ± 0.09	0.69 ± 0.08
microprop						
<i>S. fal</i>	4.53 ± 1.17 **	10.25 ± 1.30 **	1.19 ± 0.19 **	19.68 ± 1.71 **	1.84 ± 0.27 **	0.59 ± 0.08
wild						
<i>S. fal</i>	0.84 ± 0.29	5.34 ± 0.51	0.39 ± 0.04	12.28 ± 2.63	0.55 ± 0.07	0.51 ± 0.06
microprop						
<i>S. med/div</i>	3.48 ± 0.49 **	10.32 ± 1.21 **	1.06 ± 0.11 **	20.41 ± 1.43 **	2.09 ± 0.32 **	0.57 ± 0.07
wild						
<i>S. med/div</i>	1.31 ± 0.33	2.05 ± 0.49	0.70 ± 0.09	10.77 ± 1.35	0.17 ± 0.01	0.49 ± 0.04

Table 2. Cont.

Sphagnum Type/sp.	Ca	K	Mg	N	P	S
microprop <i>S. pal</i>	4.47 ± 0.58 **	15.99 ± 1.51 **	1.55 ± 0.14 **	29.27 ± 0.78 **	2.82 ± 0.17 **	0.92 ± 0.10 **
wild <i>S. pal</i>	1.34 ± 0.11	6.80 ± 0.92	0.72 ± 0.11	13.94 ± 2.33	0.80 ± 0.08	0.52 ± 0.03
microprop <i>S. pap</i>	3.14 ± 0.28 **	15.40 ± 1.28 **	1.26 ± 0.06 **	28.69 ± 2.14 **	2.80 ± 0.18 **	0.85 ± 0.07 **
wild <i>S. pap</i>	1.60 ± 0.37	2.75 ± 0.36	0.58 ± 0.05	10.63 ± 1.21	0.25 ± 0.06	0.53 ± 0.04
microprop <i>S. squ</i>	2.82 ± 0.20 **	15.55 ± 1.43	1.19 ± 0.13 *	26.66 ± 3.11 **	2.86 ± 0.33 **	0.90 ± 0.12
wild <i>S. squ</i>	1.21 ± 0.19	13.61 ± 0.75	0.90 ± 0.07	20.90 ± 2.05	1.83 ± 0.16	0.90 ± 0.08

*n* = 5; values in mg g<sup>−1</sup> dry matter ± SD. Statistically significant differences through ANOVA Tukey’s post hoc tests between micropropagated (microprop) and wild-sourced (wild) species for each macronutrient are indicated: \* *p* < 0.05; \*\* *p* < 0.001. *S. capillifolium* (*S. cap*), *S. fallax* (*S. fal*), *S. medium/divinum* (*S. med/div*), *S. palustre* (*S. pal*), *S. papillosum* (*S. pap*), *S. squarrosum* (*S. squ*).




Figure 7. Principal component analysis (PCA: correlation matrix) of *Sphagnum* element content. PCA graphs show (A) macronutrient and (B) micronutrient and trace element contents of *Sphagnum* samples with P<sub>max</sub> and respiration by dry weight.



The levels of trace elements Al, Fe, Mn, Na, Ni, and Pb were higher in wild-sourced than micropropagated samples overall (Al, Fe, Pb:  $p < 0.001$ ; Mn:  $p = 0.01$ ; Na and Ni:  $p$  NS;  $n = 60$  throughout) (Table 3), with the highest levels of Al, Fe, Mn, and Na in *S. papillosum*. The levels of micronutrients Cu and Zn were higher in micropropagated than wild-sourced samples overall ( $p < 0.001$ ;  $n = 60$ ) (although not in *S. squarrosum*). Na made up the largest proportion of the micronutrient and trace element contents in all samples. The principal component analysis (Figure 7B) showed a clustering and association with Cu and Zn in micropropagated samples together with wild-sourced *S. squarrosum*. Other samples of wild-sourced *Sphagnum* were only loosely grouped by species and more closely associated with micronutrients other than Cu and Zn. Cu and Zn were closely associated, as were Fe and Pb, and Mn and Na. *S. papillosum* appeared to have a strong association with Na and Al, and *S. capillifolium*, *S. medium/divinum* and *S. fallax* with Fe and Pb.

**Table 3.** Micronutrient and trace element contents of *Sphagnum* samples.

Sphagnum Type/sp.	Al	Cu	Fe	Mn	Na	Ni	Pb	Zn
microprop	9.75 ± 0.38	5.88 ± 0.50	52.74 ± 5.10	39.12 ± 2.37 **	338.0 ± 35.03	0.15 ± 0.46	0.12 ± 0.03 **	47.92 ± 9.82 *
<i>S. cap</i>	91.66 ± 9.86	4.48 ± 0.87	143.7 ± 12.63	64.30 ± 12.54	470.7 ± 90.91	0.60 ± 0.71	2.27 ± 0.26	24.89 ± 7.55
<i>S. cap</i>								
microprop	15.58 ± 2.53	5.65 ± 0.78 **	35.44 ± 3.99 *	28.21 ± 5.38	505.3 ± 120.3	0.33 ± 0.37	0.25 ± 0.04 **	66.08 ± 17.08 **
<i>S. fal</i>	83.05 ± 10.78	2.68 ± 0.64	198.9 ± 21.32	19.32 ± 2.95	256.8 ± 29.67	0.18 ± 0.13	1.34 ± 0.50	7.81 ± 2.19
<i>S. fal</i>								
microprop	18.20 ± 5.43	7.43 ± 0.89 **	47.35 ± 4.18	27.95 ± 3.62	430.0 ± 76.53 **	0.89 ± 0.94	0.33 ± 0.09	76.96 ± 8.94 **
<i>S. med/div</i>	62.51 ± 12.35	3.29 ± 0.66	90.15 ± 17.61	29.50 ± 13.02	899.7 ± 195.3	0.32 ± 0.24	0.70 ± 0.06	10.74 ± 2.74
<i>S. med/div</i>								
microprop	13.32 ± 3.32	8.87 ± 0.62	66.89 ± 3.66	35.81 ± 4.45	497.9 ± 110.9	0.14 ± 0.22	0.17 ± 0.03	57.57 ± 5.83 **
<i>S. pal</i>	36.95 ± 4.80	4.68 ± 0.89	46.62 ± 6.59	89.78 ± 9.53	372.1 ± 51.82	0.54 ± 0.69	0.63 ± 0.14	18.53 ± 4.28
<i>S. pal</i>								
microprop	11.20 ± 2.65 **	7.79 ± 0.90 **	50.42 ± 3.84 *	29.56 ± 2.01 **	360.7 ± 55.17 **	0.34 ± 0.49	0.24 ± 0.03 **	38.47 ± 8.75 *
<i>S. pap</i>	303.8 ± 268.4	3.72 ± 1.06	245.5 ± 175.2	171.5 ± 102.6	1159.0 ± 387.5	1.08 ± 0.34	1.65 ± 0.88	16.92 ± 8.92
<i>S. pap</i>								
microprop	7.87 ± 1.44	6.29 ± 1.65	83.87 ± 83.46	28.30 ± 2.71	303.8 ± 39.22	0 ± 0.07	0.20 ± 0.15	40.44 ± 5.45 **
<i>S. squ</i>	35.17 ± 3.67	6.33 ± 1.01	52.79 ± 5.91	92.32 ± 11.15	448.2 ± 97.71	0.24 ± 0.10	0.70 ± 0.10	67.41 ± 6.03
<i>S. squ</i>								

$n = 5$ ; values in  $\mu\text{g g}^{-1}$  dry matter  $\pm$  SD. Statistically significant differences through ANOVA Tukey's post hoc tests between micropropagated (microprop) and wild-sourced (wild) species for each element are indicated: \*  $p < 0.05$ ; \*\*  $p < 0.001$ . *S. capillifolium* (*S. cap*), *S. fallax* (*S. fal*), *S. medium/divinum* (*S. med/div*), *S. palustre* (*S. pal*), *S. papillosum* (*S. pap*), *S. squarrosum* (*S. squ*).

There was a lower N:P and N:K ratio, and a lower variation between species in micropropagated than in wild-sourced samples: N:P =  $9.61 \pm 1.24$  (CV = 12.9%) and  $31.18 \pm 19.54$  (CV = 62.7%) respectively, and N:K =  $1.85 \pm 0.09$  (CV = 4.9%) and  $3.13 \pm 1.40$  (CV = 44.8%), respectively. The N:P and N:K ratios in wild-sourced *S. squarrosum* (11.41 and 1.54) and, to a lesser extent, *S. palustre* (17.51 and 2.05) were the most similar to their micropropagated equivalents, and those of wild-sourced *S. medium/divinum* (64.59 and 5.24) and *S. papillosum* (42.56 and 3.86) the most dissimilar, respectively.

### 3.4. Sphagnum Cell Measurements and Analysis


There were significant differences between the groups of species by type (micropropagated or wild-sourced) in both chlorocyst width and number of chloroplasts ( $F = 49.9$ ,  $F = 33.7$ , respectively;  $p < 0.001$ ,  $df = 11$  for both) (Table 4). There were no significant differences (ANOVA post hoc Tukey's HSD) in chlorocyst width between micropropagated and wild-sourced species individually (Table 4), except for *S. squarrosum* ( $p < 0.05$  wild-sourced > micropropagated). The widest chlorocysts were recorded in *S. palustre* (micropropagated and wild-sourced samples). There were more chloroplasts in micropropagated compared to wild-sourced species in all but *S. squarrosum* (Table 4), and differences were significant (ANOVA post hoc Tukey's HSD) for *S. capillifolium* and *S. palustre* ( $p < 0.05$ ) and *S. papillosum*

( $p < 0.001$ ). The greatest chloroplast numbers in each type were found in *S. palustre* and *S. papillosum* (micropropagated) and *S. palustre* (wild-sourced). Physical differences between micropropagated and wild-sourced samples are limited to reduced color expression (*S. capillifolium* and *S. medium/divinum*) (Figure 1) and maturity (*S. papillosum* cell papillae) in micropropagated *Sphagnum* (Figure 8).

**Table 4.** Comparison of microscopic features between micropropagated and wild-sourced *Sphagnum* samples.

Sphagnum Species	Cell Width ( $\mu\text{m}$ )		Chloroplast No.	
	Micropropagated	Wild-Sourced	Micropropagated	Wild-Sourced
<i>S. capillifolium</i>	$7.14 \pm 0.91$	$6.65 \pm 0.65$	$16.6 \pm 2.7^*$	$10.6 \pm 1.1$
<i>S. fallax</i>	$7.14 \pm 0.83$	$6.96 \pm 0.39$	$11.1 \pm 2.7$	$7.1 \pm 1.7$
<i>S. medium/divinum</i>	$9.86 \pm 0.72$	$10.10 \pm 1.36$	$16.7 \pm 2.6$	$13.1 \pm 2.9$
<i>S. palustre</i>	$12.37 \pm 0.93$	$11.05 \pm 1.24$	$24.5 \pm 3.7^*$	$19.8 \pm 4.3$
<i>S. papillosum</i>	$9.88 \pm 0.59$	$9.84 \pm 0.31$	$25.6 \pm 3.9^{**}$	$12.2 \pm 2.2$
<i>S. squarrosum</i>	$5.65 \pm 0.59^*$	$7.30 \pm 1.36$	$11.9 \pm 2.4$	$15.1 \pm 3.1$

Mean chlorocyst (cell) width ( $\mu\text{m}$ ) and mean number of chloroplasts per chlorocyst; mean values ( $n = 9$ )  $\pm$  SD; significant differences (ANOVA Tukey's post hoc HSD) between each pair are indicated: \*  $p < 0.05$ ; \*\*  $p < 0.001$ .



**Figure 8.** Examples of *Sphagnum* samples at 1000 $\times$  magnification: micropropagated (top of pair) and wild-sourced *Sphagnum* samples. Scale bar = 50  $\mu\text{m}$ .

#### 4. Discussion

The routine micropropagation of *Sphagnum* in controlled environments has enabled a comparison of species physiology when grown under the same conditions without the confounding influences of habitat conditions and plastic responses. The maximum photosynthesis ( $P_{\text{max}}$ ) capacity and respiration rate measured on a range of *Sphagnum* species were broadly in line with the published data for these species (Table S1), although rates varied more across wild-sourced than micropropagated species, and all were typically much lower than vascular plants [9,12,33,34,42,43].

Several authors [9,10,43] suggest a rank in *Sphagnum* species photosynthesis rates, corresponding generally to the accepted phylogenetic order of growth rates and a gradient in production depending on light, moisture, and nutrient levels [34]. In general, highly productive *Sphagnum* tend to be competitive or ruderal [44] species of hollows (such as *S. squarrosum*, *S. fallax*, and *S. fimbriatum*), thriving in shaded, high moisture, and nutrient environments (i.e., low-stress) and adopting an open, loose growth habit [10]. This open, loose growth habit also allows more of the plant access to any available light for photosynthesis and faster growth rates [45]. These species also tend to be green, likely due to a high chlorophyll content [10], with strong shoot growth and large capitula [30]. *Sphagnum* with lower productivity tend to be stress-adapted species (such as *S. capillifolium*, *S. medium/divinum*, and *S. papillosum*), which grow in open (unshaded), ombrotrophic bogs, subject to high light intensity, occasional drought, and low nutrient levels (i.e., high-stress) [11,12]. In these conditions, short, dense growth forms for greater acquisition and

retention of moisture [10,24] and perhaps tolerance of photoinhibition become more important to survival than the capacity for rapid growth [9,24]. Other more generalist species (such as *S. palustre*, and some other loose hummock-forming species) thrive at points along this stress gradient. There are few *Sphagnum* in unshaded, dry sites. Thus, the capacity for photosynthesis is driven by moisture, light, and nutrient levels, and different phylogenetic traits, as described above, provide *Sphagnum* species competitive advantages to thrive in particular habitats [26]. This is reflected in our study, and to answer research question (1), whether grown in the same conditions (micropropagated) or sourced from the wild, the species with the highest capacity for photosynthesis and respiration (on a dry matter basis) was *S. squarrosum* and that with the lowest was *S. medium/divinum*, suggesting that the accepted rank, as described above, does have a phylogenetic basis.

The wide range of  $P_{\max}$  rates across samples in this study was likely due to the diversity of *Sphagnum* species and sources: micropropagated species grown in a commercial greenhouse environment (although original source material was from different sites); wild-sourced species developed each with a range of nutrient and shade regimes in their habitat. It is notable that *S. medium/divinum*  $P_{\max}$  rates were higher in shaded than open sites in a study by Bengtsson et al. [43] (Table S1). This not only demonstrates the plasticity of this species in its adaptation to a range of environmental conditions but suggests that a shaded environment is likely to promote moisture retention in *Sphagnum* due to reduced evaporative effects of wind and heat, which subsequently supports photosynthesis [10]. Regimes for the storage, hydration, acclimatization, and analysis of samples were standardized in our study. Hájek [26] suggested the water content of ‘well-hydrated’ *Sphagnum* is 1500 to 3000%, and both micropropagated and wild-sourced samples in this study were within this range. However, wild-sourced samples were at the lower end of the range and had 66% of the water content of micropropagated samples, which may have had some influence on comparative photosynthesis rates.

Micropropagated *Sphagnum* species had a markedly greater response to increasing light levels and higher  $P_{\max}$  and respiration rates than the wild-sourced ones, on a dry weight basis, which supports research question (2) that there is a difference in photosynthetic capacity and respiration between micropropagated and wild-sourced *Sphagnum* species. Across all species samples, the wild-sourced *Sphagnum*  $P_{\max}$  was 34.8% (by dry weight) that of the micropropagated *Sphagnum*. This is perhaps because the micropropagated samples in a shaded commercial greenhouse had not yet developed photo-inhibitive adaptations to light stress. However, in both micropropagated and wild-sourced samples, the competitive, shade species *S. squarrosum* had the highest rates of  $P_{\max}$  and respiration and the lowest bulk density (mass/volume), and the stress-adapted species *S. medium/divinum* had the lowest rates of  $P_{\max}$  and respiration, and the highest bulk density, demonstrating some parity in species behavior, despite the tissue-culture process.

$P_{\max}$  and respiration rates were positively related throughout as typically found across ecosystems [46], most strongly in wild-sourced species, suggesting this more mature, natural material may have reached an equilibrium within its habitat, compared to the more even photosynthetic response of micropropagated species to light. There was a higher ratio of  $P_{\max}$  to respiration in micropropagated than in wild-sourced species, despite generally higher micropropagated respiration rates, suggesting potential carbon balance benefits on application to the field, at least in the early stages of establishment.

Differences in  $P_{\max}$  between micropropagated and wild-sourced samples were particularly apparent in species from section *Sphagnum*: *S. medium/divinum*, *S. palustre*, and *S. papillosum*, where photosynthetic activity was noticeably lower in wild-sourced than micropropagated samples. The difference between  $P_{\max}$  rates of micropropagated and wild-sourced samples of *S. capillifolium* was not so marked as other species. This species is adapted to a range of peatland habitats and ecological niches, and has tolerance to shade [11]. Wild-sourced, shade-tolerant species, *S. squarrosum*, *S. fallax*, and *S. palustre*, had the highest  $P_{\max}$  levels at the lowest light levels, on a dry weight basis, which concurs with findings by Rice et al. [9] and Hájek et al. [10].



The DW bulk density of wild-sourced *Sphagnum* for each species studied was significantly greater than that of micropropagated species, which were established under favorable light and moisture regimes, and were also in the early stages of rapid, linear growth [12]. There was also a relationship between increasing bulk density and declining  $P_{\max}$  in wild-sourced *Sphagnum* samples but not in micropropagated samples. In wild-sourced samples, there were two distinct groups by dry weight bulk density: *S. fallax*, *S. palustre*, *S. squarrosum* (low bulk density), and *S. capillifolium*, *S. papillosum*, *S. medium/divinum* (high bulk density), showing obvious differences between shade-adapted, moisture-dependent species, and light-adapted, moisture-retaining species, which are only fully expressed in the natural environment.

Micropropagated *Sphagnum* species were not visually different from those wild-sourced, although some characteristics, such as *S. papillosum* cell papillae, were less developed in micropropagated samples, suggesting immaturity (Figure 8), and there was less color expression in micropropagated *Sphagnum* (Figure 1), grown in a shaded greenhouse, as secondary pigments usually only develop in full light [40] (p. 61). Chlorocyst size was not significantly different, despite the relative immaturity of micropropagated samples, apart from *S. squarrosum* species, where those of wild-sourced samples were significantly larger than micropropagated samples and the number of chloroplasts per cell was also greater. *S. squarrosum* samples were sourced from a nutrient-rich, shaded environment, beneficial to continued upward growth; the greater cell size and number of chloroplasts were likely based on maturity and optimum growing conditions. In all other species, there was a significantly greater number of chloroplasts in micropropagated than in wild-sourced samples, apart from *S. palustre*, where the difference was not statistically significant. This is consistent with micropropagated plants being in the early stages of rapid growth [12] but also not exposed to conditions of high light intensity and low moisture, and so, they were perhaps acting more like shade plants [47]. Therefore, the aspect of research question (3) examining whether any differences in chlorocyst size and chloroplast number between micropropagated and wild-sourced *Sphagnum* species may affect their photosynthesis and respiration did not lead to a clear conclusion, as any differences were not directly linked to different photosynthetic capacity. However, the high rates of photosynthesis in micropropagated *Sphagnum*, the greater number of capitula per sample, as well as similar features to wild-sourced suggest that successful establishment in optimum field conditions is likely.

The nutrient analysis of the samples allowed the further examination of the causes of difference in photosynthesis rates. Micropropagated samples contained significantly higher levels of macronutrients, essential for plant photosynthesis and growth, than wild-sourced samples, and as  $\text{CO}_2$  uptake and emission rates were also higher in micropropagated *Sphagnum*, it appears that the aspect of research question (3), which examined whether any differences in nutrient content between micropropagated and wild-sourced *Sphagnum* species may affect their photosynthesis and respiration, was supported.

The higher levels of macronutrients in micropropagated *Sphagnum* may be explained through the standard horticultural processes of nutrient application associated with micropropagated production. Conversely, wild-sourced *Sphagnum* had higher levels than micropropagated of micronutrients, which are a smaller component of plant nutritional requirements but also support plant health. Concentrations of micronutrients may have become diluted in micropropagated plants due to their rapid growth rate. Additionally, higher levels of elements considered detrimental to plant health (such as Al, Fe, and Pb) in some wild-sourced species may reflect the capability of bryophytes to absorb certain levels of pollution and act as bioindicators [4,48].

The three main macronutrients associated with plant growth and photosynthesis, N, P, and K, were strongly positively associated with  $P_{\max}$  overall in this study and reflected species' differences in macronutrient uptake and  $P_{\max}$  rates. For example, wild-sourced species with the highest content of NPK were *S. squarrosum* and *S. palustre*, and those with the lowest content were *S. papillosum* and *S. medium/divinum*, which corresponded with  $P_{\max}$  rates for those species.



In vascular plants, there is a strong positive relationship between both net photosynthesis and dark respiration and leaf N content in a wide range of global biomes [42]. Typically, mosses translocate nutrients into their tissues gradually over the growing season [49] and the N:P and N:K ratios remain balanced [50]. Granath et al. [51] and Mazziotta et al. [28] found a positive relationship between N concentration and photosynthesis rates in *Sphagnum* (as in this study). However, *Sphagnum* growth reportedly improves with increasing levels of N until a 'critical concentration' is reached, at which there is no further promotion of plant growth despite increasing levels of N in tissues [52]. In the context of aerial nitrogen pollution, Press et al. [53] found that increasing inorganic nitrogen supply slowed *Sphagnum* growth. Saturated N levels limit further P and K accumulation, and so, growth becomes P- and K-limited [4,54,55]. Excess N is also leached into the environment, promoting the growth of vascular plants that may outcompete *Sphagnum* in the field [27,55,56].

The balance between N, P, and K levels in *Sphagnum* tissues appears to be similar throughout natural bog settings, but differences may occur due to growth habit and conditions. Wang and Moore [57] reported averages of  $N = 9$ ,  $P = 0.55$ ,  $K = 7.5 \text{ mg g}^{-1} \text{ DW}$  content for 'hummock' *Sphagna* (*S. capillifolium* and *S. medium/divinum*) at the Mer Bleue ombrotrophic bog in Canada. Bragazza et al. [4] had similar values of  $N = 8.2$  and  $9.2$ ,  $P = 0.41$  and  $0.44$ ,  $K = 4$  and  $4.44 \text{ mg g}^{-1} \text{ DW}$  for hummock and lawn species, respectively, in European mires, which are reported to have higher N deposition than Canadian mires. NPK levels of wild-sourced species in this study had higher N values but were otherwise similar to European values:  $N = 12.0 \pm 2.13$ ,  $P = 0.44 \pm 0.24$ ,  $K = 4.05 \pm 1.87 \text{ mg g}^{-1} \text{ DW}$ . Wild-sourced *S. squarrosum*, however, had a higher NPK content ( $N = 20.9 \pm 2.05$ ,  $P = 1.83 \pm 0.16$ ,  $K = 13.6 \pm 0.75 \text{ mg g}^{-1} \text{ DW}$ ) and, as the sample was sourced from wet woodland, suggests that nutrients may continue to accumulate in fast-growing *Sphagnum* in conditions of shade [58] and optimum moisture [37].

All micropropagated species had a high NPK content ( $N = 24.2 \pm 4.41$ ,  $P = 2.55 \pm 0.49$ ,  $K = 13.2 \pm 2.79 \text{ mg g}^{-1} \text{ DW}$ ) due to greenhouse nutrient additions. *Sphagnum* N concentration thresholds of 11 to  $12 \text{ mg g}^{-1}$  [4,55],  $15 \text{ mg g}^{-1}$  (for *S. recurvum*—section *Cuspidata*) [59], and  $20 \text{ mg g}^{-1}$  (for a range of hummock and hollow species) [56] have been reported. Nitrogen content in wild-sourced *S. squarrosum* in this study was at the top of the reported threshold range, with no apparent limitation of P or K, although this may be due to the fast-growing nature of this species, and nutrients are readily replenished by the woodland habitat from which it was sourced. Nitrogen content in micropropagated samples was well above the highest reported threshold—as high as  $30 \text{ mg g}^{-1}$  in some samples, with no evidence of toxicity or limitation of P or K.

Growth reduction is reported when *Sphagnum* experiences a considerable increase in N availability relative to conditions at the site from which it was obtained [60], implying that *Sphagnum* tolerance of N is site-adapted. Bragazza et al. [4] found that, at elevated N levels, the N:P ratio increased to 33.8 and 33.6 and N:K ratio increased to 3.7 and 4.0 for hummock and lawn species, respectively. In this study, there were very high N:P and marginally high N:K ratios in *S. medium/divinum* (64.6 and 5.24) and *S. papillosum* (42.6 and 3.86 respectively) from open, ombrotrophic bogs compared to N:P and N:K ratios in *S. squarrosum* (11.4 and 1.54) and *S. palustre* (17.5 and 2.05, respectively) (from higher-nutrient, shaded environments), which are more aligned to ratios found in low nutrient sites by Wang and Moore [57] and Bragazza et al. [4]. In wild-sourced *Sphagnum* in this study, the lowest  $P_{\max}$  was found in *S. papillosum* and *S. medium/divinum*, and the highest in *S. squarrosum* and *S. palustre*. Although the plants with a high N:P ratio (*S. papillosum* and *S. medium/divinum*) appeared healthy and are likely adapted to these particular levels of nutrients, the limitation of P by higher levels of N (as described above) may have reduced their photosynthetic capacity [61], although Granath et al. [51] found that the negative effect of limited P was on production rather than photosynthetic rate.

This difference in stoichiometry between micropropagated and wild-sourced plants may be due both to issues of maturity [50], as levels of N may be naturally higher in young than in mature plants [52], and site adaptation in wild-sourced samples [60], i.e., plant

adaptations to environmental stressors inhibited photosynthetic potential in wild-sourced plants. Although micropropagated plants were ‘mature’ in terms of development, they could be classed as ‘young’, being grown directly from tissue culture to the point at which they are available as plug plants, ready for site application, after only several months of growth. Micropropagated plants do not appear to be P- or K-limited ( $N:P = 9.61 \pm 1.24$  and  $N:K = 1.85 \pm 0.09$ ), suggesting that the right balance of nutrients is applied in the horticultural process for these new plants.

The most important aspect of *Sphagnum* reintroduction to peatland restoration sites is rapid establishment and growth, particularly lateral growth, to make an intact carpet as quickly as possible [62]. This will progress restoration by keeping a cool, moist layer at the peat surface [63], reducing evaporation and encouraging the development of an acrotelm to promote peat accumulation [19,64–66]. The capacity of micropropagated *Sphagnum* to absorb nitrogen above normally limited thresholds may also be useful on application to ex-agricultural sites as a paludiculture crop or for restoration purposes, at least in the critical establishment phase. This study on differences between micropropagated and wild-sourced *Sphagnum* in terms of net photosynthesis, phylogenetic and adaptive traits, and nutrient assimilation highlights potential reasons for the successful establishment and growth of micropropagated products reported in large-scale restoration projects [20–22] and their potential contribution to CO<sub>2</sub> uptake. Future work could usefully determine whether the advantages conveyed in these early growth stages of micropropagated *Sphagnum* moss persist over longer time periods in the wild.

**Supplementary Materials:** The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/ijpb15040068/s1>, Table S1: Comparison of light-saturated photosynthesis and respiration rates between samples in this study and values in the literature.

**Author Contributions:** Conceptualization, A.T.K., C.D.F., J.G.R. and S.J.M.C.; methodology, A.T.K. and S.J.M.C.; software, A.T.K.; validation, A.T.K., C.D.F., J.G.R. and S.J.M.C.; formal analysis, A.T.K.; investigation, A.T.K.; resources, S.J.M.C.; data curation, A.T.K.; writing—original draft preparation, A.T.K.; writing—review and editing, A.T.K., C.D.F., J.G.R. and S.J.M.C.; visualization, A.T.K., C.D.F., J.G.R. and S.J.M.C.; supervision, S.J.M.C., C.D.F. and J.G.R.; project administration, S.J.M.C.; funding acquisition, S.J.M.C. All authors have read and agreed to the published version of the manuscript.

**Funding:** This paper was derived from a doctoral thesis (PhD) part-funded by Micropropagation Services (EM) Ltd. (trading as BeadaMoss®), match-funded by Manchester Metropolitan University (MMU). The authors wish to thank David McKendry and technical officers at MMU for their support, and are grateful to the examiners of the PhD thesis who provided valuable comments and corrections.

**Institutional Review Board Statement:** Not applicable.

**Data Availability Statement:** The original data presented in the study are openly available in Mendeley Data at DOI:10.17632/dfvmfmb3vz.1. <https://data.mendeley.com/datasets/dfvmfmb3vz/1> (accessed on 24 April 2024).

**Conflicts of Interest:** Author J.G.R. was employed by the company Jacobs UK Limited. The remaining authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest. The funding sponsors had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, and in the decision to publish the results.

## References

1. Verhoeven, J.T.; Liefveld, W.M. The ecological significance of organochemical compounds in *Sphagnum*. *Acta Bot. Neerl.* **1997**, *46*, 117–130. [CrossRef]
2. Rydin, H.; Jeglum, J.K.; Bennett, K.D. *The Biology of Peatlands*, 2nd ed.; Oxford University Press: Oxford, UK, 2013.
3. Lindsay, R.; Birnie, R.; Clough, J. *Peat Bog Ecosystems: Key Definitions*; International Union for the Conservation of Nature: Edinburgh, UK, 2014; Available online: <https://repository.uel.ac.uk/item/85870> (accessed on 24 April 2024).
4. Bragazza, L.; Tahvanainen, T.; Kutnar, L.; Rydin, H.; Limpens, J.; Hájek, M.; Grosvernier, P.; Hájek, T.; Hajkova, P.; Hansen, I.; et al. Nutritional constraints in ombrotrophic *Sphagnum* plants under increasing atmospheric nitrogen deposition in Europe. *New Phytol.* **2004**, *163*, 609–616. [CrossRef] [PubMed]

5. Fritz, C.; Lamers, L.P.; Riaz, M.; van den Berg, L.J.; Elzenga, T.J. *Sphagnum* mosses-masters of efficient N-uptake while avoiding intoxication. *PLoS ONE* **2014**, *9*, e79991. [CrossRef]
6. van Breemen, N. How *Sphagnum* bogs down other plants. *TREE* **1995**, *10*, 270–275. [CrossRef]
7. Malmer, N.; Albinsson, C.; Svensson, B.M.; Wallén, B. Interferences between *Sphagnum* and vascular plants: Effects on plant community structure and peat formation. *Oikos* **2003**, *100*, 469–482. [CrossRef]
8. Clymo, R.S. Interactions of *Sphagnum* with water and air. In *Effects of Atmospheric Pollutants on Forests, Wetlands and Agricultural Systems*; Hutchinson, T.C., Meema, K.M., Eds.; Springer: Berlin, Germany, 1987; pp. 513–530.
9. Rice, S.K.; Aclander, L.; Hanson, D.T. Do bryophyte shoot systems function like vascular plant leaves or canopies? Functional trait relationships in *Sphagnum* mosses (*Sphagnaceae*). *Am. J. Bot.* **2008**, *95*, 1366–1374. [CrossRef]
10. Hájek, T.; Tuittila, E.S.; Ilomets, M.; Laiho, R. Light responses of mire mosses—a key to survival after water-level drawdown? *Oikos* **2009**, *118*, 240–250. [CrossRef]
11. Bonnett, S.A.; Ostle, N.; Freeman, C. Short-term effect of deep shade and enhanced nitrogen supply on *Sphagnum capillifolium* morphophysiology. *Plant Ecol.* **2010**, *207*, 347–358. [CrossRef]
12. Laine, A.M.; Juurola, E.; Hájek, T.; Tuittila, E.S. *Sphagnum* growth and ecophysiology during mire succession. *Oecologia* **2011**, *167*, 1115–1125. [CrossRef]
13. Joosten, H.; Sirin, A.; Couwenberg, J.; Laine, J.; Smith, P. The role of peatlands in climate regulation. In *Peatland Restoration and Ecosystem Services: Science, Policy and Practice*; Bonn, A., Allott, T., Evans, M., Joosten, H., Stoneman, R., Eds.; British Ecological Society, Cambridge University Press: Cambridge, UK, 2016; pp. 63–76.
14. Gregg, R.; Elias, J.; Alonso, I.; Crosher, I.; Muto, P.; Morecroft, M. Carbon sequestration by habitat: A review of the evidence. In *Natural England Research Report NERR094*; Natural England: York, UK, 2021; Available online: <https://www.cabidigitallibrary.org/doi/full/10.5555/20220357895> (accessed on 1 September 2024).
15. Evans, C.; Artz, R.; Moxley, J.; Smyth, M.A.; Taylor, E.; Archer, E.; Burden, A.; Williamson, J.; Donnelly, D.; Thomson, A.; et al. *Implementation of an Emissions Inventory for UK Peatlands*; Centre for Ecology and Hydrology: Report to the Department for Business, Energy and Industrial Strategy; Centre for Ecology and Hydrology: Bangor, UK, 2017.
16. 2017 UK Greenhouse Gas emissions, final figures. In *Statistical Release: National Statistics*; Department for Business, Energy and Industrial Strategy: London, UK, 2019. Available online: [https://assets.publishing.service.gov.uk/media/5c584a12e5274a317cc0eaf7/2017\\_Final\\_emissions\\_statistics\\_-\\_report.pdf](https://assets.publishing.service.gov.uk/media/5c584a12e5274a317cc0eaf7/2017_Final_emissions_statistics_-_report.pdf) (accessed on 24 April 2024).
17. Wilson, D.; Farrell, C.; Mueller, C.; Hepp, S.; Renou-Wilson, F. Rewetted industrial cutaway peatlands in western Ireland: A prime location for climate change mitigation? *Mires Peat* **2013**, *11*(1), 1–22.
18. Renou-Wilson, F.; Moser, G.; Fallon, D.; Farrell, C.A.; Müller, C.; Wilson, D. Rewetting degraded peatlands for climate and biodiversity benefits: Results from two raised bogs. *Ecol. Eng.* **2019**, *127*, 547–560. [CrossRef]
19. Quinty, F.; Rochefort, L. *Peatland Restoration Guide*, 2nd ed.; Canadian Sphagnum Peat Moss Association and New Brunswick Department of Natural Resources and Energy: Québec, QC, Canada, 2003; Available online: [https://www1.up.poznan.pl/glinbar/wp-content/uploads/2015/03/Restoration-Guide\\_2nd\\_2003.pdf](https://www1.up.poznan.pl/glinbar/wp-content/uploads/2015/03/Restoration-Guide_2nd_2003.pdf) (accessed on 27 May 2024).
20. Caporn, S.J.; Rosenburgh, A.E.; Keightley, A.T.; Hinde, S.L.; Riggs, J.L.; Buckler, M.; Wright, N.A. *Sphagnum* restoration on degraded blanket and raised bogs in the UK using micropropagated source material: A review of progress. *Mires Peat* **2018**, *20*, 1–17.
21. Crouch, T. *Kinder Scout Sphagnum Trials: 2018 Update Report*; Moors for the Future Report: Edale, UK, 2018; Available online: [https://www.moorsforthefuture.org.uk/\\_data/assets/pdf\\_file/0027/93933/MFFP-Kinder-Scout-Sphagnum-Trials-Update-Report-2018.pdf](https://www.moorsforthefuture.org.uk/_data/assets/pdf_file/0027/93933/MFFP-Kinder-Scout-Sphagnum-Trials-Update-Report-2018.pdf) (accessed on 27 May 2024).
22. Pilkington, M.; Walker, J.; Fry, C.; Eades, P.; Meade, R.; Pollett, N.; Rogers, T.; Helliwell, T.; Chandler, D.; Fawcett, E.; et al. Diversification of *Molinia*-dominated blanket bogs using *Sphagnum* propagules. *Ecol. Solut. Evid.* **2021**, *2*, e12113. [CrossRef]
23. Bengtsson, F.; Rydin, H.; Baltzer, J.L.; Bragazza, L.; Bu, Z.J.; Caporn, S.J.; Dorrepaal, E.; Flatberg, K.I.; Galanina, O.; Galka, M.; et al. Environmental drivers of *Sphagnum* growth in peatlands across the Holarctic region. *J. Ecol.* **2021**, *109*, 417–431. [CrossRef]
24. Loisel, J.; Gallego-Sala, A.V.; Yu, Z. Global-scale pattern of peatland *Sphagnum* growth driven by photosynthetically active radiation and growing season length. *Biogeosciences* **2012**, *9*, 2737–2746. [CrossRef]
25. Hassel, K.; Kyrkjeeide, M.O.; Yousefi, N.; Prestø, T.; Stenøien, H.K.; Shaw, J.A.; Flatberg, K.I. *Sphagnum divinum* (sp. nov.) and *S. medium* Limpr. and their relationship to *S. magellanicum* Brid. *J. Bryol.* **2018**, *40*, 197–222. [CrossRef]
26. Hájek, T. Physiological ecology of peatland bryophytes. In *Photosynthesis in Bryophytes and Early Land Plants*; Hanson, D.T., Rice, S.K., Eds.; Springer: Dordrecht, The Netherlands, 2013; pp. 233–252.
27. Bubier, J.L.; Moore, T.R.; Bledzki, L.A. Effects of nutrient addition on vegetation and carbon cycling in an ombrotrophic bog. *Glob. Change Biol.* **2007**, *13*, 1168–1186. [CrossRef]
28. Mazziotta, A.; Granath, G.; Rydin, H.; Bengtsson, F.; Norberg, J. Scaling functional traits to ecosystem processes: Towards a mechanistic understanding in peat mosses. *J. Ecol.* **2019**, *107*, 843–859. [CrossRef]
29. Aldous, A.R. Nitrogen translocation in *Sphagnum* mosses: Effects of atmospheric nitrogen deposition. *New Phytol.* **2002**, *156*, 241–253. [CrossRef]
30. Laing, C.; Granath, G.; Belyea, L.R.; Allton, K.E.; Rydin, H. Tradeoffs and scaling of functional traits in *Sphagnum* as drivers of carbon cycling in peatlands. *Oikos* **2014**, *123*, 817–828. [CrossRef]

31. Black, C.C., Jr.; Mollenhauer, H.H. Structure and distribution of chloroplasts and other organelles in leaves with various rates of photosynthesis. *Plant Physiol.* **1971**, *47*, 15–23. [[CrossRef](#)] [[PubMed](#)]
32. Murashige, T.; Skoog, F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiol. Plant* **1962**, *15*, 473–497. [[CrossRef](#)]
33. Haraguchi, A.; Yamada, N. Temperature dependency of photosynthesis of *Sphagnum* spp. distributed in the warm-temperate and the cool-temperate mires of Japan. *Am. J. Plant Sci.* **2011**, *2*, 716. [[CrossRef](#)]
34. Kangas, L.; Maanavilja, L.; Hájek, T.; Juurola, E.; Chimner, R.A.; Mehtätalo, L.; Tuittila, E.S. Photosynthetic traits of *Sphagnum* and feather moss species in undrained, drained and rewetted boreal spruce swamp forests. *Ecol. Evol.* **2014**, *4*, 381–396. [[CrossRef](#)]
35. Dossa, G.G.; Paudel, E.; Wang, H.; Cao, K.; Schaefer, D.; Harrison, R.D. Correct calculation of CO<sub>2</sub> efflux using a closed-chamber linked to a non-dispersive infrared gas analyzer. *Methods Ecol. Evol.* **2015**, *6*, 1435–1442. [[CrossRef](#)]
36. Limpens, J.; Berendse, F. How litter quality affects mass loss and N loss from decomposing *Sphagnum*. *Oikos* **2003**, *103*, 537–547. [[CrossRef](#)]
37. McNeil, P.; Waddington, J.M. Moisture controls on *Sphagnum* growth and CO<sub>2</sub> exchange on a cutover bog. *J. Appl. Ecol.* **2003**, *40*, 354–367. [[CrossRef](#)]
38. Smith, A.J. *The Moss Flora of Britain and Ireland*, 2nd ed.; Cambridge University Press: Cambridge, UK, 2004.
39. Laine, J.; Flatberg, K.I.; Harju, P.; Timonen, T.; Minkinen, K.J.; Laine, A.; Tuittila, E.S.; Vasander, H.T. *Sphagnum mosses: The Stars of European Mires*; Sphagna Ky: Helsinki, Finland, 2018.
40. Atherton, I.; Bosanquet, S.; Lawley, M. (Eds.) *Mosses and Liverworts of Britain and Ireland: A field guide*; British Bryological Society: Plymouth, UK, 2010.
41. Hammer, Ø.; Harper, D.A. Past: Paleontological statistics software package for education and data analysis. *Palaeont. Electr.* **2001**, *4*, 1.
42. Reich, P.B.; Walters, M.B.; Ellsworth, D.S. From tropics to tundra: Global convergence in plant functioning. *Proc. Natl. Acad. Sci. USA* **1997**, *94*, 13730–13734. [[CrossRef](#)]
43. Bengtsson, F.; Granath, G.; Rydin, H. Photosynthesis, growth, and decay traits in *Sphagnum*—A multispecies comparison. *Ecol. Evol.* **2016**, *6*, 3325–3341. [[CrossRef](#)]
44. Grime, J.P. Evidence for the existence of three primary strategies in plants and its relevance to ecological and evolutionary theory. *Am. Nat.* **1977**, *111*, 1169–1194. [[CrossRef](#)]
45. Krebs, M.; Gaudig, G.; Joosten, H. Record growth of *Sphagnum papillosum* in Georgia (Transcaucasus): Rain frequency, temperature and microhabitat as key drivers in natural bogs. *Mires Peat* **2016**, *18*, 1–16.
46. Chapin, F.S.; Matson, P.A.; Vitousek, P.M.; Chapin, F.S.; Matson, P.A.; Vitousek, P.M. *Principles of Terrestrial Ecosystem Ecology*; Springer: Berlin/Heidelberg, Germany, 2011.
47. Marschall, M.; Proctor, M.C. Are bryophytes shade plants? Photosynthetic light responses and proportions of chlorophyll a, chlorophyll b and total carotenoids. *Ann. Bot.* **2004**, *94*, 593–603. [[CrossRef](#)] [[PubMed](#)]
48. Blagny, R.; Paliulis, D. Research into heavy metals pollution of atmosphere applying moss as bioindicator: A literature review. *Environ. Res. Eng. Manag.* **2010**, *54*, 26–33.
49. Chapin, F.S., III; Johnson, D.A.; McKendrick, J.D. Seasonal movement of nutrients in plants of differing growth form in an Alaskan tundra ecosystem: Implications for herbivory. *J. Ecol.* **1980**, *68*, 189–209.
50. Sterner, R.W.; Elser, J.J. Ecological stoichiometry: The biology of elements from molecules to the biosphere. In *Ecological Stoichiometry*; Princeton University Press: Princeton, NJ, USA, 2017.
51. Granath, G.; Strengbom, J.; Rydin, H. Direct physiological effects of nitrogen on *Sphagnum*: A greenhouse experiment. *Funct. Ecol.* **2012**, *26*, 353–364. [[CrossRef](#)]
52. Barker, A.V.; Pilbeam, D.J. (Eds.) *Handbook of Plant Nutrition*, 2nd ed.; CRC Press: Boca Raton, FL, USA, 2015.
53. Press, M.C.; Woodin, S.J.; Lee, J.A. The potential importance of an increased atmospheric nitrogen supply to the growth of ombrotrophic *Sphagnum* species. *New Phytol.* **1986**, *103*, 45–55. [[CrossRef](#)]
54. Aerts, R.; Wallen, B.O.; Malmer, N. Growth-limiting nutrients in *Sphagnum*-dominated bogs subject to low and high atmospheric nitrogen supply. *J. Ecol.* **1992**, *80*, 131–140. [[CrossRef](#)]
55. Lamers, L.P.; Bobbink, R.; Roelofs, J.G. Natural nitrogen filter fails in polluted raised bogs. *Glob. Change Biol.* **2000**, *6*, 583–586. [[CrossRef](#)]
56. Berendse, F.; van Breemen, N.; Rydin, H.; Buttler, A.; Heijmans, M.; Hoosbeek, M.R.; Lee, J.A.; Mitchell, E.; Saarinen, T.; Vasander, H.; et al. Raised atmospheric CO<sub>2</sub> levels and increased N deposition cause shifts in plant species composition and production in *Sphagnum* bogs. *Glob. Change Biol.* **2001**, *7*, 591–598. [[CrossRef](#)]
57. Wang, M.; Moore, T.R. Carbon, nitrogen, phosphorus, and potassium stoichiometry in an ombrotrophic peatland reflects plant functional type. *Ecosystems* **2014**, *17*, 673–684. [[CrossRef](#)]
58. Ma, J.Z.; Bu, Z.J.; Zheng, X.X.; Ge, J.L.; Wang, S.Z. Effects of shading on relative competitive advantage of three species of *Sphagnum*. *Mires Peat* **2015**, *16*, 1–17.
59. van der Heijden, E.; Verbeek, S.K.; Kuiper, P.J. Elevated atmospheric CO<sub>2</sub> and increased nitrogen deposition: Effects on C and N metabolism and growth of the peat moss *Sphagnum recurvum* P. Beauv. var. *mucronatum* (Russ.) Warnst. *Glob. Change Biol.* **2000**, *6*, 201–212. [[CrossRef](#)]



60. Limpens, J.; Berendse, F. Growth reduction of *Sphagnum magellanicum* subjected to high nitrogen deposition: The role of amino acid nitrogen concentration. *Oecologia* **2003**, *135*, 339–345. [[CrossRef](#)]
61. Reich, P.B.; Oleksyn, J.; Wright, I.J. Leaf phosphorus influences the photosynthesis–nitrogen relation: A cross-biome analysis of 314 species. *Oecologia* **2009**, *160*, 207–212. [[CrossRef](#)]
62. Rochefort, L. *Sphagnum*—A keystone genus in habitat restoration. *The Bryologist* **2000**, *103*, 503–508. [[CrossRef](#)]
63. Waddington, J.M.; Warner, K. Atmospheric CO<sub>2</sub> sequestration in restored mined peatlands. *Ecoscience* **2001**, *8*, 359–368. [[CrossRef](#)]
64. Lucchese, M.; Waddington, J.M.; Poulin, M.; Pouliot, R.; Rochefort, L.; Strack, M. Organic matter accumulation in a restored peatland: Evaluating restoration success. *Ecol. Eng.* **2010**, *36*, 482–488. [[CrossRef](#)]
65. Waddington, J.M.; Lucchese, M.C.; Duval, T.P. *Sphagnum* moss moisture retention following the re-vegetation of degraded peatlands. *Ecohydrology* **2011**, *4*, 359–366. [[CrossRef](#)]
66. Worrall, F.; Chapman, P.; Holden, J.; Evans, C.; Artz, R.; Smith, P.; Grayson, R. *A Review of Current Evidence on Carbon Fluxes and Greenhouse Gas Emissions from UK Peatlands*; JNCC Report: Peterborough, UK, 2011; Available online: [https://nora.nerc.ac.uk/id/eprint/15889/1/jncc442\\_webfinal.pdf](https://nora.nerc.ac.uk/id/eprint/15889/1/jncc442_webfinal.pdf) (accessed on 11 September 2024).

**Disclaimer/Publisher’s Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.