

**Please cite the Published Version**

Turner, IM, Brearley, FQ, Trethowan, LA and Utteridge, TMA (2021) A survey of aluminium accumulation in eumachia (rubiaceae). *Edinburgh Journal of Botany*, 78. pp. 1-10. ISSN 0960-4286

**DOI:** <https://doi.org/10.24823/EJB.2021.335>

**Publisher:** Royal Botanic Garden Edinburgh

**Version:** Published Version

**Downloaded from:** <https://e-space.mmu.ac.uk/636283/>

**Usage rights:**  [Creative Commons: Attribution 4.0](https://creativecommons.org/licenses/by/4.0/)

**Additional Information:** This is an open access article which first appeared in *Edinburgh Journal of Botany*

**Enquiries:**

If you have questions about this document, contact [openresearch@mmu.ac.uk](mailto:openresearch@mmu.ac.uk). Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

## A SURVEY OF ALUMINIUM ACCUMULATION IN *EUMACHIA* (RUBIACEAE)

I. M. Turner<sup>1,2</sup>, F. Q. Brearley<sup>3</sup>, L. A. Trethowan<sup>3,4</sup> & T. M. A. Utteridge<sup>4</sup>

A useful character for spotting specimens of the pantropical Rubiaceae genus *Eumachia* is the leaves drying green. The physiological reasons for this are unresolved. We investigated whether the phenomenon is related to the accumulation of aluminium. Samples of foliage from herbarium specimens of nine species of *Eumachia* (Rubiaceae), including collections from Asia-Pacific, Africa and the Americas, were analysed for elemental concentrations. For comparison, specimens of the closely related genera *Psychotria* and *Palicourea* from similar geographical regions were also analysed. Two species, *Palicourea violacea* (Panama) and *Psychotria pallens* (Brazil) were found to be aluminium hyperaccumulators ( $\geq 1\%$  Al by dry weight), with a third species, *Eumachia collina* (from New Caledonia), falling just short of the threshold with 0.99% Al. Aluminium accumulation ( $\geq 0.4\%$  Al by dry weight) occurs but is infrequent among species of *Eumachia* and appears less common than in the close relative *Palicourea*. Aluminium accumulation also appears patchy in *Psychotria sensu stricto*. It seems unlikely that herbarium greenness in *Eumachia* is directly related to aluminium accumulation.

**Keywords.** Hyperaccumulation, *Palicourea*, Palicoureaeae, *Psychotria*, Psychotrieae, shrubs.

Received 1 October 2019 Accepted 22 September 2020 Published 9 March 2021

### Introduction

The rubiaceous genus *Eumachia*, despite being large (more than 100 species) and almost pantropical (Taylor *et al.*, 2017), is poorly known to tropical botanists. This is readily explained because nearly all the species were formerly included in *Psychotria*, probably the most species-rich genus of woody plants (Davis *et al.*, 2001). Added to this, once plant taxonomists began to separate out this group from *Psychotria* various names, including *Chalaziella* and *Margaritopsis* (Andersson, 2001; Barrabé *et al.*, 2012), were used before *Eumachia* was determined to be the correct one.

Species of *Eumachia* are typically shrubs or small trees with interpetiolar stipules that become hard and straw-coloured and often fragment from the margin, and pyrenes with marginal preformed germination slits (Taylor *et al.*, 2017). Another character that can be useful in determining herbarium material is that in most species of *Eumachia* the leaves

<sup>1</sup> Singapore Botanical Liaison Officer, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, England, UK. E-mail: [i.turner@kew.org](mailto:i.turner@kew.org).

<sup>2</sup> Herbarium, Singapore Botanic Gardens, National Parks Board, 1 Cluny Road, Singapore 259569.

<sup>3</sup> Department of Natural Sciences, Manchester Metropolitan University, Chester Street, Manchester M1 5GD, England, UK.

<sup>4</sup> Identification and Naming, Science Directorate, Royal Botanic Gardens, Kew, Richmond, Surrey TW9 3AB, UK.

dry distinctly green. Although there are species of true *Psychotria* that dry green, most dry various shades of red-brown or grey-brown.

The reason for the unusual colour of dry leaves in *Eumachia* species does not seem to be known. One possibility is that it is related to the accumulation of particular elements in the plant tissues. Given the extensive distribution of the genus and the absence of a common association of the species with unusual geological substrates such as ultramafics, aluminium accumulation on the typically acid and aluminium-rich soils of the lowland wet tropics is considered the most likely explanation. Chenery (1948) noted that leaves drying yellowish green were common among aluminium accumulators. There is a report, based on a semiquantitative chemical assay, of *Eumachia acuífolia* (C.Wright) Delprete & J.H.Kirkbr., a species from Cuba, being an aluminium hyperaccumulator (Jansen *et al.*, 2000, as *Margaritopsis nudiflora* K.Schum.). To investigate this further, we measured elemental concentrations in leaf material from herbarium specimens. To provide comparators, we sought to analyse material from species of *Psychotria sensu stricto*.

## Materials and methods

The material for investigation all came from specimens in the collection of the Royal Botanic Gardens, Kew (K). Samples were chosen by going through the herbarium cupboards searching for *Eumachia* specimens with extra material of mature leaves loose in packets. These were then sampled for destructive analysis of selected elements including aluminium. Species from across the geographical range of the genus were sought. Specimens obviously from upper montane systems were avoided. To provide a comparator for each *Eumachia* specimen, a specimen of a *Psychotria* species from a similar region was also sought for use in the analysis. It transpired that for two of the American *Psychotria* species selected, the herbarium curation had not caught up with recent transfers to *Palicourea*, a large Neotropical genus. Therefore, the comparator taxon to *Eumachia* included species of both *Psychotria* and *Palicourea*. In total 18 samples were analysed. The specimens sampled are listed in the [Appendix](#).

Leaf samples obviously muddy or otherwise exhibiting soil accretion were excluded. It is possible that some of the specimens used may have been treated with mercuric chloride as a prophylactic against insect pests, or contaminated with mercury from treated specimens. This is not likely to influence concentrations of aluminium, however.

The leaf samples (c.100 mg each) were digested in 10 mL of concentrated nitric acid by using a MARSXpress microwave (CEM, Matthews, NC, USA) and made up to 100 mL in ultrapure (18 M $\Omega$ ) deionised water. Total Al, Ca, Cd, Co, Cr, Cu, Fe, K, Mg, Mn, Na, Ni, P and Pb concentrations were quantified using an iCAP 6300 Duo inductively coupled plasma optical emission spectrometer (Thermo-Finnegan, San Jose, CA, USA). Reference material (LGC 7162, Strawberry Leaves) was analysed alongside the samples. These showed good recovery for the certified elemental concentrations, but Al recovery was considerably (62%) lower than the indicative (non-certified) concentration of 1000  $\mu\text{g/g}$ . This may mean that

our results for Al are lower than the true values, but they are likely to be a good reflection of relative concentrations between species.

In a statistical analysis of a large data set, Metali *et al.* (2011) found that a threshold value of 2300–3900 µg Al/g leaf dry mass was required in tropical species to distinguish aluminium accumulators from non-accumulators. To be conservative, we used a minimum value of 4000 µg/g or 0.4% for aluminium accumulators and 10,000 µg/g or 1% for hyperaccumulators.

To visualise the complete data set, a principal components analysis (PCA) was run using the software R (R Core Team, 2020).

## Results

Evidence of aluminium accumulation (> 0.4% by dry weight) was found for four species (Table, Figure 1). These were *Palicourea conephoroides* (Rusby) C.M.Taylor, *Palicourea*

**Table.** Aluminium concentration in the leaves of 18 species from three genera in the Rubiaceae<sup>a</sup>

Species	Location	Aluminium (µg/g leaf dry mass)
<i>Eumachia abrupta</i>	Tanzania	1248
<i>Psychotria leucopoda</i>	Tanzania	66
<i>Eumachia boliviana</i>	Peru	140
<i>Palicourea conephoroides</i>	Peru	5515 <sup>b</sup>
<i>Eumachia cephalantha</i>	Brazil	667
<i>Psychotria pallens</i>	Brazil	29,628 <sup>c</sup>
<i>Eumachia collina</i>	New Caledonia	9910 <sup>b</sup>
<i>Psychotria frondosa</i>	New Caledonia	329
<i>Eumachia damasiana</i>	New Guinea	129
<i>Psychotria beaufortiensis</i>	New Guinea	1141
<i>Eumachia leptothyrsa</i>	New Guinea	174
<i>Psychotria lolokiensis</i>	New Guinea	2016
<i>Eumachia microdon</i>	Panama	66
<i>Palicourea violacea</i>	Panama	33,477 <sup>c</sup>
<i>Eumachia montana</i>	Sumbawa	197
<i>Psychotria</i> sp.	Sumbawa	296
<i>Eumachia obanensis</i>	Cameroon	1368
<i>Psychotria hypsophila</i>	Cameroon	368

<sup>a</sup> For species authorities, see the Appendix.

<sup>b</sup> Above the 4000 µg/g (0.4%) threshold for aluminium accumulation.

<sup>c</sup> Above the 10,000 µg/g (1.0%) threshold for aluminium hyperaccumulation.




Figure 1. Aluminium concentration in the leaves of 18 species from three Rubiaceae genera.


*violacea* (Aubl.) A.Rich. and *Psychotria pallens* Gardner from the Neotropics, and *Eumachia collina* (Labill.) Barrabé, C.M.Taylor & Razafim. from New Caledonia. *Palicourea violacea* and *Psychotria pallens* had Al concentrations well above the threshold for hyperaccumulation, with Al concentration in *Eumachia collina* (0.99%) falling just short.

No consistent pattern emerged from the comparison of *Eumachia* species with *Psychotria/Palicourea* in terms of foliar Al concentration. For the Asia-Pacific and Neotropical pairs, *Eumachia* species generally had lower Al concentrations than *Psychotria/Palicourea*, but for the two African comparators the trend was reversed.

The first two axes of the PCA (Figure 2) accounted for 48% of the variance in the data. The first axis (principal component 1; 32% of variance explained) correlated positively with Al concentration and negatively with the concentration of most of the other elements (see Figure 2B). The second axis (principal component 2; 16% of variance) more effectively discriminated between *Eumachia* and *Psychotria/Palicourea*. This axis opposed foliar concentrations of K, P and Mg with those of Mn, Pb and Al (see Figure 2B). *Eumachia* species generally had a higher score on this axis, with *Psychotria* and *Palicourea* species at the lower end, although there was some overlap (see Figure 2A).

## Discussion

Our sample size was small and unreplicated within species, but some patterns were evident. Of the nine species of *Eumachia* analysed, only one (*Eumachia collina* from New Caledonia) was found to be an aluminium accumulator, although it very nearly qualified as a hyperaccumulator. Although this adds to the previous record of this phenomenon, from



**Figure 2.** Results of principal components analysis: A, biplot of foliar concentrations of 14 elements; B, correlation between the axis score and elemental concentration for 18 species from three Rubiaceae genera. PC, principal component.


---

*Eumachia acuifolia* (Jansen *et al.*, 2000), it is clear that the accumulation of aluminium is not a universal or even frequent phenomenon in *Eumachia*.

The comparison between *Eumachia* and *Psychotria* foliar Al concentration found no consistent differences. One of the seven species of *Psychotria* tested was a clear aluminium hyperaccumulator. Notably, *Psychotria pallens* has foliage that dries a pale green rather than red-brown, which is the general pattern in *Psychotria sensu stricto*. Although only two species of *Palicourea* were analysed, both were found to be aluminium accumulators. One had the highest foliar aluminium concentrations recorded in this study and the other was well over the accumulator threshold. Aluminium hyperaccumulation has been previously reported in species of *Palicourea* (Chenery, 1948; Jansen *et al.*, 2000), but its prevalence in the genus is considerably greater than this, because many of the species listed in these reports have subsequently been transferred to *Palicourea*. So, for instance, of the 24 species of *Psychotria* and *Cephaelis* that Chenery (1948) reported as aluminium hyperaccumulators, only three are now recognised in *Psychotria* and the rest have all been relocated to *Palicourea*.

The taxonomic changes in these Rubiaceae genera have largely been driven by studies of phylogenetic relationships using molecular data, although the morphological overview of Taylor (1996) presaged much of the rearrangement. Phylogenetic analyses have shown that *Psychotria*, as traditionally recognised, contained disparate elements representing various clades (Nepokroeff *et al.*, 1999; Razafimandimbison *et al.*, 2014). The ongoing reorganisation has resulted in considerable change to the systematics of this part of the Rubiaceae. These include the recognition of the tribe Palicoureeae as sister to Psychotrieae (Robbrecht & Manen, 2006), the transfer of much of the heterogeneous *Psychotria* subgenus *Heteropsychotria* to *Palicourea*, and the recognition of *Eumachia* as a large pantropical genus in the Palicoureeae. The currently accepted phylogeny for the genera in this clade of the Rubiaceae is shown in [Figure 3](#).

Aluminium accumulation is widely known in the Rubiaceae (Jansen *et al.*, 2000), although apparently commoner in subfamily Rubioideae than in subfamily Cinchonoideae. Our results, and those of others (Chenery, 1948; Haridasan, 1982; Jansen *et al.*, 2000), confirm the presence of aluminium accumulation in the sister tribes Psychotrieae and Palicoureeae. The prevalence of aluminium accumulation seems mixed among the genera of these tribes. *Palicourea* has many species that are aluminium hyperaccumulators, but its cotribalist *Eumachia* is less frequently found to have tissues with abundant aluminium. Interestingly, *Palicourea* species also tend to have leaves that dry with a greenish hue (Berger, 2018), and this is listed as a character of the Palicoureeae in general (Stevens, [continuously updated](#)). *Palicourea* species often have blue fruits (Taylor & Steyermark, 2004; Berger, 2018), which is a known correlate of aluminium hyperaccumulation (Chenery, 1948), but in *Eumachia* the fruits generally ripen orange-red (Taylor *et al.*, 2017). It seems likely, therefore, that an inherent tolerance to relatively high internal aluminium concentrations is widespread and



**Figure 3.** Phylogenetic relationships of the genera in the sister tribes Psychotrieae and Palicoureeae of the Rubiaceae (after Razafimandimbison et al., 2014).

probably long established in this branch of the Rubiaceae – perhaps unsurprising for a set of genera that are very diverse across the lowland tropics, and that often grow on acid and aluminium-rich soils. It may be that in the Palicoureeae the tendency for leaves to dry green is related to the physiology of mineral uptake and deployment in the tissues, but clearly in *Eumachia* it is not always linked to aluminium accumulation.

Interestingly, the overall stoichiometry of foliar elemental concentrations appeared to show some discrimination at the generic level, with species of *Eumachia* exhibiting relatively high concentrations of the physiologically abundant elements Mg, K and P compared with species of *Psychotria* and *Palicourea*. Two of the species of *Psychotria* also had relatively high foliar Mn concentrations, accentuating the discriminatory power of the second principal component in the PCA. Most *Eumachia* specimens sampled had leaves with a very thin, membranous texture. The *Psychotria/Palicourea* specimens tended to vary from chartaceous to subcoriaceous in leaf texture rather than the tissue-thin laminas of *Eumachia*. A higher cell wall to cytoplasm ratio in the *Psychotria/Palicourea* specimens might be the explanation for the generally lower concentrations of physiologically important elements than in *Eumachia* species.

Clearly, many questions remain unanswered and there is considerable scope for further comparisons within and among widespread tropical genera such as *Eumachia* and *Psychotria*.

### Acknowledgements

We thank David McKendry (MMU) for running the ICP-OES analysis and Martin Cheek (K) for helpful comments on the manuscript.



---

## References

- Andersson L. 2001. *Margaritopsis* (Rubiaceae, Psychotrieae) is a pantropical genus. *Systematics and Geography of Plants*. 71(1):73–85. <https://doi.org/10.2307/3668754>
- Barrabé L, Buerki S, Mouly A, Davis AP, Munzinger J, Maggia L. 2012. Delimitation of the genus *Margaritopsis* (Rubiaceae) in the Asian, Australasian and Pacific region, based on molecular phylogenetic inference and morphology. *Taxon*. 61(6):1251–1268. <https://doi.org/10.1002/tax.616007>
- Berger A. 2018. Synopsis and typification of Mexican and Central American *Palicourea* (Rubiaceae: Palicoureeae), part I: the entomophilous species. *Annalen des Naturhistorischen Museums in Wien. Serie B für Botanik und Zoologie*. 120:59–140. <https://www.jstor.org/stable/26335282>
- Chenery EM. 1948. Aluminium in plants and its relation to plant pigments. *Annals of Botany*. 12(2):121–136. <https://doi.org/10.1093/oxfordjournals.aob.a083177>
- Davis AP, Bridson D, Jarvis C, Govaerts R. 2001. The typification and characterization of the genus *Psychotria* L. (Rubiaceae). *Botanical Journal of the Linnean Society*. 135(1):35–42. <https://doi.org/10.1006/bojl.2000.0390>
- Haridasan M. 1982. Aluminium accumulation by some cerrado native species of central Brazil. *Plants and Soil*. 65:265–273.
- Jansen S, Dessein S, Piesschaert F, Robbrecht E, Smets E. 2000. Aluminium accumulation in leaves of Rubiaceae: systematic and phylogenetic implications. *Annals of Botany*. 85(1):91–101. <https://doi.org/10.1006/anbo.1999.1000>
- Metali F, Salim KA, Burslem DFRP. 2011. Evidence of foliar aluminium accumulation in local, regional and global datasets of wild plants. *New Phytologist*. 193(3):637–649. <https://doi.org/10.1111/j.1469-8137.2011.03965.x>
- Nepokroeff M, Bremer B, Sytsma KJ. 1999. Reorganization of the genus *Psychotria* and tribe Psychotrieae (Rubiaceae) inferred from ITS and rbcL sequence data. *Systematic Botany*. 24(1):5–27. <https://doi.org/10.2307/2419383>
- R Core Team. 2020. R: A language and environment for statistical computing. Vienna: R Foundation for Statistical Computing. <https://www.R-project.org/>
- Razafimandimbison SG, Taylor CM, Wikström N, Pailler T, Khodabandeh A, Bremer B. 2014. Phylogeny and generic limits in the sister tribes Psychotrieae and Palicoureeae (Rubiaceae): evolution of schizocarps in *Psychotria* and origins of bacterial leaf nodules of the Malagasy species. *American Journal of Botany*. 101(7):1102–1126. <https://doi.org/10.3732/ajb.1400076>
- Robbrecht E, Manen JF. 2006. The major evolutionary lineages of the coffee family (Rubiaceae, Angiosperms). Combined analysis (nDNA and cpDNA) to infer the position of *Coptosapelta* and *Luculia*, and supertree construction based on rbcL, rps16, trnL–trnF and atpB–rbcL data. A new classification in two subfamilies, Cinchonoideae and Rubioideae. *Systematics and Geography of Plants*. 76(1):85–145. <http://www.jstor.org/stable/20649700>
- Stevens PF. **Continuously updated**. Angiosperm Phylogeny Website, version 14. <http://www.mobot.org/MOBOT/research/APweb/> [Accessed 11 June 2019.]

- Taylor CM. 1996. Overview of the Psychotrieae (Rubiaceae) in the Neotropics. *Opera Botanica Belgica*. 7:261–270.
- Taylor CM, Steyermark JA. 2004. 63. *Palicourea* Aubl., *Hist. Pl. Guiane* 172, pl. 66. 1775. In: Steyermark JA, Berry PA, Yatskievych K, Holst BK, editors. *Flora of the Venezuelan Guayana*, Volume 8 Poaceae–Rubiaceae. St Louis: Missouri Botanical Garden Press. pp. 680–691.
- Taylor CM, Razafimandimbison SG, Barrabé L, Jardim JG, Barbosa MRV. 2017. *Eumachia* expanded, a pantropical genus distinct from *Psychotria* (Rubiaceae, Palicoureeae). *Candollea*. 72:289–318. <https://doi.org/10.15553/c2017v722a6>

## Appendix

### LIST OF SPECIMENS ANALYSED FOR THE STUDY

All are from the herbarium collection of the Royal Botanic Gardens, Kew (K).

***Eumachia abrupta* (Hiern) J.H.Kirkbr.**

TANZANIA. Uzungwa Mountain NP, stream west of Kilombaro village, 12 xi 1997, P.A. & W.R.Q. Luke 5151.

***Eumachia boliviana* (Standl.) Delprete & J.H.Kirkbr.**

PERU. Coronel Portillo, Chara del Sr Cesar Vela, 25 x 1972, J. Sibunke V. 5461.

***Eumachia cephalantha* (Müll.Arg.) Delprete & J.H.Kirkbr.**

BRAZIL. Sao Paulo: Piracicaba, estrada para Limeira, 14 vii 1993, V.C. Souza et al. 6001.

***Eumachia collina* (Labill.) Barrabé, C.M.Taylor & Razafim.**

NEW CALEDONIA. Exploitation forestiere Guiraud piste en direction de Mt Ades (800–850 m), sol ferallitique sur peridotites, 25 iii 1988, Jaffre 2951.

***Eumachia damasiana* (Sohmer) Barrabé, C.M.Taylor & Razafim.**

PAPUA NEW GUINEA. New Britain: north end of Lake Daketana on Willhaumez Peninsula, 30 v 1973, R.S. Isles et al. NGF 32277.

***Eumachia leptothyrsa* (Miq.) Barrabé, C.M.Taylor & Razafim.**

PAPUA NEW GUINEA. Madang Province: Ramu Subprovince, Walium Station, 20 iv 1979, Sohmer & Katik LAE 75127.

***Eumachia microdon* (DC.) Delprete & J.H.Kirkbr.**

PANAMA. OIS, 16 vi 1924, L.A.M. Riley SERA 149.

***Eumachia montana* (Blume) I.M.Turner**

INDONESIA. Sumbawa: 1927, I. Rensch 593.

***Eumachia obanensis* (Wernham) Razafim. & C.M.Taylor**

CAMEROON. Prov. Sud-Oeust: Dept Fako, Bakingili 50–300 m trail to Mt Cameroon, 25 vii 1984, S.A. Thompson & J.E. Rawlins 1662.

***Palicourea conephoroides* (Rusby) C.M.Taylor**

PERU. Pasco: Oxapampa, Distrito Palcazu, Parque Nacional Yonachagu–Chemillen, Bosque el mirador, 750–820 m, 14 v 2003, A. Monteagudo et al. 5228.

***Palicourea violacea* (Aubl.) A.Rich.**

PANAMA. El Valle de Anter alt. 1000 m, 16 vi 1946, *P.H. Allen* 3536.

***Psychotria beaufortiensis* Valetton ex Sohmer**

INDONESIA. North New Guinea, Cycloop Mountains between Nefor and the coast, vi 1938, *E. Meijer-Drus* 29.

***Psychotria frondosa* S.Moore**

NEW CALEDONIA. Col d'Amien, Mont Pembai 800 m, foret humide schistes, 14 iv 1976, *H.S. McKee* 31039.

***Psychotria hypsophila* K.Schum. & K.Krause**

CAMEROON. Proposed Ebo NP, near the Njuna River, 7 x 2015, *M. Alvarez* 30.

***Psychotria leucopoda* E.M.A.Petit**

TANZANIA. Kwamngumi FR, 350 m, 14 xi 1986, *Ruffo & Mmari* 1980.

***Psychotria lolokiensis* S.Moore**

PAPUA NEW GUINEA. Near south-east side of Little Mt Lowes, c.16 miles north of Port Morseby, 25 iv 1967, *R. Pullen* 6813.

***Psychotria pallens* Gardner**

BRAZIL. Rio de Janeiro, 24 vii 2012, *J.A. Oliveira* et al. 168.

***Psychotria* sp.**

INDONESIA. Sumbawa: Mata, 2 i 1910, *J. Elbert* 4106.