




Please cite the Published Version

Gebrye, Tadesse , Mbada, Chidozie , Apeagyei, Phoebe and Fatoye, Francis  (2025) Prevalence of musculoskeletal disorders among garment workers: a systematic review and meta-analysis. *BMJ Open*, 15 (1). e085123 ISSN 2044-6055

DOI: <https://doi.org/10.1136/bmjopen-2024-085123>

Publisher: BMJ Publishing Group

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/637952/>

Usage rights:  [Creative Commons: Attribution-Noncommercial 4.0](https://creativecommons.org/licenses/by-nc/4.0/)

Additional Information: This is an open access article which first appeared in *BMJ Open*

Data Access Statement: No data are available. All data relevant to the study are included in the article or uploaded as supplementary information. Not applicable.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

BMJ Open Prevalence of musculoskeletal disorders among garment workers: a systematic review and meta-analysis

Tadesse Gebrye,¹ Chidozie Mbada,¹ Phoebe Apeagyei,² Francis Fatoye ^{1,3}

To cite: Gebrye T, Mbada C, Apeagyei P, *et al.* Prevalence of musculoskeletal disorders among garment workers: a systematic review and meta-analysis. *BMJ Open* 2025;**15**:e085123. doi:10.1136/bmjopen-2024-085123

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-085123>).

Received 06 February 2024
Accepted 10 January 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

¹Department of Health Professions, Manchester Metropolitan University, Manchester, UK

²Manchester Metropolitan University, Manchester, UK

³Lifestyle Diseases, Faculty of Health Sciences, North-West University, Potchefstroom, South Africa

Correspondence to

Tadesse Gebrye;
t.gebrye@mmu.ac.uk

ABSTRACT

Background Garment workers are at high risk of musculoskeletal disorders (MSDs) due to repetitive physical tasks, long working hours and varying workstations. As there is no existing epidemiological overview of MSDs among garment workers, this systematic review aimed to evaluate the global evidence on prevalence of MSDs in this population.

Methods A systematic review of the literature was conducted in accordance with Preferred Reporting Items for Systematic Reviews and Meta-Analyses guidelines. Medline, Cumulative Index to Nursing and Allied Health Literature, PubMed, Scopus and Web of Science electronic databases were searched to identify studies published in English up to November 2022. Random-effects meta-analysis was used to estimate pooled prevalence.

Results The search yielded 258 published studies, of which 14 were deemed relevant and were included in this review. The included studies reported prevalence from India (n=3), Bangladesh (n=2), Ethiopia (n=2), Thailand (n=2), Botswana (n=1), Iran (n=1), Sri Lanka (n=1), Cambodia (n=1) and Denmark (n=1) and involved 15 029 garment workers. Most of the included studies (79%) were assessed to be methodologically sound (low risk of bias). The mean age of participants ranged from 24.2 to 40 years. The prevalence of MSDs ranged from 15.5% to 92%. The pooled prevalence of MSDs from nine studies was 65.6% (95% CI 44.5% to 51.9%). Low back pain and neck pain were reported as the common MSDs in the included studies.

Conclusions The findings highlight the considerable risk of MSDs, especially low back and neck pain, linked to repetitive tasks, extended hours and inconsistent workstations. Given the heavy toll of MSDs on this workforce, targeted interventions and ergonomic improvements are crucial to mitigate the risks and improve garment workers' well-being.

INTRODUCTION

Work-related musculoskeletal disorders (WRMSDs) refer to disabling injuries to the muscles, tendons or nerves caused by work activities.¹ WRMSDs are the leading cause of work absence, indicating that the management of risks related to musculoskeletal (MSK) disorders in the workforce may not be adequate.² According to the Global Burden of Disease study, MSK conditions are the

STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ To our knowledge, this is the first systematic review summarising the prevalence of musculoskeletal disorders (MSDs) among garment workers.
- ⇒ We extracted prevalence data from 14 studies conducted across 9 different countries.
- ⇒ The current study used methodology that is clearly outlined, allowing for replication and transparency in the selection and analysis of data.
- ⇒ The use of varied questionnaires across the included studies to assess MSDs may lead to inconsistencies in measurement, limiting the generalisability of the results.

second-highest contributor (20%–33%) to global disability among those affected by such conditions.³ Among working-age individuals, MSK disorders are one of the most common causes of years lived with disability.⁴ Common risk factors for WRMSDs include awkward postures, prolonged static work, repetitive movements, manual material handling, forceful exertions and vibration.^{5 6} Major psychosocial factors, such as job dissatisfaction, work-related stress and time pressure, also play a significant role in the development of WRMSDs.⁷ Individuals affected by WRMSDs face a risk of losing income, which can lead to an increased risk of poverty.⁸ On average, 4% of the total gross national product across all countries is lost annually due to compensation, medical expenses, lost productivity and insurance costs related to work-related diseases and injuries.⁹

Workers in the garment industry are highly susceptible to WRMSDs,¹⁰ with factors such as perceived physical and psychological demands contributing to their development, as shown in a study on weaving operations.¹¹ This sector, which is labour-intensive and highly competitive globally, is crucial for economies, particularly in countries like India, where it is the second-largest industry, contributing significantly to gross domestic product, exports and employment,

Table 1 Overview of all studies included in systematic review

Study author	Country	Study design	Data source	Sample size	Male (n)	Age range (mean)	Type of MSDs/ exposure	Risk of bias
Zelee <i>et al</i> ²⁰	Ethiopia	Cross-sectional	Clinics (integrated textile factories)	7992	40%	40 years, 10 (18–69)	Not reported	Low
Sealetsa and Thatcher ²⁸	Botswana	A modified Corlett and Bishop body map questionnaire and the NASA TLX	Questionnaire	157	0%	32.25 years (SD=7.00)	Lower back, upper back and midback, shoulders, neck, and legs.	Moderate
Pal <i>et al</i> ¹⁴	India	Cross-sectional study	Face-to-face interview	222	70.22%	36–55 (42.34%)	Not reported	Low
Kebede Deyyas and Tafese ²¹	Ethiopia	Cross-sectional study	Face-to-face interview	422	12.30%	26.9 (SD=7.2)	Elbow and wrist musculoskeletal disorders	Low
Shazzad <i>et al</i> ²⁴	Bangladesh	Cross-sectional study	Questionnaire	350	17.10%	N/A	Shoulder, lower back, neck and knee	Low
Hossain <i>et al</i> ²⁵	Bangladesh	Cross-sectional study	A structured questionnaire	232	19%	31.3 years (SD=7)	Lower back pain, neck pain, neck pain, knee pain	Low
Chavavitsakulchai and Shahnava ²⁶	Thailand	Survey	Interviews based on Standardised Nordic Questionnaires	400	12%	24.2 years (SD=5)	Neck, upper back, lower back, and knees, ankles/feet, shoulders, elbows, hips/thighs	Moderate
Veisi <i>et al</i> ²⁹	Iran	Cross-sectional study	The Nordic Musculoskeletal Questionnaire	586	41.90%	Age range 16–72 years, 35.7 years (SD=11.8)	Shoulders, wrists and arms, elbows, hand	Low
Lombardo <i>et al</i> ³⁰	Sri Lanka	Cross-sectional study	Questionnaire	1058	0%	27.8 years±7.3	Back pain, knees, shoulders, hand and wrist, neck, and forearm and elbow.	Low
Van <i>et al</i> ³¹	Cambodia	Cross-sectional descriptive study	Face-to-face interviews and direct observation	702	10.70%	27.3 years (SD=7.45)	Neck, shoulder, upper back, lower back	Low
Andersen and Gaardboe ³²	Denmark	Cross-sectional study	Questionnaires	424	37.3 years (SD=8.8)		Neck pain, shoulder pain, elbow pain, forearm and/or wrist pain, hand and/or finger pain	Low

Continued

Table 1 Continued

Study author	Country	Study design	Data source	Sample size	Male (n)	Age range (mean)	Type of MSDs/ exposure	Risk of bias
Mokhasi ²²	India	Prospective observational study	The Nordic questionnaire:	430	0%	29.36 years±6.96	Neck pain, shoulder, elbow, wrist, upper back, lower back, foot/ ankle, knees.	Moderate
Homsombat and Chaiklieng ²⁷	Thailand	Cross-sectional study	The Standardise Nordic Questionnaire	1674	0%	Ranged between 30 and 60 years old	Low back, neck pain, elbow	Low
Ravichandran et al ²³	India (Tamil Nadu)	Cross sectional study	The Standardise Nordic Questionnaire	380	60.80%	30.53	Neck pain, knee and low back pain	Low

MSDs, musculoskeletal disorders; NASA TLX, The National Aeronautics and Space Administration's (NASA) Task Load Index (TLX).




Figure 1 Flow diagram of publications included and excluded in the review.

including for millions of women and rural workers.^{12–14} In Bangladesh, the garment sector accounts for 84% of exports,¹⁵ while countries such as India and Cambodia employ millions of people in this industry,¹⁴ with the garment sector representing a significant portion of the labour force.¹⁴ The industry has also seen rapid growth in nations like Brazil, China, India and Vietnam, where the workforce has expanded substantially.¹⁶

Understanding the potential risk factors for WRMSDs is essential for quality improvement and reducing the incidence of these disorders among garment workers. Numerous studies have explored the prevalence of MSK disorders in this population. Consequently, we conducted a systematic review of studies reporting the prevalence of MSK disorders across various countries and time periods, with the goal of providing updated insights into the prevalence of these conditions among garment workers. We aggregated prevalence estimates from different regions and countries and analysed the prevalence of MSK disorders.

METHODS

The study was designed as a systematic review and meta-analysis of published literature on the prevalence of MSK disorders among garment workers. It was performed and reported following the Preferred Reporting Items for Systematic Reviews and Meta-Analyses checklist for systematic reviews of intervention.¹⁷ A protocol for this systematic review was prospectively registered on PROSPERO and can be found at https://www.crd.york.ac.uk/PROSPERO/display_CRD42022368808.

Search strategy

The authors conducted a search of several databases, including Ovid MEDLINE, CINAHL, PubMed, Scopus and Web of Science, for studies published from their inception up to 30 November 2022, that reported on the prevalence of MSDs among garment workers. The search

terms included combinations of keywords such as musculoskeletal conditions, MSDs, tailor, seamstress, garment worker, designer, pattern cutter, prevalence and epidemiology (online supplemental appendix 1). These terms were linked using conjunctions like “AND” and “OR.” The search was limited to articles published in English. Additionally, the authors reviewed all relevant references and related systematic reviews from various regions.

Inclusion and exclusion criteria

Studies were deemed eligible for inclusion if they met the following criteria: the population studied consisted of garment workers (aged 18 years and older), and the focus was on musculoskeletal (MSK) disorders related to garment fit. The review included all studies published in English, irrespective of study design, publication status or publication date. We excluded reviews, editorials, conference papers, case reports or series and animal studies. Three independent reviewers (TG, PA and MCE) initially assessed the titles and abstracts, followed by the full texts of the studies to determine if they met the inclusion criteria. In cases of disagreement, the final decision was made by a fourth reviewer (FF).

Study selection and assessment of methodological quality

After removing duplicates, one reviewer (TG) screened all titles, abstracts and full-text articles, while a sample of these was checked by a second reviewer (PA and MCE). Any discrepancies were resolved through discussion and consensus with a fourth reviewer (FF). The full texts of the identified studies were then assessed against the inclusion and exclusion criteria.

The risk of bias in the included studies was evaluated using a previously developed risk of bias tool.¹⁸ This assessment tool includes 10 items that address both external and internal validity (online supplemental appendix 2). Each study's overall risk of bias was classified as low, moderate or high. Studies receiving 9 or 10 ‘yes’ answers were considered to have a low risk of bias; those with 7 or 8 ‘yes’ answers were considered to have a moderate risk of bias, and studies with 6 or fewer ‘yes’ answers were considered to have a high risk of bias.

Data extraction and analysis

The data were extracted into a standardised Excel spreadsheet, which included study variables such as the first author's name, year of publication, country, study design, data source, sample size, sex, age range (mean), response rate, types of MSDs, overall prevalence, prevalence in males and prevalence in females. All data were double-checked by other researchers to ensure accuracy.

A descriptive and statistical analysis of the extracted results was conducted. The prevalence for each study was calculated using Comprehensive Meta-analysis software (Biostat, New Jersey, USA), V.3 for Windows. Pooled prevalence estimates and 95% CIs were computed using a random-effects meta-analysis model, which is robust and appropriate when there is significant heterogeneity

in prevalence across studies.¹⁹ The prevalence of MSDs among garment workers and the total number of patients in each study population were used as variables. Weighted events were reported with 95% CIs, and heterogeneity was assessed using I^2 values.

Patient and public involvement

Patients and the general public were not involved in this study.

RESULTS

Identification and description of studies

The literature search identified 258 citations (CINHAL and MEDLINE=62, PubMed=102, Scopus=24, Web of Science=70). After removing 27 duplicates, 167 studies were excluded following title and abstract screening. A total of 64 studies were deemed potentially relevant and underwent full-text evaluation. Ultimately, 14 articles met the inclusion criteria and provided prevalence data on MSDs among garment workers. Of these 14 studies, 11 were assessed as having a low risk of bias, while the remaining 3 were classified as having a moderate risk of bias (table 1). The flow chart illustrating the literature selection process is presented in figure 1.

Characteristics of the included studies

A total of 14 studies reporting data on the prevalence of MSDs among garment workers were included (table 1). These studies enrolled 15 029 participants, with sample sizes ranging from 157 to 7992 individuals. The mean age of participants varied from 24.2 to 40 years. The studies encompassed a broad range of demographic characteristics and were published between 1993 and 2022. The studies were conducted in Ethiopia,^{20 21} India,^{14 22 23} Bangladesh,^{24 25} Thailand,^{26 27} Botswana,²⁸ Iran,²⁹ Sri Lanka,³⁰ Cambodia³¹ and Denmark.³² One study employed a prospective observational method,²² while the others used a cross-sectional design.

Prevalence

The prevalence of MSDs among garment workers ranged from 15.5% to 92% (table 2). Sri Lanka reported the highest prevalence, while Cambodia had the lowest. Two studies^{25 29} found that the proportion of females with MSDs was higher than that of males. For instance, one study reported that the prevalence of MSDs in females was approximately 10% higher than in males.²⁹ In the random effects meta-analysis (n=9), the pooled prevalence of MSDs was 0.656 (95% CI 0.445 to 0.819) (figure 2).

DISCUSSION

The present study aimed to consolidate evidence on the prevalence of MSKs among garment workers. The findings revealed a high prevalence of MSK disorders, ranging from 15.5% to 92%. The studies included in the analysis provided prevalence data from Ethiopia, India, Bangladesh, Thailand, Botswana, Iran, Sri Lanka, Cambodia and

Denmark. Overall, the studies were assessed to have a low risk of bias.

Consistent with our study, a review of health issues among garment factory workers identified MSDs as one of the major health concerns.³³ MSK problems were the most prevalent among all the conditions examined, with a prevalence of 78.89%. Additionally, the mean 12-month prevalence of work-related MSDs in Europe's secondary industries was found to be high, with back pain (60%), neck pain (51%) and shoulder pain (50%) reported.³⁴ Overall, the results suggest that factory workers, particularly those in garment factories, are highly exposed to MSK problems, which are the most common health issues among them.

Evidence shows that 60% of women working as sewing machine operators suffer from back pain.³⁵ A study conducted among ready-made garment workers in Bangladesh revealed that the 12-month prevalence of MSDs was highest in the lower back for women (24.7%), followed by the neck (23.7%) and knees (17.7%). For men, the highest prevalence of MSDs was in the neck (21.7%), followed by the knees (13%), lower back (13%) and upper back (10.9%).²⁵ A study investigating the 1-year prevalence of spinal pain and its consequences—such as pain, function, sick leave and healthcare use—found gender differences in sick leave patterns and healthcare utilisation. It also noted that a small proportion of individuals with pain used a significant amount of healthcare

Table 2 Reported prevalence (%) of MSD among garment workers

Study author	Prevalence total (95% CI)	Prevalence in males (95% CI)	Prevalence in females (95% CI)
Zele <i>et al</i> ²⁰	29%	N/A	N/A
Sealetsa and Thatcher ²⁸	37%	N/A	N/A
Pal <i>et al</i> ¹⁴	71%	N/A	N/A
Kebede Deyyas and Tafese ²¹	Elbow (40%) Wrist (37.7%)	N/A	N/A
Shazzad <i>et al</i> ²⁴	77.1% (shoulder=17.9%, lower back=15.2%, neck=13.8%) and knee=10.8%).	N/A	N/A
Hossain <i>et al</i> ²⁵	N/A	Neck=21.7%, Knees=13%), Lower back=13%	Lower back=24.7%, Neck=23.7% and knees=17.7%
Chavalitsakulchai and Shahnava ²⁶	50%	N/A	N/A
Veisi <i>et al</i> ²⁹	87.7%	82,1%	91.60%
Lombardo <i>et al</i> ³⁰	15.5% (back pain=57.3%, knees=31.7%, shoulders=9.1%, hand and wrist=7.3%, neck=6.7%, and forearm and elbow=3.0%).	N/A	N/A
Van <i>et al</i> ³¹	92% (95% CI=90.0% to 94.0%)	N/A	N/A
Andersen and Gaardboe ³²	Neck pain=26.6%, shoulder pain=25.2% and elbow pain=4.5%	N/A	N/A
Mokhasi ²²	Lower back (70%), followed by the upper back (40%)	N/A	N/A
Homsombat and Chaiklieng ²⁷	Without severity: lower back=41.75%, shoulders=41.21%, neck=41.10% and elbows=40.80% Considering severity: low backs=61.47%, neck=61.23% and thigh=58.06%	N/A	N/A
Ravichandran <i>et al</i> ²³	77.6% (neck=32.1%, knee=28.7% and low back pain=26.6%)	N/A	N/A
N/A, not available.			




Figure 2 Forest plot of prevalence of MSDs among garment workers. MSDs, musculoskeletal disorders.

resources.³⁶ One contributing factor may be that women often perform dual roles, such as childcare, which can lead to increased stress and psychological problems.³³

Biomechanics plays a crucial role in the prevention and management of WRMSDs in the garment industry.³⁷ It is the scientific field that applies mechanical principles to biological systems.³⁸ MSDs are strongly associated with factors such as peak hand force, the percentage of time spent in forceful hand exertion and the rate of forceful hand repetitions.³⁹ Studies consistently indicate that repetitive movements are a key risk factor for shoulder pain.⁴⁰ Additionally, working in static postures for extended periods with minimal or no breaks has been identified as a significant cause of neck pain among garment workers.^{22,32} Further evidence also shows that the incidence of WRMSDs increases with the age of workers.⁴¹ Therefore, a deeper understanding of tissue tolerance under varying loading conditions, along with proper workstation design, may help reduce garment workers' vulnerability to MSDs.

This review has both strengths and potential limitations. A systematic and rigorous approach was used to identify relevant studies on the prevalence of MSDs among garment workers. However, the use of different questionnaires across the included studies to investigate MSDs may limit the generalisability of the results. Diagnosing MSDs in various settings and countries can be challenging due to the adoption of specific surgical procedures, and the prevalence data may be based on self-reported measures.⁴² Additionally, since only English-language publications were reviewed, relevant literature in other languages may have been overlooked. Except for

two studies,^{25,29} the results were not reported by gender. Despite these limitations, this is the first systematic review on this topic to provide a meta-analysis.

CONCLUSIONS

MSDs are highly prevalent among garment workers, particularly affecting the low back and neck. This prevalence significantly impacts health service utilisation. Therefore, it is essential to gain insights and develop targeted programmes to prevent MSDs among garment workers. By effectively summarising the prevalence of MSDs, the findings from this study could play a key role in shaping decisions related to workstations and in addressing the physical challenges and risks faced by garment workers.

Contributors TG and FF contributed to the conception and design of the study. TG, MCE and PA made substantial contributions to literature search, data collection, statistical analysis and the original drafting and revising of the manuscript. FF and PA provided professional comments on the manuscript. FF is responsible for the overall content as guarantor. All authors read and approved of the final manuscript for publication.

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient and public involvement Patients and/or the public were not involved in the design, or conduct, or reporting, or dissemination plans of this research.

Patient consent for publication Not applicable.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement No data are available. All data relevant to the study are included in the article or uploaded as supplementary information. Not applicable.

Supplemental material This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

Open access This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

ORCID iD

Francis Fatoye <http://orcid.org/0000-0001-7976-2013>

REFERENCES

- Forsman M, Fan X, Rhén I-M, *et al*. Concerning a Work Movement Velocity Action Level Proposed in 'Action Levels for the Prevention of Work-Related Musculoskeletal Disorders in the Neck and Upper Extremities: A Proposal' by Inger Arvidsson *et al*. (2021). *Ann Work Expo Health* 2022;66:130–1.
- Crawford JO, Berkovic D, Erwin J, *et al*. Musculoskeletal health in the workplace. *Best Pract Res Clin Rheumatol* 2020;34:101558.
- James SL, Abate D, Abate KH. Global, regional, and national incidence, prevalence, and years lived with disability for 354 diseases and injuries for 195 countries and territories, 1990–2017: a systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;392:1789–858.
- Safiri S, Kolahi A-A, Hoy D, *et al*. Global, regional, and national burden of neck pain in the general population, 1990–2017: systematic analysis of the Global Burden of Disease Study 2017. *BMJ* 2020;368:m791.
- Sim J, Lacey RJ, Lewis M. The impact of workplace risk factors on the occurrence of neck and upper limb pain: a general population study. *BMC Public Health* 2006;6:1–0.
- Choobineh A, Tabatabaei SH, Tozihian M, *et al*. Musculoskeletal problems among workers of an Iranian communication company. *Indian J Occup Environ Med* 2007;11:32–6.
- David G, Woods V, Li G, *et al*. The development of the Quick Exposure Check (QEC) for assessing exposure to risk factors for work-related musculoskeletal disorders. *Appl Ergon* 2008;39:57–69.
- Parvin F, Rafiquzzaman M, Parvez MS, *et al*. Ergonomic compatibility between sewing workstation and Bangladeshi apparel workers. *Int J Occup Saf Ergon* 2024;27:1–16.
- Takala J, Niu S. Responses to the equity challenge in safety and health at work: improvement of working conditions in equitable bases. 27th International Congress on Occupational Health; 2003:1–12.
- International Labour Organization. ILO introductory report: global trends and challenges on occupational safety and health. 2011.
- Telaprolu N, Anne SD. Physical and psychological work demands as potential risk factors for musculoskeletal disorders among workers in weaving operations. *Indian J Occup Environ Med* 2014;18:129–34.
- Hamja A, Maalouf M, Hasle P. The effect of lean on occupational health and safety and productivity in the garment industry – a literature review. *Prod Manuf Res* 2019;7:316–34.
- Taplin IM, Winterton J, Winterton R. Understanding Labour Turnover in a Labour Intensive Industry: Evidence from the British Clothing Industry*. *J Management Studies* 2003;40:1021–46.
- Pai A, Dasgupta A, Sadhukhan SK, *et al*. How common are aches and pains among garment factory workers? A work-related musculoskeletal disorder assessment study in three factories of south 24 Parganas district, West Bengal. *J Family Med Prim Care* 2021;10:917–21.
- Berg A, Chhaparia H, Hedrich S, *et al*. What's next for Bangladesh's garment industry, after a decade of growth, 6. McKinsey & Company; 2021.1–7.
- Nordas HK. *The global textiles and clothing industry post agreement on textile and clothing. discussion paper (5)*. Geneva: World Trade Organisation, 2023.
- Page MJ, McKenzie JE, Bossuyt PM, *et al*. Updating guidance for reporting systematic reviews: development of the PRISMA 2020 statement. *J Clin Epidemiol* 2021;134:103–12.
- Hoy D, Brooks P, Woolf A, *et al*. Assessing risk of bias in prevalence studies: modification of an existing tool and evidence of interrater agreement. *J Clin Epidemiol* 2012;65:934–9.
- Stroup DF, Berlin JA, Morton SC, *et al*. Meta-analysis of observational studies in epidemiology: a proposal for reporting. *JAMA* 2000;283:2008–12.
- Zelev Y, Kumie A, Deressa W, *et al*. Registered health problems and demographic profile of integrated textile factory workers in Ethiopia: a cross-sectional study. *BMC Public Health* 2021;21:1526.
- Kebede Deyyas W, Tafese A. Environmental and organizational factors associated with elbow/forearm and hand/wrist disorder among sewing machine operators of garment industry in Ethiopia. *J Environ Public Health* 2014;2014:732731.
- Mokhasi VR. Fore-Warred Is Fore-Armed: Effect of Musculoskeletal Disorders on Sickness Absenteeism. *Cureus* 2022;14:e30481.
- Ravichandran SP, Shah PB, Lakshminarayanan K, *et al*. Musculoskeletal problems among workers in a garment industry, at Tirupur, Tamil Nadu. *Indian J Community Health* 2016;28:269–74.
- Shazzad MN, Ahmed S, Haq SA, *et al*. Musculoskeletal symptoms and disorders among 350 garment workers in Bangladesh: A cross-sectional pilot study. *Int J Rheum Dis* 2018;21:2063–70.
- Hossain MD, Aftab A, Al Imam MH, *et al*. Prevalence of work related musculoskeletal disorders (WMSDs) and ergonomic risk assessment among readymade garment workers of Bangladesh: A cross sectional study. *PLoS One* 2018;13:e0200122.
- Chavalitsakulchai P, Shahnavaz H. Musculoskeletal disorders of female workers and ergonomics problems in five different industries of a developing country. *J Hum Ergol (Tokyo)* 1993;22:29–43.
- Homsombat T, Chaiklieng S. Physical Fitness and Muscular Discomfort among Informal Garment Female Workers in Udonthani Province, Thailand. *J Med Assoc Thai* 2017;100:230–8.
- Sealetsa OJ, Thatcher A. Ergonomics issues among sewing machine operators in the textile manufacturing industry in Botswana. *Work* 2011;38:279–89.
- Veisi H, Choobineh A, Ghaem H, *et al*. Upper extremity musculoskeletal symptoms among Iranian hand-woven shoe workers. *Work* 2020;67:129–39.
- Lombardo SR, Vijitha de Silva P, Lipscomb HJ, *et al*. Musculoskeletal symptoms among female garment factory workers in Sri Lanka. *Int J Occup Environ Health* 2012;18:210–9.
- Van L, Chaiear N, Sumananont C, *et al*. Prevalence of musculoskeletal symptoms among garment workers in Kandal province, Cambodia. *J Occup Health* 2016;58:107–17.
- Andersen JH, Gaardboe O. Prevalence of persistent neck and upper limb pain in a historical cohort of sewing machine operators. *Am J Ind Med* 1993;24:677–87.
- Lillypet S, Jain T, Joseph B, *et al*. Health problems among garment workers: A narrative literature review. *JOHE* 2017;6:114–21.
- Govaerts R, Tassignon B, Ghillebert J, *et al*. Prevalence and incidence of work-related musculoskeletal disorders in secondary industries of 21st century Europe: a systematic review and meta-analysis. *BMC Musculoskelet Disord* 2021;22:751.
- Roy S. Garments industry in India: some reflections on size distribution of firms. Working paper 05/2010, 34. Institute for Human Development; 2010.1–16.
- Linton SJ, Hellsing AL, Haldén K. A population-based study of spinal pain among 35–45-year-old individuals. Prevalence, sick leave, and health care use. *Spine (Phila Pa 1976)* 1998;23:1457–63.
- Das S, Krishna Moorthy M, Shanmugaraja K. Analysis of Musculoskeletal Disorder Risk in Cotton Garment Industry Workers. *J Natural Fibers* 2023;20:1–10.
- Gitajn IL, Rodriguez EK. Biomechanics of musculoskeletal injury. In: *Biomechanics in applications*. IntechOpen, 2011.
- Harris-Adamson C, Eisen EA, Kapellusch J, *et al*. Biomechanical risk factors for carpal tunnel syndrome: a pooled study of 2474 workers. *Occup Environ Med* 2015;72:33–41.
- van der Windt DA, Thomas E, Pope DP, *et al*. Occupational risk factors for shoulder pain: a systematic review. *Occup Environ Med* 2000;57:433–42.
- Berberoglu U, Tokuc B. Work-related musculoskeletal disorders at two textile factories in Edirne, Turkey. *Balkan Med J* 2013;1:23–7.
- Carotti M, Salaffi F, Beci G, *et al*. The application of dual-energy computed tomography in the diagnosis of musculoskeletal disorders: a review of current concepts and applications. *Radiol Med* 2019;124:1175–83.