


Please cite the Published Version

Jafarian, Fahimeh-Sadat, Jafari-Harandi, Mahmonir, Yeowell, Gillian  and Sadeghi-Demneh, Ebrahim (2025) Comparative efficacy of lumbar and pelvic support on pain, disability, and motor control in women with postpartum pelvic girdle pain: a three-armed randomized controlled trial. BMC Musculoskeletal Disorders, 26 (1). 100 ISSN 1471-2474

DOI: <https://doi.org/10.1186/s12891-025-08350-3>

Publisher: BioMed Central

Version: Supplemental Material

Downloaded from: <https://e-space.mmu.ac.uk/638255/>

Usage rights:  [Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0](https://creativecommons.org/licenses/by-nc-nd/4.0/)

Additional Information: This is an open access article which first appeared in BMC Musculoskeletal Disorders

Data Access Statement: The datasets used and analysed during the current study are available from the corresponding author (E.S-D) at sadeghi@rehab.mui.ac.ir. upon reasonable request.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)

The TIDieR (Template for Intervention Description and Replication) Checklist*:

Information to include when describing an intervention and the location of the information

Item number	Item	Where located **	
		Primary paper (page or appendix number)	Other † (details)
1.	BRIEF NAME Provide the name or a phrase that describes the intervention.	8	_____
2.	WHY Describe any rationale, theory, or goal of the elements essential to the intervention.	8	_____
3.	WHAT Materials: Describe any physical or informational materials used in the intervention, including those provided to participants or used in intervention delivery or in training of intervention providers. Provide information on where the materials can be accessed (e.g. online appendix, URL).	8	_____
4.	Procedures: Describe each of the procedures, activities, and/or processes used in the intervention, including any enabling or support activities.	8	_____
5.	WHO PROVIDED For each category of intervention provider (e.g. psychologist, nursing assistant), describe their expertise, background and any specific training given.	8	_____
6.	HOW Describe the modes of delivery (e.g. face-to-face or by some other mechanism, such as internet or telephone) of the intervention and whether it was provided individually or in a group.	8	_____
7.	WHERE Describe the type(s) of location(s) where the intervention occurred, including any necessary infrastructure or relevant features.	8	_____

	WHEN and HOW MUCH		
8.	Describe the number of times the intervention was delivered and over what period of time including the number of sessions, their schedule, and their duration, intensity or dose.	8	_____
	TAILORING		
9.	If the intervention was planned to be personalised, titrated or adapted, then describe what, why, when, and how.	8	_____
	MODIFICATIONS		
10.†	If the intervention was modified during the course of the study, describe the changes (what, why, when, and how).	N/A	_____
	HOW WELL		
11.	Planned: If intervention adherence or fidelity was assessed, describe how and by whom, and if any strategies were used to maintain or improve fidelity, describe them.	N/A	_____
12.‡	Actual: If intervention adherence or fidelity was assessed, describe the extent to which the intervention was delivered as planned.	N/A	_____

** **Authors** - use N/A if an item is not applicable for the intervention being described. **Reviewers** – use ‘?’ if information about the element is not reported/not sufficiently reported.

† If the information is not provided in the primary paper, give details of where this information is available. This may include locations such as a published protocol or other published papers (provide citation details) or a website (provide the URL).

‡ If completing the TIDieR checklist for a protocol, these items are not relevant to the protocol and cannot be described until the study is complete.

* We strongly recommend using this checklist in conjunction with the TIDieR guide (see *BMJ* 2014;348:g1687) which contains an explanation and elaboration for each item.

* The focus of TIDieR is on reporting details of the intervention elements (and where relevant, comparison elements) of a study. Other elements and methodological features of studies are covered by other reporting statements and checklists and have not been duplicated as part of the TIDieR checklist. When a **randomised trial** is being reported, the TIDieR checklist should be used in conjunction with the CONSORT statement (see www.consort-statement.org) as an extension of **Item 5 of the CONSORT 2010 Statement**. When a **clinical trial protocol** is being reported, the TIDieR checklist should be used in conjunction with the SPIRIT statement as an extension of **Item 11 of the SPIRIT 2013 Statement** (see www.spirit-statement.org). For alternate study designs, TIDieR can be used in conjunction with the appropriate checklist for that study design (see www.equator-network.org).