







Please cite the Published Version

Hacket-Pain, Andrew , Szymkowiak, Jakub , Journé, Valentin , Barczyk, Maciej K., Thomas, Peter A., Lageard, Jonathan , Kelly, Dave  and Bogdziewicz, Michał  (2025) Growth decline in European beech associated with temperature-driven increase in reproductive allocation. Proceedings of the National Academy of Sciences, 122 (5). e2423181122 ISSN 0027-8424

DOI: <https://doi.org/10.1073/pnas.2423181122>

Publisher: National Academy of Sciences

Version: Published Version

Downloaded from: <https://e-space.mmu.ac.uk/638328/>

Usage rights:  Creative Commons: Attribution-Noncommercial-No Derivative Works 4.0

Additional Information: This is an open access article published in Proceedings of the National Academy of Sciences.

Data Access Statement: Datasets data have been deposited in OSF <https://osf.io/eujt7/> (68).

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from <https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines>)



Growth decline in European beech associated with temperature-driven increase in reproductive allocation

Andrew Hacket-Pain^{a,1} , Jakub Szymkowiak^{b,c} , Valentin Journé^c , Maciej K. Barczyk^c, Peter A. Thomas^d, Jonathan G. A. Lageard^e, Dave Kelly^f , and Michał Bogdziewicz^{c,1}

Affiliations are included on p. 6.

Edited by Donald Ort, University of Illinois at Urbana Champaign, Urbana, IL; received November 13, 2024; accepted December 16, 2024

Climate change is impacting forests in complex ways, with indirect effects arising from interactions between tree growth and reproduction often overlooked. Our 43-y study of European beech (*Fagus sylvatica*) showed that rising summer temperatures since 2005 have led to more frequent seed production events. This shift increases reproductive effort but depletes the trees' stored resources due to insufficient recovery periods between seed crops. Consequently, annual tree ring increments have declined by 28%, dropping from a stable average of 1.60 mm y⁻¹ between 1980 and 2005 to 1.16 mm y⁻¹ thereafter. Importantly, this growth decline occurred without an accompanying trend in summer drought, indicating that altered reproductive patterns—not moisture stress—are driving the reduction. This creates a “perfect storm”: Increased reproductive effort drains resources, viable seed output falls due to the loss of mast-seeding benefits via pollination and lower seed predation, and the ongoing growth decline reduces current carbon uptake and future reproductive potential. These compounding factors threaten the sustainability of Europe's most widespread forest tree. Our findings unveil a critical yet underrecognized indirect mechanism by which climate change endangers forest ecosystems, emphasizing the need to consider interactions between demographic processes when assessing species vulnerability to climate change.

climate change | tree demography | tree growth | tree rings | mast seeding

The influence of climate change on global forest demographics is now evident, often characterized by increased mortality rates (1, 2), shifts in growth rates (3), and changes in fecundity (4, 5). While the direct impacts of climate on these metrics have been extensively examined, the interaction among demographic rates suggest the possibility of indirect effects (6, 7). For instance, mortality rates can increase when extreme weather events coincide with reproductive phases that deplete plant reserves (8, 9). This is particularly pertinent for species that mast, i.e., forgo regular reproduction in favor of concentrating seed production into sporadic, large-scale events, resulting in strong resource depletion (8, 10, 11). For example, in the tree *Distylium lepidotum*, masting depletes stored starch, resulting in slower regrowth or increased susceptibility to dieback following drought conditions (12). Consequently, alterations in reproductive patterns due to climate change, which are now increasingly documented (13, 14), may indirectly affect mortality and growth rates. These indirect effects remain largely unexplored, primarily due to the scarcity of long-term data on both seed production and subsequent impacts on growth or mortality (7). Nonetheless, acknowledging their existence is essential. These indirect effects can shape trends in demographic rates, even without the expected changes in climatic conditions that are known to directly influence these rates.

Weather effects on tree growth are typically understood in terms of photosynthesis and wood formation processes, yet weather also affects resource allocation, such as between growth and reproduction (8, 15, 16). In masting plants, allocation to reproduction can vary dramatically between years; for example, in European beech (*Fagus sylvatica*), allocation to fruit production can vary 34-fold among years, from 508 g m⁻² y⁻¹ to 15 g m⁻² y⁻¹, which negatively correlates with allocation to growth (17). In years of high seed production, more carbon may be invested in reproduction than aboveground growth (18, 19). Besides resource trade-offs, indirect effects such as the replacement of leaf buds with flower buds also affect growth (17, 20). These processes might exacerbate whole-tree shortages in carbon supply during high seed years, especially in the early growing season when carbon demand is high (21).

The strong links between masting and weather render masting potentially sensitive to climate change (22). For instance, masting in Japanese oak (*Quercus crispula*) is correlated with warm springs, resulting in shortened intervals between mast years from a 4-y cycle

Significance

Climate change effects on forest growth and reproduction are widely reported, but indirect effects from their interactions are rarely demonstrated. In a 43-y study of European beech, rising summer temperatures led to more frequent seed production (masting), increasing total reproductive investment. This increased reproductive effort depleted stored resources, causing a 28% reduction in annual growth rates, even without increased drought stress. Diminished growth further reduces future reproductive potential, creating a negative feedback loop. A “perfect storm” of declining growth and reduced viable seed output threatens the sustainability of Europe's most widespread forest tree. We reveal an indirect mechanism by which climate change endangers forests, highlighting the importance of interactions between demographic processes when assessing species sensitivity to climate change.

Author contributions: A.H.-P., J.S., D.K., and M.B. designed research; A.H.-P., J.S., P.A.T., J.G.A.L., D.K., and M.B. performed research; A.H.-P., J.S., and M.B. analyzed data; and A.H.-P., J.S., V.J., M.K.B., P.A.T., J.G.A.L., D.K., and M.B. wrote the paper.

The authors declare no competing interest.

This article is a PNAS Direct Submission.

Copyright © 2025 the Author(s). Published by PNAS. This article is distributed under [Creative Commons Attribution-NonCommercial-NoDerivatives License 4.0 \(CC BY-NC-ND\)](https://creativecommons.org/licenses/by-nc-nd/4.0/).

¹To whom correspondence may be addressed. Email: Andrew.Hacket-Pain@liverpool.ac.uk or michalbogdziewicz@gmail.com.

This article contains supporting information online at <https://www.pnas.org/lookup/suppl/doi:10.1073/pnas.2423181122/-/DCSupplemental>.

Published January 28, 2025.

to a 2-y cycle as temperatures rise (23). In European beech, masting is triggered by summer temperature cues (24, 25). Elevated summer temperatures in recent decades have increased the frequency of weather cues, leading to more frequent initiation of reproduction, and dampening interannual variability of masting; a trend now detected across Europe (26, 27). In principle, masting only affects how a constant reproductive allocation (the fraction of all plant resources devoted to reproduction) is distributed among years to increase reproductive efficiency (28), but it is possible that changing cue frequency might also lead to changes in reproductive allocation. To the extent that changed reproductive investment limits resources allocated to growth, such disrupted temporal allocation to reproduction has the potential to impact growth patterns and long-term growth trends (16). However, the indirect effects of climate change on growth, through effects on reproduction, have not yet been investigated.

Beech is considered a highly drought-sensitive species, and defoliation, dieback, and mortality responses to recent severe summer droughts have raised concerns about the species' long-term future under climate change (16, 29, 30). In addition to the response to individual extreme droughts, growth declines are occurring across the species distribution, driven by increased summer aridity, with stronger declines at drier sites (3, 31, 32). Nonetheless, while the link between European beech masting and interannual variation in growth is well established (17, 33), the potential for longer-term changes in masting to influence growth trends remains untested (16). In this study, we explore the potential effects of warming-driven disruptions in masting patterns on radial growth using populations of European beech (*F. sylvatica*) in which seed production has been monitored annually for 43 y (1980–2022).

To untangle how the coupled growth-reproduction system has responded to climate warming, we cored 57 individuals in which seed production has been monitored as a part of the English Beech Mast Survey since 1980 (34). Tree rings offer annual measures of radial tree growth. They are widely used to study long-term changes in growth patterns (3, 35) and allow us to match individual-level seed production data with growth variation. We first tested how tree-level annual growth rate, as measured by tree ring increments, is correlated with tree-level seed production in the current and previous year and with summer drought. Seed production in the previous year was included as strong masting-driven depletion of resources could produce a carry-over effect and reduce growth the following year (36). Next, we evaluated the temporal trends in growth and attributed the trend to variations in seed production and drought, all in the context of changes in tree size. We predicted that more regular reproduction caused by a warming climate (26, 34) will reduce growth through the existence of a trade-off between masting and growth, even if the United Kingdom is not currently experiencing increasing summer drought due to climate change (37). To better evaluate the impact of reproduction on each tree's remaining resources (including nitrogen, which is important for reproductive resource dynamics: (38), we estimated net plant resources each year following (39). For each tree, this estimates unspent resources as that year's residual from a regression of cumulative seed production against time (a proxy for cumulative resource gain), thereby integrating the impact of all recent reproduction. Next, we tested for the temporal change in the negative association between radial growth and summer temperatures in the preceding year. That correlation arises through indirect effects, as high summer temperatures trigger masting the following year, diminishing growth (16, 33). As warming has reduced the sensitivity of seed production to summer temperatures (26), we expected that to translate into weaker effects of summer temperatures on growth increments.

Results

Despite no change in summer drought (May–July water deficit) over the most recent four decades, European beech annual radial growth has declined by 28%. During the period 1980–2005, the estimated growth rate was a relatively stable average of 1.6 mm per year, but this progressively declined to 1.16 mm per year by 2022 (Fig. 1), after accounting for the effect of increasing tree size over the study period. The onset of this growth decline coincided with the timing of the shift in European beech masting at our site, where annual observations over multiple decades show a changed state to more regular and less synchronized reproduction (“masting breakdown”) as a result of warming temperatures (*SI Appendix*, Fig. S1) (5, 34). Concurrently, and in agreement with the UK Environment Agency analysis of summer precipitation and hydrological records, we detected no trends in summer drought, as measured as the ratio of precipitation to potential evapotranspiration (P-PET) for the period May–July (*SI Appendix*, Fig. S2 and Table S1).

As expected, masting was an important driver of high-frequency growth variation, with narrower growth rings in years of high-seed years (Fig. 2). Summer drought and allocation to reproduction combined to determine annual radial growth, which included a lagged effect of the previous year's reproduction on growth. Alternative models that included vapor pressure deficit or temperature instead of water deficit showed similar results but had lower model performance S2. Growth was reduced most strongly when high seeding co-occurred with drought (Fig. 2). For example, in years when seeding failed and in the absence of summer drought, the estimated growth rate was 2.2 mm y^{-1} , but less than 1 mm y^{-1} when mast years co-occurred with drought. The effect of seed production on growth ($\beta = -0.10$, $\text{SE} = 0.03$) was similar to that of drought ($\beta = 0.13$, $\text{SE} = 0.03$) during the prebreakdown period (1980–2005). While the effect of drought on growth remained




Fig. 1. Decline in European beech growth over the last four decades A) Estimated annual tree ring increments; estimate derived from a generalized linear mixed model that included tree ID ($N = 57$) and site ($N = 7$) as random intercepts, while year (fitted as B-spline) and tree DBH were included as predictors. Shading shows a 95% CI. The dashed vertical line shows the year 2006, at which masting in our populations changed the state into more regular and desynchronized seed production, called masting breakdown (*SI Appendix*, Fig. S1). Note that information on seed production is not included in this model.




Fig. 2. High-frequency variation in growth rate is related to summer drought and allocation to reproduction. (A) Effect sizes of lagged seed production, production in the current year, summer drought, and previous year DBH on annual growth rates. Slopes and associated SE are estimated with a GLMM with Gamma family error distribution that included tree ID ($N = 57$) and site ($N = 7$) as random intercepts. (B) Relationship between the tree ring width and seed production pre-2006 (orange) and post-2006 (blue); estimates and associated 95% CI are derived from the same model as presented at (A). The inset density plot shows the distribution of observations. Surface plots at (C) and (D) show estimated growth rates across combinations of current-year seed production and summer water deficit, with the convex hulls defined by observations (black points). Predictions are derived from the same GLMM as slopes presented at (A). The y-axis range at (D) may suggest that post 2006 period was less dry compared to before 2006, but we detected no trends in summer water deficit (*SI Appendix, Table S1*), and that shift in y-axis range is largely driven by unusually wet 2007 (*SI Appendix, Fig. S2*).

largely unchanged over the last four decades, the effect of the current and previous year reproduction of growth has decreased in the last two decades (postbreakdown, 2006–2022) (Fig. 2).

Trend attribution analysis showed that the European beech growth decline was explained mainly by allocation to reproduction (Fig. 3). Allocation to seed production was the only significant contributor to the trend in annual growth rate, with the sensitivity of growth to increasing allocation to reproduction -0.002 (mm y^{-1} ring increment reduction per additional seed counted). In turn, consistent with a lack of trend in drought, summer water deficit was not a contributor to growth trends (Fig. 3).

Reconstruction of tree-level stored resources from seed production data, following the method of (39), indicates that more frequent reproduction associated with masting breakdown has led to progressive resource depletion (Fig. 4). In the first decades of monitoring (1981–2006), estimated marginal mean stored resources fluctuated stably above zero (18.5–29.7). However, after 2006, when more frequent weather cues caused plants to flower

more often and less in synchrony, resources progressively declined, with negative means recorded in 2012–2016 (-30.3) and 2017–2022 (-40.5) (Fig. 4 and *SI Appendix, Fig. S4*). This analysis suggests that a change in seed production pattern to more regular masting depleted plant resources due to insufficient recovery periods between seed crop production.

The strength of the relationship between growth and the previous year's summer temperature declined by two-thirds over time, with effect size -0.33 ($\text{SE} = 0.02$) before 2006, and -0.12 ($\text{SE} = 0.02$) after that year (Fig. 5A). Annual growth was estimated as 2.7 mm for temperatures below 18°C in the prebreakdown period, which is reduced by 30% to 1.91 mm in the postbreakdown period. At the same time, at a temperature of 22°C , growth was estimated as 0.85 mm, which increased 1.6-fold to 1.36 mm (Fig. 5A). This is consistent with the observation that warming has broken the link between seed production and previous years' summer temperature (Fig. 5B and C), weakening the relationship between growth and seed production (Fig. 2A and B).




Fig. 3. Contribution of the predictor variables to European beech growth trend. The analysis, based on 7 sites and 57 trees (1980–2022), indicated that change in patterns of seed production is the main contributor to the observed decrease in European beech annual growth rate (tree ring increments). The difference between the modeled contributions and the observed trend was considered an unknown contribution to the temporal variation of annual growth. The numbers alongside the bars show the sensitivity of ring width to predictor changes (see *Materials and Methods: Trend attribution*). Error bars for associated contributions indicate SE. See *Materials and Methods* for information about the methodology used to calculate the contributions.

Warm summer temperatures in the previous year are no longer strongly associated with high seed production (Fig. 5C), when growth is lowest. Pre-2006, seed production was largely concentrated in years that followed a sequence of cold and hot summers (Fig. 5B). Postbreakdown (2006–2022), seed production was weakly associated with previous summer temperatures (Fig. 5C).

Discussion

Evidence gathered in this study indicates that climate-driven shifts to more frequent reproduction may explain the observed decline in European beech growth rate. Consequently, we show a previously unidentified and indirect mechanism for climate change-driven growth decline in European beech, that may be operating covertly alongside a direct drought-driven decline already reported in other populations (3, 32, 35). The role of reproduction in mediating the response of growth to warming is consistent with the proposal of ref. 16 that shifts in allocation to reproduction may contribute to reported growth responses to nitrogen deposition in Europe. It is also consistent with an observation that under experimentally induced drought, holm oak (*Quercus ilex*), strawberry tree (*Arbutus unedo*), European beech, and Norway spruce (*Picea abies*) maintain fecundity at the expense of reduced growth (40, 41). In our case, growth decline occurs despite no long-term trend in summer drought stress, because climate warming changes the temporal allocation of resources to reproduction. While this can explain the recent growth decline in the absence of increased summer drought stress in UK European beech forests, we suspect that this driver may also be contributing to the reported growth declines across Europe (3), where consistent warming-driven changes in masting are also occurring (27). A priority for future research is understanding how the reproduction-driven growth decline mechanism reported here operates in conjunction with increases in summer drought stress, and the effect of concurrent warming (i.e., “hotter droughts”) (30).

Growth decreased in our populations because reproductive effort increased. Typically, variations in masting—the allocation of seeds across years—are assumed not to affect the long-term mean reproductive allocation (28). However, temporal changes in reproductive allocation are rarely tested, particularly in masting species, due to high

interannual and interindividual variability. Previous studies on the English Beech Mast Survey data initially reported a recent increase in mean seed production (34), but this was later attributed mainly to a gradual increase in tree sizes over time (5). Estimating an individual plant’s reserves based on its cumulative seed production is a more sensitive method to detect changes in reproductive allocation because it is less affected by short-term variability and allows the effects of large seed crops to persist over several years. Hence, the observed decrease in plant reserves from higher reproductive allocation post-breakdown plausibly explains the growth declines. Importantly, our findings provide a rare demonstration in a mast-seeding tree that changes in the frequency of weather cues can alter both the temporal pattern of reproduction and the mean level of investment. Although increased reproductive allocation under climate change was initially considered a potential consequence of climate warming (42), resource-based models (39, 43) later suggested that changes in reproductive allocation were unlikely. Our results challenge this notion, highlighting a mechanism by which climate change can impact tree growth through altered reproductive strategies.

Shifting allocation to reproduction is an important contributor to the observed decline in growth, despite the weakening of the growth-reproduction trade-off. Post masting breakdown, the positive growth response to low-to-medium seed production (0 to 150 seeds) weakens compared to the prebreakdown period, i.e., years of low seed production are no longer associated with such a large positive growth response. We interpret this as a consequence of increasingly resource-limited conditions, caused by more




Fig. 4. Temporal decline after 2006 in the reconstructed stored plant resources suggests an increase in relative reproductive allocation postmasting-breakdown (blue bins). Resources are estimated from measured individual-level seed production, following the (39) method, see text. Boxplots are based on tree-level averages in each bin (shown as swarmed grey points), while the nonbinned data are presented in *SI Appendix, Fig. S4*. Colors show periods before (orange) and after (blue) 2006, when masting changed the state to less interannually variable and synchronized reproduction (masting breakdown; *SI Appendix, Fig. S1*). The estimated marginal mean contrasts between bins are provided in *SI Appendix, Fig. S5*.




Fig. 5. Weakening sensitivity of annual growth rate to previous year summer temperatures mirrors the similar pattern in association between seed production and summer temperatures. (A) Association between annual tree ring increments and June–July mean max temperatures in the previous year, predicted for the period before (orange) and after (blue) 2006, when masting changed the state to less interannually variable reproduction. Prediction lines are derived from a GLMM with tree ID ($N = 57$) and site ($N = 7$) as random intercepts, shading shows associated 95% CI. Surface plots at (B) and (C) show estimated seed production across combinations of summer (June–July) temperatures one (T1) and 2 y (T2) before seed fall, with the convex hulls defined by observations (black points). Predictors are derived from zero-inflated negative binomial GLMM with tree ID and site used as random intercepts.

frequent reproduction. Strong depletion of carbon and other nutrients in association with masting is widely documented (10, 11, 44), and fruit removal experiments in trees prove that reproduction depletes resources (10, 44). We hypothesize that more frequent reproduction is depleting internal resource reserves, inhibiting the strong positive growth response to low seed production. This effect is not offset by the corresponding dampening of growth response to peak seed production because high seed production years are rare, particularly postbreakdown.

The transition in masting patterns toward more consistent reproduction, and the weaker effects of seed production on growth, paralleled a diminishing correlation between growth and the previous summer temperature. This lagged correlation is ubiquitous in European beech tree-ring chronologies, and results from an indirect effect of summer temperature-cued masting and a growth-reproduction trade-off (33, 45). The correlation is typically stronger in older or larger trees (46), probably because investment in reproduction increases with age (18, 47). In contrast, we found the opposite trend. The decline in correlation over time resulted from the weakening of the growth reproduction trade-off, combined with the reduced sensitivity of seed production to summer temperatures. These effects are likely due to shortened intervals between seeding events, limiting the accumulation of reserves for substantial seeding efforts (26, 39). Our findings align with this narrative; historically, a strong association between summer weather and masting led to predictable and synchronized peak seed production years following warm summers, accompanied by significant negative growth anomalies. However, currently, such warm summers are less reliably linked to high seed production, resulting in less predictable growth responses. That suggests that the breakdown in masting is linked to distinct signals in tree-ring chronologies, i.e., weakened correlations between growth and previous summer temperatures, which might help identify effects described here even in the absence of direct monitoring of both seed production and growth.

The indirect effects of altered reproductive patterns on growth rates can vary depending on the reproductive biology of the species under consideration. In European beech, and other species where masting is triggered by high temperatures (47–49), warming which exceeds the capacity of the species to adapt or acclimate could potentially amplify annual allocation to reproduction and decrease its interannual variation (22, 27), resulting in declining radial growth.

Conversely, in species where reproduction is stimulated by low temperatures or suppressed by high temperatures, warming may reduce the frequency of high-seed years, enabling higher investment in growth. For instance, in *Beilschmiedia tawa*, reduced frequency of low winter and summer temperature cues resulted in reproductive failure in warmer sites (50). On the other hand, masting in other species like North American conifers seems to be largely insensitive to weather trends, which could render their reproductive patterns resistant to climate change (51). Consequently, as the climate warms, reproduction frequency may decrease or increase, depending on the masting mechanisms in operation (22), with potentially predictable responses of growth. The growth response to a change in reproductive allocation will also depend on the strength of trade-offs between seed production and growth (52, 53). Tree ring measurements offer the potential to reconstruct growth patterns in populations in which seed production has been monitored for decades (54), opening the avenues for testing these hypotheses.

In summary, the breakdown of masting within our populations correlates with a decline in growth, even without concurrent trends in summer drought (3, 32). This reveals a previously unrecognized indirect pathway by which climate change can influence growth trends—through its effect on tree reproduction. The documented decline in growth presents a concerning scenario for European beech populations, which now face a cascade of negative effects under warming temperatures. Declining growth indicates reduced vitality, reduced future reproduction due to smaller plant size, lower carbon uptake, increased vulnerability to future climate extremes, and could serve as a precursor to dieback and mortality (55, 56, 57). Moreover, despite increased reproductive effort, the breakdown in masting means the trees actually produce fewer viable seeds (5). Reduced flowering synchrony and decreased interannual variability lead to higher seed predation and decreased pollination efficiency, resulting in up to an 80% decrease in viable seed production (5, 34). While life history theory predicts that long-lived plants should avoid reproduction that lowers survival (52, 58), an abruptly changed environment may render the regulation of reproduction maladaptive, with broad consequences for demographic rates. Thus, European beech faces a “perfect storm” under climate change: lower growth, lower carbon uptake, higher flowering effort, and lower viable seed production, all at a time when robust regeneration and carbon uptake are crucial for forests across Europe.

Materials and Methods

Study System and Data Collection. European beech is a major forest-forming species in temperate Europe, with high economic and ecological importance (59). Beech is a model masting species, with seed production characterized by large interannual variation and synchrony (21, 60). High seed production negatively correlates with aboveground productivity (17, 19, 36). Across the species range, annual growth is mainly driven by early summer moisture availability (31, 61).

The English Beech Mast Survey was established in 1980, with seed production sampled at 15 sites across England annually (5). The ground below each tree was searched for seeds for 7 min, and the collected seeds were later sorted and counted in the laboratory. Our past studies on these European beech populations have shown that interannual variability and among-trees synchrony of beech seed production declined by ~30% over the past four decades, which includes less frequent reproductive failures (34). The state of masting changed abruptly after 2005, the last highly synchronized year. Post-2005, masting switched to more regular and less synchronized reproduction, termed masting breakdown (26). Interannual variation of seed production associated with masting increases reproductive efficiency through reduced seed predation and improved pollination efficiency (62, 63). Consequently, masting breakdown decreased pollination efficiency and increased seed predation, reducing viable seed production by 50 to 80%, depending on tree size (5). Trend attribution analysis suggested that temperature that rose ~ 1°C over the last four decades was responsible (34). Proximally, the masting breakdown is caused by increased frequency of weather cues that trigger flowering (26). Recurrent cueing increased the fraction of years when flowering is triggered, which has led to less frequent failures and more regular reproduction (26, 34), a trend now reported over the majority of European beech range (27).

In 2022, we cored 57 trees across 7 sites. Cores were extracted using an increment corer, at 1.3m above ground level, and perpendicular to any slope to avoid tension wood. Further, 1 to 2 cores were sampled per tree, air-dried, and then prepared using standard dendrochronological methods. Polished cores were imaged using a flatbed scanner, and ring width was measured using CooRecorder v9.8.1 (64).

We extracted daily weather data for each site from the corresponding 0.1° grid cell of the E-OBS dataset (65). The summer water deficit was calculated as $P - PET$, with P standing for precipitation and PET for potential evapotranspiration, summed from May 1st to July 31st (3, 31). Evapotranspiration PET was calculated based on the Thornthwaite equation (66) using the R package SPEI (67). Data used for analysis is openly available (68).

Data Analysis.

Trends in annual growth. We have started our analysis by testing for the temporal trends in annual tree ring increments in sampled trees. To this end, we have fitted a generalized linear mixed model (GLMM) that included tree ID ($N = 57$) and site ($N = 7$) as random intercepts. We used the Gamma family error and log link. Tree ring chronologies typically contain low frequency trends associated with changes in tree size and competition. However, all our sampled trees were reproductively mature and at least canopy codominant when first added to the masting survey, so we did not statistically detrend ring width. Instead, our model included year and tree diameter at breast height (DBH) in the year prior to growth as predictors. The year effect was fitted with a B-spline to allow for nonlinear trends.

High-frequency variation in growth increments. To test for the relationship between annual tree ring increments, summer drought, and seed production, we have fitted a GLMM, with a Gamma family error term and log link. TreeID and siteID were used as random intercepts, while drought (May–July water deficit), seed production in the current year (T), and seed production in the previous year ($T-1$), and tree size (DBH) in the previous year ($T-1$) were used as fixed effects. We also tested alternative models where May–July water deficit was replaced by May–July vapour pressure deficit or May–July temperature.

Trend attribution analysis. To attribute the temporal trends in annual tree ring increments to its possible drivers, we used the temporal contribution method (47, 69). First, using the tree ID and site as the random intercepts, we modeled annual growth as a function of drought (May–July water deficit), seed production (current and past, T and $T-1$), and tree size. We then used the full model to predict the change in ring increments during the study period (1980–2022). We first calculated the observed trend (slope estimate \pm SE of the slope estimate) in our data using GLMMs with random intercepts. We then calculated the trend predicted by the full model and the trends predicted by the same model but maintaining the

predictors constant one at a time (for example, drought is held constant, using the mean values per site, while all other predictors change according to the observations). The difference between the trend predicted by the full model and when one variable was controlled was the contribution of that predictor variable to the change in the response variable. The difference between all individual contributions and the observed trend was considered to be unknown contributions. Finally, we calculated the average ring width sensitivities to predictor changes by taking the differences between the full model trend and the trends from the models with the predictors held constant and dividing it by the trends of the predictor variables. All errors were calculated using the error-propagation method (69).

Reconstructing stored resources. To estimate how change in masting patterns affected relative allocation to reproduction, we reconstructed individual tree stored resources from seed production data, following the approach developed by (39). In brief, this analysis includes fitting a linear model of cumulative reproduction (summed seed count) vs cumulative years (as a surrogate of resource acquisition over time). We used generalized linear mixed models (GLMMs) implemented via the “lme4” package (70) with plant ID and site ID as random intercept and year as random slope. The random effect structure allowed fitting a unique intercept for each plant which estimates stored resources of a plant at the beginning of the monitoring period (39), while random slope allowed heterogeneous resource acquisition of individuals over time (71). As that analysis requires long-term, continuous observations, we used a subset of trees from the English Beech Mast Survey that have been monitored for at least 30 y ($N = 61$).

To test whether estimated stored resources, and therefore relative allocation to reproduction, have changed over time, we have binned the stored resources into 5-y bins, and calculated the tree-level mean for each bin. That was done to smooth over the large year-to-year variation associated with masting (5, 34). We adjusted the bins to have 2006 as one of the bin borders; in consequence, the first and last bins have 6 y. The differences among bins were tested with an LMM model that included tree and site ID as random intercepts.

Growth and seed production sensitivity to weather cues. We tested for the temporal change in the association between radial growth and summer temperatures in the preceding year with a Gamma family, log-link GLMM. The model included tree ID and site ID as random intercepts and the previous year’s mean maximum summer (June–July) temperature as a fixed effect and tree size (DBH) as a covariate. A similar model was constructed for seed production.

Here, we used zero-inflated negative binomial GLMM with log-link, and two fixed factors: seed production in the previous year ($T-1$) and 2 y before seed fall ($T-2$). That is because the relationship between the $T1$ summer temperature is conditional on the $T2$ summer temperature; cold $T2$ summer enhances the response to $T1$ temperature (24, 25). In addition, we included individual-level seed production in $T1$ as a covariate. Both models (with growth and seeding as a response) were fitted to two data subsets; 1980–2005 and 2006–2022. The year 2006 is when European beech masting at our site changed state to more regular and less synchronized reproduction (SI Appendix, Fig. S1) (5, 34).

Data, Materials, and Software Availability. Datasets data have been deposited in OSF <https://osf.io/eujt7/> (68).

ACKNOWLEDGMENTS. This study was supported by the Polish National Science Centre grant 2019/35/D/NZ8/00050. We acknowledge The National Trust (Killerton House and Gardens), Pebblebed Heaths Conservation Trust, Natural England, The Nettlebed Estate, Urban Green Newcastle and Newcastle City Council, and Brighton and Hove City Council for providing permission to collect tree-ring material. We thank Sam Egan, Charlie Knowles and Joep Van Dormolen for help with increment coring and sample preparation, and the Arboriculture Department of Brighton and Hove Council also assisted in retrieving tree-ring samples. We are grateful to J.R. Packham and G.M. Hilton who established the beech mast survey in 1980 and the various field assistants who have helped with the survey over the last five decades.

Author affiliations: ^aDepartment of Geography and Planning, School of Environmental Sciences, University of Liverpool, Liverpool L69 7ZT, United Kingdom; ^bPopulation Ecology Research Unit, Institute of Environmental Biology, Faculty of Biology, Adam Mickiewicz University, 61-614 Poznan, Poland; ^cForest Biology Center, Institute of Environmental Biology, Faculty of Biology, Adam Mickiewicz University, 61-614 Poznan, Poland; ^dSchool of Life Sciences, Keele University, Staffordshire ST5 5BG, United Kingdom; ^eDepartment of Natural Sciences, Manchester Metropolitan University, Manchester M1 5GD, United Kingdom; and ^fSchool of Biological Sciences, University of Canterbury, Christchurch 8140, New Zealand

1. C. Senf *et al.*, Canopy mortality has doubled in Europe's temperate forests over the last three decades. *Nat. Commun.* **9**, 1–8 (2018).
2. H. Hartmann *et al.*, Climate change risks to global forest health: Emergence of unexpected events of elevated tree mortality worldwide. *Annu. Rev. Plant Biol.* **73**, 673–702 (2022).
3. E. M. del Castillo *et al.*, Climate-change-driven growth decline of European beech forests. *Commun. Biol.* **5**, 163 (2022).
4. M. D. Redmond, F. Forcella, N. N. Barger, Declines in pinyon pine cone production associated with regional warming. *Ecosphere* **3**, 1–14 (2012).
5. M. Bogdziewicz *et al.*, Reproductive collapse in European beech results from declining pollination efficiency in large trees. *Glob. Change Biol.* **29**, 4595–4604 (2023).
6. J. S. Clark, D. M. Bell, M. C. Kwit, K. Zhu, Competition–interaction landscapes for the joint response of forests to climate change. *Glob. Change Biol.* **20**, 1979–1991 (2014).
7. J. S. Clark *et al.*, Continent-wide tree fecundity driven by indirect climate effects. *Nat. Commun.* **12**, 1242 (2021).
8. J. D. Lauder, E. V. Moran, S. C. Hart, Fight or flight? potential tradeoffs between drought defense and reproduction in conifers. *Tree Physiol.* **39**, 1071–1085 (2019).
9. C. Corona, A. C. Leeper, J. M. LaMontagne, Differential defoliation and mortality of white spruce and balsam fir by eastern spruce budworm. *For. Ecol. Manage.* **508**, 120042 (2022).
10. A. Sala, K. Hopping, E. J. McIntire, S. Delzon, E. E. Crone, Masting in whitebark pine (*Pinus albiculis*) depletes stored nutrients. *New Phytol.* **196**, 189–199 (2012).
11. Q. Han, D. Kabeya, Recent developments in understanding mast seeding in relation to dynamics of carbon and nitrogen resources in temperate trees. *Ecol. Res.* **32**, 771–778 (2017).
12. T. Nakamura *et al.*, Tree hazards compounded by successive climate extremes after masting in a small endemic tree, *Distylium lepidotum*, on subtropical islands in Japan. *Glob. Change Biol.* **27**, 5094–5108 (2021).
13. A. Hackett-Pain, M. Bogdziewicz, Climate change and plant reproduction: Trends and drivers of mast seeding change. *Philos. Trans. R. Soc. B* **376**, 20200379 (2021).
14. A. Hackett-Pain, F. Roig, D. Ascoli, C. LeQuense, M. Hadad, Reconstruction of *Araucaria araucana* cone production reveals warming intensifies regionally synchronised masting. *Ecosphere* *p.*, in press (2024).
15. D. A. Norton, D. Kelly, Mast seeding over 33 years by *Dacrydium cupressinum* Lamb. (*rimu*) (*Podocarpaceae*) in New Zealand: The importance of economies of scale. *Funct. Ecol.* **2**, 399–408 (1988).
16. C. Leuschner, Drought response of European beech (*Fagus sylvatica* L.)—A review. *Perspect. Plant Ecol. Evol. Syst.* **47**, 125576 (2020).
17. H. Müller-Haubold, D. Hertel, C. Leuschner, Climatic drivers of mast fruiting in European beech and resulting C and N allocation shifts. *Ecosystems* **18**, 1083–1100 (2015).
18. H. Genet, N. Bréda, E. Dufréne, Age-related variation in carbon allocation at tree and stand scales in beech (*Fagus sylvatica* L.) and sessile oak (*Quercus petraea* (Matt.) Liebl.) using a chronosequence approach. *Tree Physiol.* **30**, 177–192 (2009).
19. H. Müller-Haubold, D. Hertel, D. Seidel, F. Knutzen, C. Leuschner, Climate responses of aboveground productivity and allocation in *Fagus sylvatica*: A transect study in mature forests. *Ecosystems* **16**, 1498–1516 (2013).
20. M. J. Vergotti, M. Fernández-Martínez, S. C. Kefauver, I. A. Janssens, J. Peñuelas, Weather and trade-offs between growth and reproduction regulate fruit production in European forests. *Agric. For. Meteorol.* **279**, 107711 (2019).
21. M. Mund *et al.*, It is not just a “trade-off”: Indications for sink- and source-limitation to vegetative and regenerative growth in an old-growth beech forest. *New Phytol.* **226**, 111–125 (2020).
22. M. Bogdziewicz *et al.*, Evolutionary ecology of masting: Mechanisms, models, and climate change. *Trends Ecol. Evol.* **39**, 851–862 (2024).
23. M. Shibata, T. Masaki, T. Yagihashi, T. Shimada, T. Saitoh, Decadal changes in masting behaviour of oak trees with rising temperature. *J. Ecol.* **108**, 1088–1100 (2020).
24. G. Vacchiano *et al.*, Spatial patterns and broad-scale weather cues of beech mast seeding in Europe. *New Phytol.* **215**, 595–608 (2017).
25. V. Journé *et al.*, Summer solstice orchestrates the subcontinental-scale synchrony of mast seeding. *Nat. Plants* **10**, 367–373 (2024).
26. M. Bogdziewicz *et al.*, Climate warming causes mast seeding to break down by reducing sensitivity to weather cues. *Glob. Change Biol.* **27**, 1952–1961 (2021).
27. J. Foest *et al.*, Widespread breakdown in masting in European beech due to rising summer temperatures. *Glob. Change Biol.* **30**, e17307 (2024).
28. D. Kelly, Nutrient scarcity cannot cause mast seeding. *Nat. Plants* **6**, 760–762 (2020).
29. A. Gebler *et al.*, Potential risks for European beech (*Fagus sylvatica* L.) in a changing climate. *Trees-Struct. Funct.* **21**, 1–11 (2007).
30. A. Buras, A. Rammig, C. S. Zang, Quantifying impacts of the 2018 drought on European ecosystems in comparison to 2003. *Biogeosciences* **17**, 1655–1672 (2020).
31. R. Weigel *et al.*, Summer drought exposure, stand structure, and soil properties jointly control the growth of European beech along a steep precipitation gradient in northern Germany. *Glob. Change Biol.* **29**, 763–779 (2023).
32. S. Klesse *et al.*, No future growth enhancement expected at the northern edge for European beech due to continued water limitation. *Glob. Change Biol.* **30** (2024).
33. A. J. Hackett-Pain *et al.*, Climatically controlled reproduction drives interannual growth variability in a temperate tree species. *Ecol. Lett.* **21**, 1833–1844 (2018).
34. M. Bogdziewicz, D. Kelly, P. A. Thomas, J. G. A. Lagueard, A. Hackett-Pain, Climate warming disrupts mast seeding and its fitness benefits in European beech. *Nat. Plants* **6**, 88–94 (2020).
35. A. S. Jump, J. M. Hunt, J. Penuelas, Rapid climate change-related growth decline at the southern range edge of *Fagus sylvatica*. *Glob. Change Biol.* **12**, 2163–2174 (2006).
36. A. J. Hackett-Pain, J. G. Lagueard, P. A. Thomas, Drought and reproductive effort interact to control growth of a temperate broadleaved tree species (*Fagus sylvatica*). *Tree Physiol.* **37**, 744–754 (2017).
37. Environment Agency, *Annex to The Review of The Research and Scientific Understanding of Drought* (Environment Agency, Bristol, 2023).
38. T. Abe *et al.*, Parameterisation and validation of a resource budget model for masting using spatiotemporal flowering data of individual trees. *Ecol. Lett.* **19**, 1129–1139 (2016).
39. M. Rees, D. Kelly, O. N. Bjørnstad, Snow tussocks, chaos, and the evolution of mast seeding. *Am. Nat.* **160**, 44–59 (2002).
40. M. Bogdziewicz, M. Fernández-Martínez, J. M. Espelta, R. Ogaya, J. Penuelas, Is forest fecundity resistant to drought? Results from an 18-yr rainfall-reduction experiment. *New Phytol.* **227**, 1073–1080 (2020).
41. B. D. Hesse *et al.*, Mature beech and spruce trees under drought – Higher C investment in reproduction at the expense of whole-tree NSC stores. *Environ. Exp. Bot.* **191**, 104615 (2021).
42. M. McKone, D. Kelly, W. Lee, Effect of climate change on mast-seeding species: Frequency of mass flowering and escape from specialist insect seed predators. *Glob. Change Biol.* **4**, 591–596 (1998).
43. A. Satake, Y. Iwasa, Pollen coupling of forest trees: Forming synchronized and periodic reproduction out of chaos. *J. Theor. Biol.* **203**, 63–84 (2000).
44. I. L. Roncé *et al.*, Reproduction alternation in trees: Testing the resource depletion hypothesis using experimental fruit removal in *Quercus ilex*. *Tree Physiol.* **43**, 952–964 (2023).
45. G. Piovesan, B. Schirone, Winter North Atlantic Oscillation effects on the tree rings of the Italian beech (*Fagus sylvatica* L.). *Int. J. Biometeorol.* **44**, 121–127 (2000).
46. A. J. Hackett-Pain, L. Cavin, A. D. Friend, A. S. Jump, Consistent limitation of growth by high temperature and low precipitation from range core to southern edge of European beech indicates widespread vulnerability to changing climate. *Eur. J. For. Res.* **135**, 897–909 (2016).
47. M. B. Pesendorfer *et al.*, Investigating the relationship between climate, stand age, and temporal trends in masting behavior of European forest trees. *Glob. Change Biol.* **26**, 1654–1667 (2020).
48. M. Fernández-Martínez, S. Vicca, I. A. Janssens, J. M. Espelta, J. Peñuelas, The North Atlantic Oscillation synchronises fruit production in western European forests. *Ecography* **40**, 864–874 (2017).
49. J. G. Zamorano, T. Hokkanen, A. Lehtikainen, Climate-driven synchrony in seed production of masting deciduous and conifer tree species. *J. Plant Ecol.* **11**, 180–188 (2018).
50. O. M. M. Yukich-Clendon *et al.*, Global change explains reduced seeding in a widespread New Zealand tree: Indigenous Tuhoe knowledge informs mechanistic analysis. *Front. For. Glob. Change* **6**, 1172326 (2023).
51. J. M. LaMontagne, M. D. Redmond, A. P. Wion, D. F. Greene, An assessment of temporal variability in mast seeding of North American Pinaceae. *Philos. Trans. R. Soc. B* **376**, 20200373 (2021).
52. S. C. Thomas, *Age-Related Changes in Tree Growth and Functional Biology: The Role of Reproduction*. pp. 33–64 (2011).
53. T. W. Patterson, C. H. Greenberg, A. Hackett-Pain, Acorn production, climate, and tree-ring growth of five oak species in southern Appalachian forests. *For. Ecol. Manage.* **546**, 121310 (2023).
54. A. Hackett-Pain *et al.*, Mastree+: Time-series of plant reproductive effort from six continents. *Glob. Change Biol.* **28**, 3066–3082 (2022).
55. M. Dobberty, Tree growth as indicator of tree vitality and of tree reaction to environmental stress: A review. *Eur. J. For. Res.* **124**, 319–333 (2005).
56. S. Gillner, N. Rüger, A. Roloff, U. Berger, Low relative growth rates predict future mortality of common beech (*Fagus sylvatica* L.). *For. Ecol. Manage.* **302**, 372–378 (2013).
57. A. Neycken, M. Scheggia, C. Bigler, M. Lévesque, Long-term growth decline precedes sudden crown dieback of European beech. *Agric. For. Meteorol.* **324**, 109103 (2022).
58. J. R. Obeso, The costs of reproduction in plants. *New Phytol.* **155**, 321–348 (2002).
59. C. Leuschner, H. Ellenberg, Beech and mixed beech forests. *Ecol. Cent. Eur. For.: Veg. Ecol. Cent. Eur.* **1**, 351–441 (2017).
60. S. G. Nilsson, U. Wastljung, Seed predation and cross-pollination in mast-seeding beech (*Fagus sylvatica*) patches. *Ecology* **68**, 260–265 (1987).
61. I. Dorado-Liñán *et al.*, Jet stream position explains regional anomalies in European beech forest productivity and tree growth. *Nat. Commun.* **13**, 2015 (2022).
62. D. Kelly, The evolutionary ecology of mast seeding. *Trends Ecol. Evol.* **9**, 465–470 (1994).
63. I. S. Pearce, W. D. Koenig, D. Kelly, Mechanisms of mast seeding: Resources, weather, cues, and selection. *New Phytol.* **212**, 546–562 (2016).
64. R. S. Maxwell, L. A. Larsson, Measuring tree-ring widths using the Coorecorder software application. *Dendrochronologia* **67**, 125841 (2021).
65. R. C. Cornes, G. van der Schrier, E. J. van den Besselaar, P. D. Jones, An ensemble version of the E-OBS temperature and precipitation data sets. *J. Geophys. Res.-Atmos.* **123**, 9391–9405 (2018).
66. C. Thornthwaite, An approach toward a rational classification of climate. *Geogr. Rev.* **38**, 55–94 (1948).
67. S. Beguería, S. Vicente-Serrano, Spei: Calculation of the standardised precipitation evapotranspiration index (version 1.7, R Package, 2017). <https://cran.r-project.org/web/packages/SPEI/index.html>. Accessed 11 November 2024.
68. A. Hackett-Pain, J. Szymkowiak, M. Bogdziewicz, Data from “Growth decline in European beech associated with temperature-driven increase in reproductive allocation.” OFS. <https://osf.io/eujt7/>. Deposited 28 December 2024.
69. M. Fernández-Martínez, J. Maspons, Tempcont: Temporal contributions on trends using mixed models (version 0.1.0, R Package, 2019). <https://cran.r-project.org/web/packages/TempCont/index.html>. Accessed 11 November 2024.
70. D. Bates, M. Mächler, B. M. Bolker, S. C. Walker, Fitting linear mixed-effects models using lme4. *J. Stat. Softw.* **67**, 1–48 (2015).
71. E. E. Crone, L. Polansky, P. Lesica, Empirical models of pollen limitation, resource allocation, and mast seeding by a bee-pollinated wildflower. *Am. Nat.* **166**, 396–408 (2005).