

Please cite the Published Version

Martinho, Diogo V., Rebelo, André, Clemente, Filipe Manuel, Costa, Renato, Gouveia, Élvio R., Field, Adam , Casonatto, Juliano, van den Hoek, Daniel, Durkalec-Michalski, Krzysztof, Ormsbee, Michael J. and Sarmento, Hugo (2025) Nutrition in CrossFit® – scientific evidence and practical perspectives: a systematic scoping review. Journal of the International Society of Sports Nutrition, 22 (1). 2509674 ISSN 1550-2783

DOI: https://doi.org/10.1080/15502783.2025.2509674

Publisher: Taylor & Francis

Version: Published Version

Downloaded from: https://e-space.mmu.ac.uk/639886/

Usage rights: (cc) BY

Creative Commons: Attribution 4.0

Additional Information: This is an open access article published in Journal of the International Society of Sports Nutrition, by Taylor & Francis.

Data Access Statement: The data that support the findings of this study are available from the corresponding author, DVM, upon reasonable request.

Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines)





ISSN: 1550-2783 (Online) Journal homepage: www.tandfonline.com/journals/rssn20

Nutrition in CrossFit® – scientific evidence and practical perspectives: a systematic scoping review

Diogo V. Martinho, André Rebelo, Filipe Manuel Clemente, Renato Costa, Élvio R. Gouveia, Adam Field, Juliano Casonatto, Daniel van den Hoek, Krzysztof Durkalec-Michalski, Michael J. Ormsbee & Hugo Sarmento

To cite this article: Diogo V. Martinho, André Rebelo, Filipe Manuel Clemente, Renato Costa, Élvio R. Gouveia, Adam Field, Juliano Casonatto, Daniel van den Hoek, Krzysztof Durkalec-Michalski, Michael J. Ormsbee & Hugo Sarmento (2025) Nutrition in CrossFit® – scientific evidence and practical perspectives: a systematic scoping review, Journal of the International Society of Sports Nutrition, 22:1, 2509674, DOI: <u>10.1080/15502783.2025.2509674</u>

To link to this article: https://doi.org/10.1080/15502783.2025.2509674

9	© 2025 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.	+	View supplementary material 🕝
	Published online: 05 Jun 2025.		Submit your article to this journal 🕑
hl	Article views: 1319	Q	View related articles 🕝
CrossMark	View Crossmark data 🗹		



👌 OPEN ACCESS 🖲 🗅

Check for updates

Nutrition in CrossFit[®] – scientific evidence and practical perspectives: a systematic scoping review

Diogo V. Martinho (D^{a,b}, André Rebelo (D^{c,d}, Filipe Manuel Clemente (D^{e,f,g}, Renato Costa^a, Élvio R. Gouveia (D^{b,h}, Adam Field (Dⁱ, Juliano Casonatto (Dⁱ, Daniel van den Hoek (D^k, Krzysztof Durkalec-Michalski (D^{l,m}, Michael J. Ormsbee (D^{n,o,p,q}) and Hugo Sarmento (D^{a,r})

^aUniversity of Coimbra, Faculty of Sport Sciences and Physical Education, Coimbra, Portugal: ^bLaboratory of Robotics and Engineering Systems, Interactive Technologies Institute, Funchal, Portugal; ^cUniversidade Lusófona, CIDEFES, Centro de Investigação em Desporto, Educação Física e Exercício e Saúde, Lisboa, Portugal; ^dCOD, Center of Sports Optimization, Sporting Clube de Portugal, Lisboa, Portugal; ^eInstituto Politécnico de Viana do Castelo, Rua Escola Industrial e Comercial de Nun'Álvares, Escola Superior Desporto e Lazer, Viana do Castelo, Portugal; Sport Physical Activity and Health Research & Innovation Center, Viana do Castelo, Portugal; ^gGdansk University of Physical Education and Sport, Gdańsk, Poland; ^hUniversity of Madeira, Department of Physical Education and Sport, Funchal, Portugal; Manchester Metropolitan University, Department of Sport and Exercise Science, Institute of Sport, Manchester, UK; ^JUniversity of North Paraná, Research Group in Physiology and Physical Activity, Londrina, Brazil; ^kUniversity of the Sunshine Coast, School of Health, Petrie, Queensland, Australia; Poznan University of Physical Education, Department of Sports Dietetics, Poznan, Poland; "Charles University, Sport Sciences-Biomedical Department, Praque, Czechia; "Florida State University, Department of Health, Nutrition, and Food Sciences, Tallahassee, FL, USA; °Florida State University, Institute of Sports Sciences and Medicine, Tallahassee, FL, USA; PUniversity of KwaZulu-Natal, Discipline of Biokinetics, Exercise, and Leisure Sciences, Durban, South Africa; ^qUniversity of Arkansas for Medical Sciences, Center for Aging and Longevity, Geriatrics, Little Rock, AR, USA; ' University of Coimbra, CIPER, FCDEFUC, Coimbra, Portugal

ABSTRACT

Background: CrossFit[®] sessions and competitions are characterized by high-intensity challenges that combine aerobic and anaerobic activities with short recovery periods. As a result, effective nutritional practices play a crucial role in optimizing performance and enhancing recovery. Therefore, nutritional practices are central to optimizing performance and accelerating recovery. This review aims to summarize current evidence on nutritional and ergogenic aid responses to CrossFit[®] practice.

Methods: The search was conducted in four electronic databases (PubMed, Web of Science, Scopus and SportDiscus). Gray literature was also extracted for studies exploring the nutritional habits of CrossFit[®] participants as well as the ingestion of ergogenic aids. In addition, a meta-analysis was conducted to examine the impacts of dietary habits and ergogenic aids on performance.

Results: Forty-nine studies met the eligibility criteria and were included in the current review. Carbohydrate intake was below the recommendations for athletes, although protein ingestion

ARTICLE HISTORY

Received 18 August 2024 Accepted 15 May 2025

KEYWORDS

Carbohydrates; caffeine; energy; performance; strength; conditioning

© 2025 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group.

This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/ licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. The terms on which this article has been published allow the posting of the Accepted Manuscript in a repository by the author(s) or with their consent.

CONTACT Diogo V. Martinho 🖾 dvmartinho92@hotmail.com 🗈 University of Coimbra, Faculty of Sport Sciences and Physical Education, Avenida Conímbriga Pavilhão 3, Coimbra 3040-248, Portugal

Supplemental data for this article can be accessed online at https://doi.org/10.1080/15502783.2025.2509674

remains adequate. High-carbohydrate diets had a positive effect on CrossFit[®] performance. The evidence concerning the effects of a ketogenic diet on performance is limited. When used prior to or during the workout, the impact of carbohydrates on CrossFit[®] performance was negligible, whereas the effect of caffeine was significant. Ergogenic aids, particularly creatine and protein, are commonly used by CrossFit[®] participants.

Conclusion: The standard diets recommended to CrossFit[®] participants need to be revised because they are characterized by lower values of carbohydrates. Caffeine should be used prior to or during the CrossFit[®] sessions. Regarding the impact of ergogenic aids on recovery, future studies are needed.

1. Background

CrossFit[®] has gained worldwide popularity over the last decade. According to data from the World Metrics Report [1], the number of CrossFit[®] affiliations increased by 118% from 2005 to 2015. The CrossFit[®] Open, an event that allows competitors to participate for three weeks, included 344,396 participants in 2024. In CrossFit[®], athletes are required to train distinct components of fitness (cardiorespiratory, stamina, strength, flexibility, power, speed, coordination, agility, balance, and accuracy) and a variety of movements (e.g. deadlift, squat, power clean, push-ups, pull-ups, burpees, rowing, running, cycling) at high intensities and with short periods of recovery [2,3]. Optimizing performance and recovery are two central aspects of success in CrossFit[®] participation [4–6]. Many studies have described the physical and physiological aspects of CrossFit[®] workouts [4,7,8] and have focused on examining the time needed to recover from a specific workout [9-11]. While characterizing CrossFit in terms of physical and physiological demands based on a single workout requires caution, data from Spanish CrossFit® participants indicate that restoring normal levels of physical performance and creatine kinase typically takes 48 to 72 hours [12]. Among Brazilian practitioners, 48 hours of recovery allowed them to reestablish physical performance, whereas creatine kinase values were higher 72 hours after the CrossFit[®] training than before the workout [9]. Therefore, adequate nutritional practices and appropriate nutritional ergogenic aids can optimize performance and accelerate recovery [13,14].

The dietary recommendations for CrossFit[®] participants relate to following the Paleoor Zone Diets [15]. The Paleo Diet emphasizes the consumption of unprocessed foods, while the Zone Diet focuses on maintaining stable glucose levels through a macronutrient distribution of 40% carbohydrates (CHO), 30% protein, and 30% fat. Both dietary practices tend to neglect the importance of CHO [16,17], which is surprising given the wealth of evidence pertaining to the impact of ingesting CHO before, during, and after training to improve performance and reduce fatigue [18–20]. Two recent reviews published in 2021, including 14 and 13 studies, summarized the effects of dietary and supplementary interventions among CrossFit[®] participants with a focus on performance [21,22]. However, assessments of energy and macronutrient intake among CrossFit[®] participants are limited. This gap needs particular attention to determine whether CrossFit[®] participants meet the nutritional recommendations established for athletes. Additionally, neither of the reviews addressed the prevalence of supplements used in CrossFit[®]. Given the rising popularity of CrossFit[®], an increase in the number of studies in the field of nutrition over the past few years is expected. Therefore, the available information regarding nutritional issues in CrossFit[®] needs to be systematically reviewed and summarized.

To improve the dietary practices of nutritionists working with CrossFit[®] athletes, this systematic scoping review aims to do the following: (1) summarize the dietary practices of participants, (2) describe nutritional and energetic intake, (3) examine the acute and chronic effects of nutritional ergogenic aids on performance and recovery, and (4) identify the gaps in the available literature and provide suggestions for future research.

2. Methods

The present scoping review conformed to the standards set by the latest revision of the Cochrane guidelines [23] and followed the written instructions proposed by the PRISMA 2020 guidelines [24] as well as the respective extension for scoping reviews [25]. The protocol was also preregistered on the INPLASY plataform (doi:https://doi.org/10.37766/ inplasy2024.6.0059).

2.1. Eligibility criteria

Original manuscripts published in peer-reviewed journals, master dissertations, doctoral theses, research reports, doctoral dissertations, conference presentations, abstracts, and clinical trials written in English, Portuguese, and Spanish were included in the present review. The Participants, Intervention, Comparator, Outcomes, and Study Design (PICOS) framework was used to define the eligible studies for the present review: Participants – adult CrossFit[®] participants; Intervention – studies that described dietary practices or examined the effects of nutrition on performance or recovery; Comparator – studies assessing the impact of nutrition on performance or recovery versus a placebo or control; Outcomes – energy and/or macronutrient intake, energy expenditure, energy balance, performance, and/or recovery; Study design – observational and interventional studies. No restrictions were applied in terms of publication date.

2.2. Search information and information sources

The search strategy included the combination of the following terms: ((*nutrition** OR "*nutritional strategy*" OR "*nutritional intervention*" OR *diet**, *carbohydrate* OR *glucose* OR *protein* OR *collagen* OR *fat* OR *ketone** OR *antioxidant** OR *vitamin* OR *polyphenol** OR *fruit* OR *creatine* OR *caffeine* OR *nitrate** OR *beetroot* OR "*tart cherry*" OR *beta alanine* OR *sodium bicarbonate* OR *supplement**OR *energy** OR *macronutrient** OR *micronutrient** OR *mineral** OR *electrolyte**) AND *CrossFit*®). The search was divided into two different phases: 1) identification of studies via databases and 2) identification of studies via other methods (dissertation and thesis databases; gray literature databases; trial results in platforms; other systematic reviews; and reference lists of included studies). Four electronic databases were consulted: PubMed, Web of Science, Scopus and SportDiscus. The full search strategy for each database can be found in Supplementary Material 1. The Open Access Thesis and Dissertations

4 🕒 D. V. MARTINHO ET AL.

database (https://otad.org) was used to check search for master dissertations and doctoral theses. Following the Cochrane guidelines, two platforms were used to identify trial registers (https://clinicaltrials.gov/; https://trialsearch.who.int/) [26]. The references of other systematic reviews on the same topic were consulted in the Web of Science database to identify other potential references. The search strategy adopted was similar to that mentioned above, including the Boolean connector "AND" with the term "systematic review." The titles of the reference lists of those studies included in the present review were consulted to identify additional studies. The search strategy was conducted on the same day for all databases and platforms (11 June 2024).

Dedicated computer software was used for reference management, facilitating deduplication and screening steps (EndNoteTM 21.0, ClarivateTM). Following the automatic omission of duplicates, two authors manually screened the remaining references for their relevance (DVM/HS). The titles and abstracts were screened first. The full texts of the studies were then screened to ensure that the studies met the eligibility criteria. Two researchers (DVM/HS) conducted screenings, and when necessary, a third author (AR) was contacted to resolve any disagreements.

2.3. Data extraction and items

The first author (DVM) created a template to organize the relevant data. The Microsoft Excel[®] document included four sheets: (1) prevalence of nutrition use, (2) data concerning dietary and nutritional intake, (3) the effects of dietary interventions on performance, and (4) the effects of nutrition on performance and recovery. To examine the prevalence of nutritional aids, the following information was retrieved: sample size, percentage of nutritional ergogenic aid use, and percentage consumed by athletes. Data related to energetic and nutritional intake included the relative and absolute values of daily energy expenditure, intake, and macronutrients. Information surrounding the effects of specific diets on performance considered the study design, sample size, intervention, performance outputs, specificity of the performance variable (i.e. CrossFit[®] movements or functional capacities), and mean and standard deviation of both conditions. Corresponding information was collected for studies that tested the effects of nutritional aids on performance and recovery. The corresponding authors were individually contacted when the data were not reported. When the data were presented graphically, specific software was used to extract the data (http://www.getdata-graph-digitizer.com). This software has been shown to be accurate and precise in extracting mean and standard deviation values from graphs [27].

2.4. Statistical analysis

2.4.1. Energy and nutritional intake

Sex, sample size, and the means and standard deviations of energy and macronutrient intake were retrieved from each study. The overall means of energy, CHO, and protein contents were split by sex. In parallel, data were organized according to the year of study publication to estimate the tendency of energetic, CHO, and protein intake across time using the moving averages.

2.4.2. Percentage of use of nutritional ergogenic aids

Supplementation prevalence was examined based on sample size and percentage of supplements used by CrossFit[®] participants. The number of cases was calculated based on the preceding information, and a random effect model was considered. The mean prevalence, 95% confidence intervals, and I² were retrieved for the analysis. The I² reflects the proportion of true variance to observed variance, contrasting the true and observed effects.

2.4.3. Interventions (diets and ergogenic aids)

Standardized mean differences corrected by the degrees of freedom and expressed as effect size (i.e. *Hedges's g*) were used to compare CrossFit[®] performance after the implementation of diets with higher values of CHO or specific acute nutritional strategies (CHOs and caffeine). The effect size calculation was based on a random effects model. The effect sizes were interpreted as follows: < 0.2, trivial; 0.2–0.6, small; > 0.6–1.2, moderate; > 1.2–2.0; large; > 2.0–4.0, very large; and > 4.0, extremely large [28]. An integrative approach was chosen to combine multiple effect sizes of the same study because they represent different features of CrossFit[®] performance [29]. Heterogeneity was assessed via the l² and qualitatively described as follows: low (l² <25%), moderate (25–75%), and > high (l² >75%) [30]. The l² and 95% confidence intervals were used to investigate heterogeneity. The bias of publication was graphically inspected with a funnel plot and statistically verified with Egger's test (observed Hedge's g values were contrasted with the respective standard errors). The trim-and-fill method of Duval and Tweedie was used to adjust for potential publication biases [31].

The meta-analysis was performed via Comprehensive Meta-Analysis software version 2.2.064 (BiostatTM, Englewood, NJ, USA).

2.4.4. Risk of bias

The risk of bias was evaluated via two different tools according to the study design: (1) the Quality Assessment Tool for Observational Cohort and Cross-Sectional Studies [32] and (2) the PEDro scale, an 11-item validated tool used to measure the risk of bias and statistical reporting of clinical trials (https://pedro.org.au/english/resources/pedro-scale/). The first tool included fourteen items about the research question, study population, groups recruited from the same population and uniform eligibility criteria, sample size justification, exposure assessed prior to measurement outcome, sufficient timeframe to observe an effect, different levels of the exposure effect, exposure measurement, repeated exposure assessment, outcome measurement, blinding of outcome assessors, follow-up rate and statistical analysis. Each item was analyzed individually and assigned a designation of yes, "no," or 'not reported or applicable. The 11-item PEDro scale presents questions about the eligibility criteria, group allocation, group similarities at baseline, blinding procedures, completion rates of the outcome measures, statistical analyses, and reporting of outcome measures. Each item was assigned a value of "yes" (corresponding to 1 point) or "no" (corresponding to 0 points). The first item is not used to calculate the PEDro score. The methodological quality of the interventional studies was interpreted using the following criteria [33]: 0-3 points was considered "poor" quality, 4-5 points was considered "fair" quality, 6-8 points was considered "good" quality, and 9-10 points was considered "excellent" quality. Two authors (DVM and AR) independently assessed the risk of bias. In the event of disagreements, a third author (HS) was consulted, and a final decision was reached by consensus. No studies were excluded based on their assessed risk of bias.

3. Results

3.1. Study identification and selection

The search was conducted in four databases, and 784 records were identified. Of these, 281 were removed because they were identified as duplicates. Thus, the titles and abstracts of 502 articles were screened. After this process, 67 manuscripts remained potentially eligible for the present review. Six reasons were identified to exclude 18 reports: observational studies of supplement intake (n = 3) that did not include control or placebo groups (n = 3), one case study (n = 1), outcomes were not recovery or performance (n = 3), the sample was not mentioned as CrossFit[®] participants or athletes (n = 6), and duplicate data (n = 2). Duplicate data refer to records extracted from the same original papers and abstracts published at conferences. Forty-nine papers extracted from the four databases met the inclusion criteria and were included in the present review. The reference lists of the 49 papers were individually reviewed, and two additional references were considered for the current review. Six master's or doctoral theses were identified via the Open Access Thesis and Dissertations database. One master's thesis was not publicly available (https://digitalcommons.lib.uconn.edu/gs theses/134/). The first author (DVM) contacted the author of the thesis, and it was confirmed that access to the PDF document was unavailable. Consequently, this record was not retrieved for the present review. Another record was found in the International Clinical Trials Registry Platform of the World Health Organization. Six records were identified via other sources and were combined with 51 original papers. The present review included 57 records, as shown in Figure 1.

3.2. Study characteristics

Table 1 summarizes the primary information extracted from each study about the use of ergogenic aids and dietary and nutritional patterns of CrossFit[®] participants (e.g. sample size, origin of sample, objective, methods used, and main results) [34–56]. Studies were conducted mainly on Brazilian participants (n = 16, ~70%), and four included American participants (~17%). The total number of CrossFit[®] participants in this group of studies was 4,560. Two main topics emerged from Table 1: (1) food, energy, and nutritional intake [34,35,37,40–45,47–52,54,55] and (2) the use of supplements [36,38–40,43,44,46,50,52,53,56] The percentage and type of supplements used were evaluated via questionnaires, whereas the daily energy and nutritional intake were estimated via 24-hour recall methods or food diaries.

Table 2 presents the studies that focus on investigating the effects of a specific type of diet on physical performance [57–63]. The sample sizes of these studies ranged from 11 to 27 male and female CrossFit[®] participants. Three studies tested the effects of diets with a considerable percentage of carbohydrates [58,59,62], three papers focused on the impact of fat (i.e. a ketogenic diet) [60,61,63], and one study investigated the influence



*The reviews of Quaresma et al. (2021) and Silva de Souza et al. (2023) included 14 and 13 papers about nutrition and supplements, respectively.

Figure 1. Flow chart of records identification, screening and inclusion in the present review articleCHO (carbohydrates).

of fasting on performance [57]. The time of interventions and performance outputs varied across studies.

The characteristics of the studies (study design, dosage, timing, and main findings) of ergogenic supplements are shown in Table 3. Sodium bicarbonate [66,67,72,73,79,85] carbohydrates [64,70,86,87], caffeine [65,74,80,84], betaine [69,81], capsiate [71], beta-alanine [75], tribulus terrestris [76,78], nitrate [82,88], and beetroot juice [83] were tested in CrossFit[®] participants. The combination of different supplements was also investigated in four studies [68,77,89,90]. Most of the supplements were ingested before the CrossFit[®] workout [65–68,70,71,73,74,76,79– 85,87,89,90], two studies investigated the effects of supplements during the workout [64,87], and one study tested the impact of CHO intake prior to and during the workout [86].

3.3. Results of individual studies and meta-analysis

3.3.1. Energy and nutritional intake

Nine studies [34,37,41,42,45,48,49,52,55] aimed to analyze nutritional and dietary intake. Data from 10 CrossFit[®] participants with six months of experience were reported as medians and, consequently, are not included in the figures [55]. The average and standard deviation of energy intake data points extracted from the literature combining male and female participants revealed a mean ingestion of 2247 ± 606 kcal.day⁻¹ (Figure 2). Although only two studies exclusively included female participants [41,48], the intake of females (1746 ± 40 kcal.day⁻¹) is substantially lower than that of male CrossFit[®] participants (2360 ± 174 kcal.day⁻¹). CHO intake was greater in females than in males (females: $3.4 \text{ g.kg}^{-1} \text{ day}^{-1}$; males: $3.2 \text{ g.kg}^{-1} \text{ day}^{-1}$), but protein ingestion was comparable in both sexes (Figure 3). The average intakes of CHO and protein in both groups were $3.6 \text{ g.kg}^{-1} \text{ day}^{-1}$ and $1.7 \text{ g.kg}^{-1} \text{ day}^{-1}$, respectively. The moving averages of energy and CHO

8 😔 D. V. MARTINHO ET AL.

			Methodology – nutritional/ergogenic	
Study	Sample characteristics	Country	aids issues	Main results
De Jesus et al. [34]	4 high performance male and female athletes (19–30 yrs); training experience: NB	Brazil	24 hours recall	 Nutritional intervention ↑ energy and macronutrients intake
Brustolin et al. [35]	20 CrossFit® male and female participants (28.7 ± 4.5 yrs); training experience: NR.	Brazil	Interview	 Hunce. 40% performed 3–4 meals per day. Fruits, vegetables, cereals, legumes and milk and dairy products are † selected. 25% used supplements. 75% ingested a pre-workout 30 minutes prior to the CrossFit® section: banana, oats, cinnamon and honey. 70% reported that ingested 1 hour post CrossFit® workout: different foods.
Dos Santos et al. [36]	112 CrossFit® male and female participants of different competitive levels (28.9 ± 7.6 yrs); training experience: > 1 yr.	Brazıl	Questionnaire	 78% of participants used supplements. Supplements were mainly used to 1 performance. >50% of participants used supplements reported to use supplements at least 5 or more times per week. Protein and creatine were used by 63% and 48% of participants, respectively.
Rezende et al. [37]	20 male and female CrossFit® participants; training experience: NR.	Brazil	24 hours recall	 → energy intake and relative protein between groups. → % CHO and ↑ fat in CrossFit[®] participants.
Faria et al. [38]	113 CrossFit® male and female participants; training experience: < 3 mths to >2 yrs.	Brazil	Questionnaire	 75% of participants used supplements. 81% of participants used whey protein († recovery and muscle mass). 53% of participants used creatine († performance and muscle mass). 53% of participants used BCAAs († recovery).
Comerlatto et al. [39]	217 male and female CrossFit® participants (33.7 ± 5.2 yrs); training experience: NR.	Brazil	Questionnaire	 61% of participants used supplements. 53% ingested whey protein. 31% ingested creatine.
Filho et al. [40]	160 male and female CrossFit® participants (25–30 yrs); training experience: NR.	Brazil	Questionnaire: supplementation; intake; food pattern: food frequency questionnaire	 98% of participants used supplements. Creatine was ingested by 61% of participants. Why protein was ingested by 28% of participants. ↓ number of participants used supplements prior or post training. Fruits, legumes, milk and dairy products are ↑ selected.

Table 1. Studies about ergogenic aids statistics, energetic, and nutritional characteristics of CrossFit[®] participants.

(Continued)

			Methodology –	
Study	Sample characteristics	Country	aids issues	Main results
Pearson and Jenkins [41]	443 CrossFit [®] male and female participants (36.6 \pm 11.4 yrs); training experience: > 6 mths	US	Dietary health questionnaire	 Dietary intake was associated weight status, sex, age, exercise and nutritional goals.
Gonçalves et al. [42]	25 CrossFit [®] male and female athletes (32.0 ± 8.9 yrs); training experience: NR.	Brazil	Food diary	 The mean energy intake was 2844 ± 724 kcal.day⁻¹. The relative intake of CHO and protein were 4.6 g.kg⁻¹·day⁻¹ and 2.2 g.kg⁻¹·day⁻¹, respectively. 64% of athletes consumed ↓ CHO than the recommendations. ↓ values of micronutrients (calcium, potassium, selenium, vitamins A, B9, B12, and D) ingestion were also noted.
Mesquita and Cavalcanti [43]	24 CrossFit [®] male and female participants (33.6 yrs); training experience: NR.	Brazil	Food frequency questionnaire	 † salad, vegetables, roots, fruits, legumes, rice or pasta, milk and dietary products, meat, fish and eggs. 75% of the food ingested was derived from protein sources. 4 of participants used supplements (21%).
Brisebois et al. [44]	2576 CrossFit® male and female participants (33.6 yrs); training experience: NR.	US, Canada, UK, Australia	Questionnaire	 - 60% of participants practiced a specific diet. - 82% use supplements twice per week. - Macro Counting, intermittent fasting, paleo diets were 1 reported. - 2015 supplements were described. - Protein was the most used supplement (51%), followed by creatine (23%) and pre- workouts (21%). - Nutrition was recognized as determinant for CrossFit[®] performance. - CHO and protein were widely recognized as determinant for performance.
Vieira et al. [45]	12 male recreational CrossFit® participants (30.2 ± 5.5 yrs); training experience: NR.	Brazil	Food diary	 performance. The mean energy intake was 2561 ± 593 kcal.day⁻¹. The relative intakes of CHO and protein were 3.5 ± 1.2 g.kg⁻¹.day⁻¹ and 1.5 ± 0.5 g.kg⁻¹.day⁻¹, respectively. Fat represents 34% of total daily energy intake
Higino and Freitas [46]	450 male and female CrossFit® participants; training experience: < 3 mths to >2 yrs.	Brazil	Questionnaire	 - 71% of participants used supplements. - 31% of participants ingested protein supplements.
				(Continued)

Table 1. (Continued).

10 🕒 D. V. MARTINHO ET AL.

		_	Methodology – nutritional/ergogenic	
Study	Sample characteristics	Country	aids issues	Main results
Kutch [47]	73 female CrossFit [®] participants; training experience: 3.3 yrs.	US	Food diary	 Most of CrossFit participants did not meet the nutritional athletic recommendations. 30% were ↓ 30 kcal.kgFFM⁻¹. dav⁻¹.
Gogojewicz et al. [48]	62 male and female CrossFit* participants; training experience: > 6 mths.	Poland	Food diary	 The energy intake and expenditure of male participants were 2265 ± 417 kcal.day⁻¹ and 2828 ± 316 kcal.day⁻¹. Corresponding values for females were 1736 ± 407 kcal.day⁻¹ and 2598 ± 286 kcal.day⁻¹. The protein values were similar in male and female groups (1.6 g.kg⁻¹.day⁻¹). Females ingested 3.9 g.kg⁻¹.day⁻¹ of CHO whilst males ingested 3.3 g.kg⁻¹.day⁻¹. The % of fat intake was 30% in both groups. Among females, ↓ calcium, iron, zinc, acid folic and ↑ phorehours.
Zaykova [49]	25 male and female professional CrossFit® participants; training experience: 3.5 yrs	Bulgaria	Questionnaire	 Daily energy intake ↑ than daily energy expenditure. Only five participants ingested >2.0 a ka⁻¹·dav⁻¹ of protein.
Brescansin et al. [50]	30 CrossFit [®] male and female participants (29.4 ± 9.1 yrs); training experience: NR.	Brazil	Food frequency questionnaire	 - 33.3% of participants use supplements. - Milk and diary products, eggs, legumes, cereals, fruits were consumed at least once a week.
Terry [51]	21 male and female CrossFit® participants; training experience: NR.	US	Food frequency questionnaire	 Female CrossFit[®] participants had ↓ calcium intake. Male CrossFit[®] participants had ↓ calcium and magnesium intake
Fayad [52]	15 male and female CrossFit® athletes (25–41 yrs); training experience: NR.	Brazil	24 hours recall	 60% of athletes ingested supplements. Whey protein and creatine were used by 53% and 26% of athletes, respectively. Mean energy intake was 1739 kcal.day⁻¹ (range: 835–1739 kcal.day⁻¹). Relative intake of CHO was 2.3 g.kg⁻¹·day⁻¹. 67% of athletes ingested >1.7 g.kg⁻¹·day⁻¹ of protein.
Lins et al. [53]	50 male and female CrossFit [®] participants (30.2 ± 5.6 yrs); training experience: < 1 mth to >1 yr.	Brazil	Questionnaire	 80% of participants ingested supplements. All participants that ingested supplements used whey protein, creatine, glutamine and BCAAs.

Table 1. (Continued).

(Continued)

-	•			
Study	Sample characteristics	Country	Methodology – nutritional/ergogenic aids issues	Main results
Pacheco et al. [54]	10 male and female CrossFit [®] participants (27.9 ± 7.4 yrs); training experience: NR.	Portugal	Food frequency questionnaire	 CrossFit[®] participants tend to consume ↓ dairy products, vegetables, legumes and sweets and pastries.
Bueno et al. [55]	10 male CrossFit® participants (18–50 yrs); training experience: > 6 months.	Brazil	Food diary	 The median energy intake 1409 kcal.day⁻¹. The median relative intake of protein, CHO and fats were 4.0 g.kg⁻¹·day⁻¹, 11.5 g.kg⁻¹·day⁻¹, 2.0 g.kg⁻¹·day⁻¹, respectively. Micronutrients (vitamins C, E, K) were ↓ the recommendations whilst B6 and B12 were ↑. Calcium and magnesium were ↓ the recommendations and iron and zinc ↑.
Freitas et al. [56]	88 male and female CrossFit [®] athletes; training experience: < 1 yr to >3 yrs.	Portugal	Questionnaire	 − 76.1% of CrossFit® athletes ingested at least one supplement. − Protein, creatine, and BCAAs are ↑ by CrossFit® athletes.

Table 1. (Continued).

tended to decrease from 2019 to 2024 (Figures 4 and 5). Protein intake has remained reasonably stable over the last five years (Figure 6).

3.3.2. Percentage of use of ergogenic aids

The prevalence of supplement use ranged from 25% to 98%. The mean prevalence was 72% (95% CI: 63% to 79%). The I^2 was 93%, which indicates that 93% of the observed variance in effects is real (Figure 7). Creatine (range: 26% to 61%) and protein (range: 100% to 28%) were the most prevalent ergogenic aids ingested by CrossFit[®] participants in seven studies [21,38–40,44,46,50,52,53,56].

3.3.3. Dietary interventions

Three different diets were considered in the analysis: high CHO [58], vegan [59], and Mediterranean [62]. Positive Hedges's g values indicate the benefits of CHO diets. Three studies provided data on the effects of diets with a greater percentage of CHO than that of customary diets on CrossFit[®] performance (pooled n = 31; Figure 8). The effect of CHO diets on CrossFit[®] performance was significantly moderate (Hedge's g = 0.487; 95% CI: 0.110 to 0.886). No trimmed studies were identified, and Egger's regression intercept was nonsignificant (Egger's intercept = 5.44, p = 0.563), indicating no risk of publication bias. Heterogeneity was low. Two studies with different designs (i.e. crossover and parallel) investigated the effects of a ketogenic diet on aerobic outputs assessed in the laboratory [61] and CrossFit[®] performance [60].

Data for the ketogenic diet were limited to two studies [60,61]. The former tested the effects of diet on laboratory outcomes (\dot{VO}_{2max}), time to exhaustion, and maximal power),

t2	
Ē	
ğ	
. <u> </u>	
Ē	
al	
<u></u>	
Ë	
Ч	
ŝ	
14	
2	
.⊆	
S	
p	
Ξ	
eU	
ž	
ē	
ъ	
1	
ŝ	
Ę,	
:₩	
5	
Ö	
ts	
2	
Ť	
Ð	
e	
÷	
σ	
é	
÷≣	
Ľ	
×	
e U	
at	
ቲ	
Ś	
ie.	
P	
E.	
2	
e	
9	
Ца	
-	

	Main results	 Blood lactate ↑ eating trial at baseline but, at post-exercise was ↑ fasting trial. → RPE, handgrip strength, jumping, CrossFit[®] performance time. 	 Relative peak power 1 in both groups. Maximal speed 1 in mediterranean diet. Time to attain peak power 1 in mediterranean diet. Power drop 1 in mediterranean diet. Jump height and time 1 in Mediterranean diet. Fran performance and chin-up test 1 in Morth crouns. 	 Number of repetitions on deadlift Number of repetitions on deadlift and squat ↑ in both groups. → fight gone bad performance. 	 	(Continued)
	Time	24 hours	8 weeks	4 weeks	4 weeks	
Diet	Control	Non-fasting: meal prior 2–3 hours before the exercise (50% CHO, 25% fats, 25% protein)	Control group: habitual diet	Mixed diet (adjustments of macronutrients – CHO: 297.7 \pm 53.9 g.kg ⁻¹ ; protein: 130.1 \pm 23.7 g.kg ⁻¹ ; fat: 98.9 \pm 12.7 g.kg ⁻¹)	Customory diet	
	Intervention	Fasting: consumed only water	Mediterranean diet (50% CHO, proteins were calculated based on training section, fats were distributed for the remaining calories)	Vegan diet (adjustments of macronutrients – CHO: 395.3 ± 63.1 g,kg ⁻¹ ; protein: 113.7 ± 23.4 g,kg ⁻¹ ; fat: 62.3 ± 21.2 g,kq ⁻¹)	Ketogenič diet ats: >75%, protein: 1.7 g.kg ^{−1} , CHO: ≤ 5%)	
Study	design	Crossover	Parallel	Parallel	Crossover (f	
	Sample characteristics	11 female CrossFit [®] participants (30.9 ± 3.4 yrs); training experience: > 2 yrs.	22 male and female CrossFit [®] participants; training experience: > 1 yr.	20 male and female CrossFit [®] participants; training experience: > 1 vr.	22 male and female CrossFit* participants; training experience: > 2 yrs.	
	Study	Eroglu et al. [57]	Ficarra et al. [58]	Durkalec- Michalski et al. [59]	Durkalec- Michalski et al. [60]	

12 🕑 D. V. MARTINHO ET AL.

Table 2. (Continued).	

		Study		Diet		
Study	Sample characteristics	design	Intervention	Control	Time	Main results
Kephart et al. [61]	12 male and female CrossFit® participants; training experience: NR.	Parallel	Ketogenic diet	Customary diet	12 weeks	 → maximal repetition of squat and power clean, push-ups, 400- m running in both groups. - Customary diet ↑ maximal oxvoren untake
Escobar et al. [62]	18 male and female CrossFit® participants; training experience: NR.	Parallel	CHO: 6–8 g.kg ^{–1} day ^{–1}	Lower CHO: < 6 g.kg ⁻¹ 'day ⁻¹	3 days	 CrossFit* performance ↑ CHO CrossFit* performance ↑ CHO diet. ← maximal oxygen uptake, blood Llacrate in both orous.
Gregory et al. [63]	27 male and female CrossFit participants [®] ; training experience: > 1 month.	Parallel	Ketogenic diet (CHO <50 grams per day)	Customary diet	6 weeks	- Ketogenic diet and customary diet \leftrightarrow physical performance.

amnle characteristics	Study desirin	Summement	Intervention	Timing	Main findings
g yrs.	Double- blind crossover trial	Cyclodextrin (CHO derived from plant starch).	309	During	
ale tes; ence:	Double- blind crossover trial	Caffeine	3 mg.kg _{BM} ⁻¹ , 6 mg. kg _{BM} ⁻¹ , 9 mg.kg _{BM} ⁻¹	70 minutes prior to the test	 → of CrossFit* performance (number of repetitions in both groups). - 6 mg/kg_{BM}⁻¹ CAF ↑ number of repetitions. - 6 mg/kg_{BM}⁻¹ CAF ↑ number of number of repetitions. - 6 mg/kg_{BM}⁻¹ in comparison to CAF 3 mg. kg_{BM}⁻¹ - 6 mg/kg_{BM}⁻¹ CAF ↓ reaction and more rimes
ale en ce:	Double- blind crossover trial trial	Sodium bicarbonate	0.15 gkg _{FFM} -', 0.25 gkg _{FFM} -', 0.35 gkg _{FFM} -1	2 hours prior to exercise	 1 blood pH 0.35 g/sg r_{FM} ⁻¹ and 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ to the baseline values at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood pH 0.35 g/sg r_{FM} ⁻¹ NaHCO₃ at recovery. 1 blood cells, than control at recovery. 1 blood cells than control at recovery. 2 blood st recovery. 2 blood cells than control at recovery. 2 blood cells than control at recovery.

(Continued)

Table 3. Ergogenic aids interventions in CrossFit® participants.

Sample characteristics a 17 advanced male [67] CrossFit* trained athletes; training experience: > 1 yr. [68] participants*, training experience: > 2 yrs. [69] 43 male recreationally training experience: > 1 yr. [69] and regularly training experience: > 1 yr. [69] and regularly trained CrossFit* participants; training experience: > 1 yr. et al. 9 male CrossFit* participants;	Study design Double- blind crossover trial blind crossover trial blind crossover trial	Sodium bicarbonate Caffeine + sodium bicarbonate Betaine CHO	Intervention Dosage 0.30 g'kg ⁻¹ ingested with the meal with the meal 0.1 g'kg ⁻¹ (CAF) + 0.1 g'kg ⁻¹ (NaHCO ₃) 2.5 g'kg ⁻¹ , 5.0 g'kg ⁻¹ 2.5 g'kg ⁻¹ , 5.0 g'kg ⁻¹ 1.0 g'kg ⁻¹ grape + banana juice; 1.0	Timing 2 hours prior to exercise 20, 90, 60 minutes before the protocol 3 weeks: 2.5 g'kg ⁻¹ – three capsules in the morning and two in the evening; 5.0 g'kg ⁻¹ – four capsules in the afternoon, and three in the afternoon, and three in the evening 60 minutes before the protocol	Main findings - NaHCO ₃ ↑ the performance in Fran CrossFit* challenge. - Performance in 500 m rowing and jumping ↔ - NaHCO ₃ ↑ blood pH in comparison to placebo. - No differences between groups were found for lactate, heart rate and RPE. - No significant effects of - No significant effects of - HR _{max} ↑ CAF and CAF + NaHCO ₃ in comparison to control group. - RPE during Cindy challenge ↓ CAF compared to NaHCO ₃ , CAF + NaHCO ₃ and control group. - RPE during Cindy challenge ↓ CAF compared to NaHCO ₃ , CAF + NaHCO ₃ and control group. - RPE during Cone Bad performance but, no significant differences were found compared to placebo group. - ↔ performance in Wingate test in both groups.
training experience: > 3 months. 17 male trained CrossFit [®] participants; training experience: > 1 yr. I. 11 female CrossFit [®] athletes; training experience: > 2 yrs.	Double- blind crossover trial Double- blind crossover trial	Capsiate Sodium bicarbonate	g kg ⁻ Whey Maximize 12 mg, 24 mg 0.3 g kg ⁻¹	45 minutes before the protocol Individualized approach	 Performance, HR and RPE was ↔ in CAP and placebo. NaHCO₃ † 2.2% rowing performance in comparison to placebo group. Mean power ouput ↑ NaHCO₃ (<i>p</i> < 0.05).

JOURNAL OF THE INTERNATIONAL SOCIETY OF SPORTS NUTRITION 🕒 15

ale CrossFit® articipants;	design Double- blind	Supplement Sodium bicarbonate	Intervention Dosage 0.3 g'kg ⁻¹	Timing 90 ⁻ 150 minutes prior to the test	Main findings – No effects of NaHCO ₃ on RPE.
berience:	parallel trial Double- blind crossover trial	Caffeine	6 mg kg ⁻¹	60 minutes prior to the protocols	 CAF ↑ local muscular endurance and number of repetitions. Squat maximum repetition was ↔ in both groups.
l female its; xperience: hs	Double- blind crossover trial	Beta-alanine	6.4 g.day ⁻¹	28 days. 4 times per day: 3 hours of interval (especially after meals)	 No differences between groups in performance, RPE, sleep quality, mood and well-being.
male nts; experience:	Parallel single- blind trial	Tribulus terrestris	770 mg	6 weeks: 30 minutes prior to CrossFit® workout	 Muscle biomarkers: ↑ LDH in placebo group whilst LDH ↓ in TT supplementation. Inflammatory markers: ↑ CRP placebo group whilst CRP ↓ in TT supplementation but without significance. Total Oxidant Status ↑ in TT group. Performance ← in both groups.
ssFit® training ce: > 6	Crossover single- blind trial	Carbohydrates + protein	500 ml carbohydrate- protein with different ratios (2:2, 3:1)	60 minutes prior to the test and immediately before	 Performance in CHO + protein and placebo groups was ↔ considering the number of repetitions in CrossFit[®] challenges.
male ants; experience: 6.	Parallel single- blind trial	Tribulus terrestris	770 mg	6 weeks	 T ↑ bench press performance significantly whilst the performance in control group ↔. CrossFit[®] performance, RPE, hormonal indicators ↔ in both groups.

Table 3. (Co	intinued).					
		Study		Intervention		
Study	Sample characteristics	design	Supplement	Dosage	Timing	Main findings
Toledo et al. [79]	30 male experienced CrossFit [®] participants; training experience: 2 2 vrs	Double- blind crossover trial	Sodium bicarbonate	0.3 g'kg ⁻¹	60 minutes prior to the test	– Number of repetitions, lactate, average and maximal heart rate, were \leftrightarrow in both conditions.
Stein et al. [80]	30 male CrossFit® participants; training experience: > 6 months.	Double- blind crossover trial	Caffeine	5 mg.kg ⁻¹	60 minutes prior to the test	– Number of repetitions and RPE \leftrightarrow in both groups.
Moro et al. [81]	29 male and female CrossFit [®] participants; training experience: > 6 months.	Double- blind crossover trial	Betaine	2.50 g (dosages of 1.25 g)	6 weeks: first dosage in morning, second dosage 60 minute before the work	 Significant improvements on back squat were noted for BET. No differences between groups were found for the remaining performance variables.
Ricordi et al. [82]	18 male and female beginner CrossFit [®] participants; training experience: > 6 months.	Double- blind crossover trial	Nitrate	10 g.kg ⁻¹	2 hours before CrossFit® practice	 CrossFit[®] performance was ↔ in both groups.
Garnacho- Castano et al. [83]	12 male CrossFit® participants; training experience: > 2 yrs.	Double- blind crossover trial	Beetroot juice	~808 mg	3 hours prior to exercise	 BJ ↑ the number of repetitions (after including resting time in the protocol). Jumping and blood lactate were ⇔in both conditions. SPO₂ decrease was ↑ BJ condition. Decline in jumping performance was ↑ BJ condition.
Fogaça et al. [84]	9 male experienced CrossFit® participants; training experience: 2 yrs.	Double- blind crossover trial	Caffeine	6 mg.kg ⁻¹	60 hours prior to exercise.	 Negligible differences between CAF and placebo groups were found for propulsive and peak power, bench press, jump squat, countermovement jumps and handgrip strength. RPE and DOMS were ← in both groups. Concentrations of CK and CPR ← in prost and 24 hours note workout

(Continued)

<mark>Table 3. (Cc</mark>	ontinued).			Intervention		
C41.141	Cample characteristics	Study	C	Intervention	Timine	mailen Baian
Durkalec- Michalski et al. [85]	21 recreational male and female CrossFit® participants; training experience: > 2 yrs.	Double- blind crossover trial	Sodium bicarbonate	Progressive dosages: 150 mg.kg ⁻¹ . Each 2 days, for 8 days, 1 25% of sodium bicarbonate dosage.	10 days: 3 doses per day. Training days was taken 1.5 hours before training sessions.	- Performance in FGB ↑ when participants ingested NaHCO ₃ . - Derived from incremental cycling test, maximum workload and HR were \leftrightarrow in both groups. - Workload at ventilatory threshold was 4.6% higher ($p < 0.05$) in NaHCO ₃ than placebo. - Time to attain ventilatory threshold was 5.1% higher ($p < 0.05$) in NaHCO ₃ than placebo. - HR at ventilatory threshold was significantly higher in NaHCO ₃ than placebo. - Time to attain ventilatory threshold was 5.1% higher ($p < 0.05$) in NaHCO ₃ than placebo. - Time to attain ventilatory threshold was 5.1% higher ($p < 0.05$) in NaHCO ₃ than placebo. - Pyruvate was significantly higher in NaHCO ₃ than placebo.
Rountree et al. [86]	8 CrossFit® trained males; training experience: > 3 months.	Double- blind crossover trial	СНО	16 g	Immediately prior and during exercise.	 Performance was ↔ in FGB on both trials.
Howarth et al. [87]	11 CrossFit® athletes; training experience: NR.	Crossover trial	СНО	Ad libitium	During the workout.	– Performance and RPE were \leftrightarrow in both groups.
Kramer et al. [88]	12 male CrossFit [®] athletes; training experience: > 4 months.	Double- blind crossover trial	Nitrate	8 mmol·d	6 days. Two times per day: morning and evening.	- No differences between groups were noted for CrossFit [®] performance (Grace and 2 km rowing). - The trend was maintained for physical performance with the exception of Wingate protocol. - Substantial improvements were noted for the NO_3^- group (6.6%) on peak power output.
						רמווווומבמי

Table 3. (Continued).

$\overline{\mathbf{O}}$
a.
≝
_
·=
<u>+</u>
_
0
· •
()
U.
9
9
<u>о</u>
<u>ю</u>
е З. (О
le 3. (C
ble 3. (C
able 3. (C
able 3. (C

		Study		Intervention		
Study	Sample characteristics	design	Supplement	Dosage	Timing	Main findings
Outlaw et al. [89]	 29 male and females CrossFit[®] participants; training experience: 6 months. 	Crossover	Pre-workout: tart cherry extract, beet root Extract, green tea and P black tea extract. Post-workout.	Pre-workout: 19 g ost-workout: females (20 g protein, 40 g CHO), males (40 g protein, 80 g CHO).	6 weeks: prior to the session.	 Performance on Wingate mean power and VO_{2max} † in the supplementation group. CrossFit[®] performance † on supplementation group but, without statistical sionificance.
Jacobs et al. [90]	 16 male and females CrossFit[®] participants; training experience: 6 months. 	Crossover	Commercial pre-workout supplement with beta-alanine, grape seed extract, coenzyme Q10, brown rice extract, natural caffeine, vitamins and minerals.	NR	20 minutes prior to Cindy workout.	- Cindy performance † in pre-workout supplement.



Figure 2. Mean energy intake of studies with CrossFit® participants.



Figure 3. Mean CHO and protein intake of studies with CrossFit® participants.



Figure 4. Mean values of energy intake and moving averages plot against the year of publication of the included studies (De Jesus et al. [34]; Rezende et al. [37]; Pearson and Jenkins [41]; gonçalves et al. [42]; Vieira et al. [45]; Gogojewicz et al [48]; Zaykova [49]).



Figure 5. Mean values of CHO intake and moving averages plot against the year of publication of the included studies (De Jesus et al. [34]; Pearson and Jenkins [41]; gonçalves et al. [42]; Vieira et al. [45]; Gogojewicz et al. [48]; Zaykova [49]; Fayad [52]).



Figure 6. Mean values of protein intake and moving averages plot against the year of publication of the included studies (De Jesus et al. [34]; Rezende et al. [37]; Pearson and Jenkins [41]; gonçalves et al. [42]; Vieira et al. [45]; Gogojewicz et al [48]; Zaykova [49]; Fayad [52]).

Study name				Event rate and 95% CI
	Event rate	Lower limit	Upper limit	
Brustolin et al.[35]	0,250	0,108	0,478	=-
Dos Santos et al. [36]	0,777	0,690	0,844	
Faria et al. [38]	0,752	0,664	0,823	
Comerlatto et al. [39]	0,631	0,565	0,693	
Filho et al. [40]	0,981	0,943	0,994	
Brisebois et al. [44]	0,820	0,805	0,834	
Higino & Freintas [46]	0,711	0,668	0,751	
Lins et al. [53]	0,800	0,667	0,889	
Fayad [52]	0,600	0,297	0,842	│ │ │ →■─│
Brescansin et al.[50]	0,333	0,190	0,516	│ │ │ -■┤ │
Freitas et al. [56]	0,761	0,662	0,839	
	0,721	0,633	0,794	
				-1,00 -0,50 0,00 0,50 1,00

Figure 7. Meta-analysis of the use of ergogenic aids among CrossFit® participants.

whereas the latter focused on CrossFit[®] performance (maximal repetition squat, maximal repetition power clean, number of push-ups, 400-m running). The mean differences between the ketogenic and customary diets did not indicate a positive effect of the ketogenic diet on performance. Given the differences in the outcomes between studies, conducting a meta-analysis for the ketogenic diet was impossible.



Figure 8. Forest plot of effects of CrossFit[®] performance after the implementation of CHO or customary diets. The black diamond reflects the overall result.

3.3.4. Ergogenic aid interventions

Considering the number of studies and outcomes analyzed, the meta-analysis was feasible for CHO and caffeine.

One study of caffeine [65] reported multiple comparisons (low-dose vs. medium-dose vs. high-dose vs. placebo) for three outcomes of CrossFit[®] performance (repetitions on Fight Gone Bad round 1, repetitions on Fight Gone Bad round 2, repetitions on Fight Gone Bad round 1, repetitions on Fight Gone Bad round 3). The average value of the medium dose was considered for the analyses, and the effect sizes of three rounds collapsed. Four crossover studies examined the impact of caffeine ingestion prior to the workout on CrossFit[®] performance (pooled n = 73). Overall, the effects of caffeine on performance are moderate (Hedges' g = 0.371) but not significant. Two trimmed studies were identified, and the adjusted Hedge's value decreased to 0.144 (95% Cl: -0.409 to 0.696), which is interpreted as a small effect. Egger's regression intercept did not identify the risk of biased publication (Egger's intercept = 2.506, p = 0.05) (Figure 9).

Three crossover studies investigated the effects of CHO intake prior to and/or during the workout on CrossFit[®] performance (pooled n = 40). CHO ingested did not affect physical performance (Hedges' g = -0.130; 95% CI: -0.652 to 0.391, p = 0.624), as shown in Figure 10. One trimmed study was identified when the Hedge's g values were plotted against the standard error. The adjusted Hedges' value for random effects was 0.063 (95%



Figure 9. Forest plot of effects of CrossFit[®] performance considering CHO or placebo intake prior or during the workout. The black diamond reflects the overall result.



Figure 10. Forest plot of effects of CrossFit[®] performance considering caffeine or placebo intake prior to the workout. The black diamond reflects the overall result. AMRAP (as many rounds as possible).

CI: -0.492 to 0.613). The risk of publication bias was identified for CHO intake (Egger's intercept = 5.489, p = 0.003).

Sodium bicarbonate supplementation prior to CrossFit[®] performance was examined in three studies [67,72,79], but the findings across these studies were inconsistent. One study reported no significant differences in Fran time or the time to complete the 500-m rowing test between the sodium bicarbonate and placebo groups [67]. However, sodium bicarbonate supplementation positively affected both time and power during the 2 km rowing test [72] as well as several repetitions performed during Cindy's workout [79].

Scientific evidence about these and other ergogenic aids used near training or competition (e.g. nitrate, beetroot juice) to optimize performance is limited. Only three studies focused on the impact of ergogenic aids on recovery [66,76,84].

3.3.5. Risk of bias

D. V. MARTINHO ET AL.

24

The risk of bias in each study indicated that the power sampling calculation was not reported in 19 studies. Twenty-two studies were cross-sectional, the exposures were not measured over time, and most assessors were aware of the participants' exposure (18 studies). In five studies, the blinding of assessors could not be determined. The PEDro scale of the interventional studies ranged from 5 to 11 points. Sixteen studies did not present or found significant differences between groups at baseline, which can influence the outcomes, and 11 records did not clearly explain the eligibility criteria used to recruit CrossFit[®] participants (Supplementary Material 2). Overall, 19 studies were classified as excellent, 9 studies as good and 4 studies as fair.

4. Discussion

This review summarizes the evidence concerning nutrition and supplementation use among CrossFit[®] participants and identifies potential gaps to guide future research. Relevant findings emerged from the current scoping review: (1) energy intake in CrossFit[®] participants is substantially lower in females than in males, although studies focused exclusively on females are limited; (2) the intake of CHO is below the recommendations for athletes, whereas the ingestion of protein is adequate; (3) CrossFit[®] participants tend to neglect CHO intake; (4) the use of ergogenic aids is considerable, with creatine and protein being the most reported supplements to optimize performance and recovery, respectively; (5) analyses of diets with higher contents of CHO or fat (i.e. a ketogenic diet) show that CHO has a moderate effect on CrossFit[®] performance, whereas a ketogenic diet does not affect performance; (6) considerable research using different supplements has been conducted in CrossFit[®] participants; however, only the data on caffeine and CHO are to generate an effect size on performance; and (7) the evidence concerning the effects of supplements on recovery is limited and warrants future research.

4.1. Energy and nutritional intake

Achieving adequate daily energy and nutrient requirements in sports is crucial for optimizing exercise performance, manipulating body composition, and facilitating recovery [18]. However, the definition of guidelines for CrossFit[®] practitioners is particularly challenging because only one study reported the total daily energy expenditure based on data derived from a heart rate monitoring device [48]. The latter study estimated daily energy expenditures of 2598 kcal.day⁻¹ and 2828 kcal.day⁻¹ for female and male participants, respectively [48]. Compared with the overall mean daily energy intake found in the current review, we report negative energy balances of -468 kcal.day⁻¹ and -852kcal.day⁻¹ for males and females, respectively. A negative energy balance was noted in endurance runners [91], but these values could also be explained by the limitations of the instruments used to estimate energy intake [92,93]. A meta-analysis of 11 studies used doubly labeled water to examine the accuracy of self-reported energy intake in athletes and reported that daily energy intake was underreported by 19% [93]. The studies reporting energy intake included in the present review used 24-hour recall [34,37,52], food diaries [42,45,48] or questionnaires [41]. Taking into account the daily energy expenditure reported in Polish CrossFit[®] participants as a reference [48], adjusting the mean total energy intake of the current review by 19% results in an energy balance of -20kcal for males, whereas a significant negative energy balance persists for females (-520 kcal). These findings suggest that individuals who follow female CrossFit[®] participants or athletes must frequently assess their energy intake, energy expenditure, body weight, and body composition to avoid periods of chronic energy restriction.

CrossFit[®] participants did not meet the nutritional recommendations of CHO specified for athletes [13,18]. Considering the range of 6–12 g.kg^{-1.}day⁻¹ CHO, the mean values of the current review are below the lower limit of the guidelines [94]. Although the lower intake of CHO differs between sports [91,95], this issue is particularly alarming in CrossFit[®] because high-intensity workouts characterize sessions. Nevertheless, the lower values of CHO identified in this review were expected because the recommendations for CrossFit[®] participants promoted high protein intake and low CHO ingestion [96]. The decrease in CHO intake has become more pronounced over time. Thus, the impacts of nutritional guidelines for CrossFit[®] participants on muscle and hepatic glycogen need to be reviewed to optimize performance and recovery [20,97]. Given the variability in exercise load, volume and rest among CrossFit[®] workouts, the quantities of CHO that should be ingested in different workouts need to be determined. Using the same participants, the maximal oxygen uptakes for the Cindy (complete as many rounds as possible in 20 minutes: 5 pull-ups, 10 push-ups, 15 air squats) and Fran (complete 21-15-9

26 🛭 😔 D. V. MARTINHO ET AL.

repetitions for time: thrusters, pull-ups) workouts were 34 and 29 ml.kg⁻¹.min⁻¹, respectively [7], whereas the mean heart rate on a workout that focused on complete rounds for time was substantially higher than that of a challenge where participants should perform as many rounds as possible in 5 minutes [98]. The metabolic demands of different workouts vary significantly, and appropriate quantities of energy and CHO should be ingested based on the type of session and the phase of the season following the concept of nutritional periodization available for endurance and power sports [99–102]. The current review does not provide specific guidelines for CHO intake among CrossFit[®] participants. Given the high intensity of CrossFit[®] workouts, athletes should consider increasing their CHO consumption to optimize performance and recovery. Additionally, it is important for athletes to be familiar with adequate sources of CHO.

Regarding protein ingestion, the *International Society of Sports Nutrition* recommends intakes of 1.4–2.0 g.kg⁻¹ for maintaining and building muscle mass [103], whereas 1.6 g.kg⁻¹ is defined as the upper cutoff value to promote changes in fat-free mass [104]. The mean protein intake reported in this review is consistent with the recommended values. CrossFit[®] participants and nutritionists should be aware that greater fat mass can affect performance [105,106] and discriminate athletes by competitive level [105]. Additional questions, such as the type, timing, and distribution of protein intake across the day, warrant further research. Given the importance of protein intake, it is unsurprising that it has been reported as one of the most prevalent ergogenic aids ingested by CrossFit[®] participants.

4.2. Prevalence of ergogenic aids

The meta-analysis of the prevalence of ergogenic aid usage revealed an average intake of ~ 72%. The most reported supplements are protein and creatine, with the aim of optimizing recovery and performance. The benefits of creatine supplementation are associated with increased muscle phosphocreatine and regeneration of phosphocreatine during exercise recovery [107]. The short- and long-term benefits of creatine supplementation on strength and resistance activities are well documented in the literature, as are the timing and dosage necessary for athletes [108,109]. Therefore, it is unsurprising that creatine has been reported as one of the most preferred supplements used by CrossFit[®] participants. One potential disadvantage of creatine is in body weight due to intracellular water retention [107]; however, gains in body weight could be particularly beneficial in supporting the extreme loads imposed by CrossFit[®].

The type of protein ingested was reported in three studies [38–40], with whey protein being the most commonly used protein by CrossFit[®] participants. Whey protein is rapidly digestible and contains a high proportion of essential amino acids (including leucine), which determines the protein quality to optimize muscle protein synthesis and body composition [103,110]. The attainment of protein intake guidelines for resistance training and skeletal muscle hypertrophy adaptations, independent of the timing and quality, has not been compared between whey protein and other types of protein [111]. Therefore, supplementation with whey protein in conjunction with dietary sources of protein is central among CrossFit[®] practitioners [110,112].

4.3. Dietary interventions

The recommended dietary prescription for CrossFit[®] participants is to follow two different types of diets: the Zone and the Paleo diets. The Zone Diet is characterized by meals and snacks with a macronutrient distribution of 40% CHO, 30% protein, and 30% fat, whereas the Paleo Diet recommends the ingestion of fruits, vegetables, lean meats, and fish and the avoidance of dairy foods, legumes, and grains with a macronutrient distribution of 35–45% CHO, 20–35% protein, and the remaining percentage from fat. Both dietary approaches are characterized by a lower ingestion of CHO and an increased percentage of protein [113]. These diets lack a scientific basis for CrossFit[®]. As such, dietary prescriptions among CrossFit[®] participants need further evidence to support their use.

The Atkins diet is classified as a nonketogenic, low-carbohydrate diet, and the potential role of this dietary approach is to deplete glycogen stores, increase fat oxidation, and optimize gluconeogenesis [113]. The ketogenic diet, characterized by relatively high levels of fat intake, may impact on CrossFit[®] performance [60,61] via a mechanism similar to that of the Atkins diet: it minimizes glycogen stores, maximizes fat oxidation, and provides ketone bodies as a potential substrate for muscle tissue and the central nervous system [114]. Recently, the increase in fat oxidation stimulated by a ketogenic diet was demonstrated to be due to an increase in the oxygen cost (i.e. less efficiency) without increasing performance among elite racewalkers [115]. Because CHO metabolism results in a greater ratio of reducing NADH to FADH₂ than fat does, more energy is produced per unit of oxygen when CHO are used as fuel in the electron transport chain [116]. The presence of oxygen and other factors (e.g. availability of coenzyme A, carnitine palmitoyl transferase) are a determinant and a critical factor in explaining the shift of CHO and fats in endurance events [117]. Moreover, at high intensities, oxygen cannot be used; consequently, the metabolism of fat via beta-oxidation is not considered. Therefore, ketogenic diets for athletes in high-intensity sports, such as CrossFit[®], did not significantly improve performance outputs compared with those of the control groups [60,61]. Another issue associated with the ketogenic diet is the difficulty of participants adhering to the diet [118].

In contrast to a ketogenic diet, applying dietary prescriptions with a higher percentage of carbohydrates was moderately beneficial for CrossFit[®] performance outputs. Although the focus of a vegan diet is the amount of protein ingested $(1.5-2.0 \text{ g.kg}^{-1} \cdot \text{day}^{-1})$, the study that examined the effects of this type of diet on performance and biomarkers was classified as a diet with a relatively high content of CHO. CHO were reported to be ingested at 4.5–5.5 g.kg⁻¹ \cdot day for a vegan diet [60]. Diets focused predominantly on CHO appear to satisfy athletes' energy needs, support high-intensity activities, and improve body composition [113].

4.4. Acute strategies

Many supplements have been used before or during workouts by CrossFit[®] participants. Most supplements lack scientific evidence, and only two studies present sufficient data on the potential effects on performance: CHO and caffeine.

28 🕒 D. V. MARTINHO ET AL.

The ergogenic effects of carbohydrates have been considered during exercise because they represent a nutritional strategy that delays glycogen depletion and improves performance [119,120], and the recommendations for CHO ingestion during exercise consider the time spent exercising. For example, in a soccer match, the ingestion of 30–60 g.hour⁻¹ or 60 g of carbohydrates before each half is recommended [121,122], whereas in endurance sports, 60 g.hour⁻¹ is advised in efforts >2.5 hours [20]. This review did not confirm the effect of CHO on CrossFit® performance. Two studies [64,86] examined the effects of carbohydrate ingestion in short-performance protocols. These results suggest that athletes may use muscle and hepatic glycogen from carbohydrates ingested during the hours before exercise to complete workouts. In other words, the protocols used did not allow the depletion of glycogen stores. A third study considered a long workout [86]; however, the intake of carbohydrates occurred during the workout. Consequently, future studies should investigate the ingestion of carbohydrates during CrossFit[®] challenges, taking into account the duration of exercise and intensity. Additional guestions about carbohydrate supplementation include tolerance to higher quantities [20,123] and practical application during high-intensity workouts. Consequently, further research should consider testing the impact of carbohydrates within training sessions and not solely on a specific part of the workout or between competitions that occur on the same day. In fact, the effects of carbohydrates on resistance training tend to be positive, particularly when training sessions exceed 45 minutes [124]. Carbohydrate intake can be especially beneficial for athletes who engage in longer training periods or have multiple training sessions in a single day.

Caffeine is a popular ergogenic aid in the context of sports [125]. The effect of caffeine ingestion on CrossFit[®] performance was small and nonsignificant. Nevertheless, the results were consistent with a previous meta-analysis that examined the effects of the ingestion of moderate doses (3–6 mg.kg⁻¹) on the mean power output and time to complete a trial. However, both analyses in this study noted significant effects [126]. A combination of 20 studies also revealed that caffeine ingestion significantly improved strength (1 repetition maximum test) and power (vertical jump) outputs [127]. A narrative review also highlighted the positive effect of caffeine intake on muscular endurance, strength, and power in a resistance context [128]. The nonsignificant effects found in this review could be explained by the demands of CrossFit[®], which requires endurance, strength, and power capacities.

The study design, dosage, timing, and formula can also affect the caffeine response [129]. The placebo effect was noted in a study of 48 cyclists, with those who received a placebo and thought they ingested caffeine, improving exercise performance [130]. Consequently, ensuring the blinding of participants, which was guaranteed in the four studies included in this meta-analysis, is important. Although the timing and dosage are two factors that need further investigation, the quantity and timing of caffeine ingested by samples included in the meta-analysis ranged from 3 to 6 mg.kg⁻¹ and from 60 to 70 minutes, respectively. The dosage followed the recommendations for caffeine intake, with lower dosages (2 mg.kg⁻¹) having no effect on exercise performance [131]. Higher dosages of caffeine should be considered in maximal repetition efforts [132]. This issue warrants further study in CrossFit[®] participants because muscular strength and endurance are central to improving performance. The ingestion of caffeine 60 minutes before exercise is widely accepted to

achieve the ergogenic effects of this supplement [133]. Nevertheless, the caffeine response depends on genetic and epigenetic factors (e.g. sex, habitual caffeine use, smoking) [134].

4.5. Limitations and future research

The present study provides a broad overview of the nutritional evidence in CrossFit[®], but some limitations should be recognized: (1) Exercise and daily energy expenditure reference values are scarce among CrossFit[®] participants. Nevertheless, the assessment of exercise energy expenditure is challenging because the error associated with wearable monitors ranges from 15.1% to 57.0% [135]. (2) The meta-analysis testing the effects of carbohydrates and caffeine used a limited number of studies. (3) Multiple ergogenic aids were used among CrossFit[®] participants, but robust conclusions could not be drawn given the variability in outcomes and interventions. For example, a well-designed, double-blind, randomized controlled trial revealed beneficial effects of sodium bicarbonate on CrossFit® performance (time trial on a 2 km rowing test); however, the data are limited to one study, and the authors determined the peak blood sodium bicarbonate for each participant [72]. Among the 17 participants, sodium bicarbonate improved the Fran time and time to complete 500 m rowing [67], whereas no effects were found in performance for Cindy's workout [79]. These inconsistent findings indicate that the differences in the effects of sodium bicarbonate on performance and recovery between training sessions or competitions need further investigation. (4) Few studies have focused on recovery, and this issue needs particular attention given the high volume and intensity of training exposure in CrossFit[®]. The studies included in this review have limitations that should not be ignored: sample size and the definition of a CrossFit[®] participant or athlete. The training characteristics of individuals from different competitive levels are not equivalent (see Supplementary Material 3 to compare training sessions of beginner, intermediate, and advanced practitioners of CrossFit[®]). This issue must be considered when the eligibility criteria are defined [136].

5. Conclusions and practical applications

The present scoping review provides new insights into nutrition and CrossFit[®], which might yield changes in practice. Although the ingestion of protein was appropriate, CHO consumption was below the recommended limit for athletes, and CHO consumption has decreased in recent years. Nutritionists need to educate coaches and athletes about the importance of CHO in high-intensity sporting activities, such as CrossFit[®]. Diets with high CHO values show performance benefits, and recommendations surrounding the Zone or Paleo diet need to be revised. CHO intake is essential for optimizing performance, whereas protein intake can influence body composition, which, in turn, may affect performance. Therefore, athletes should pay special attention to the ingestion of these two macronutrients by managing their intake effectively. Data available on the ketogenic diet remain limited, although the existing studies do not demonstrate performance benefits. The use of nutritional aids is common among CrossFit[®] participants, with creatine and protein being the most reported. The benefits of these nutritional aids for performance and recovery are well documented, and consequently, they should be incorporated into the regimen of CrossFit[®] participants. Given the variability of CrossFit[®] workouts, future studies examining the long-term

30 🕒 D. V. MARTINHO ET AL.

impact of creatine on performance and protein on recovery are needed. When consumed acutely, CHO had no effect on performance; however, ingesting 6 mg.kg⁻¹ caffeine 60–70 minutes before exercise had a small effect on performance. In summary, monitoring carbohydrate and protein intake is essential to determine whether CrossFit[®] participants meet the recommended guidelines. Regarding the scientific evidence surrounding supplements, both creatine and protein should be consumed, while caffeine may be utilized prior to training sessions.

Disclosure statement

No potential conflict of interest was reported by the author(s).

Funding

This research was funded by the Portuguese Recovery and Resilience Program (PRR), IAPMEI/ANI/ FCT under the Agenda [C645022399–00000057] (eGamesLab).

ORCID

Diogo V. Martinho () http://orcid.org/0000-0003-0825-4032 André Rebelo () http://orcid.org/0000-0003-2441-9167 Filipe Manuel Clemente () http://orcid.org/0000-0001-9813-2842 Élvio R. Gouveia () http://orcid.org/0000-0003-0927-692X Adam Field () http://orcid.org/0000-0002-2600-6182 Juliano Casonatto () http://orcid.org/0000-0001-5397-5694 Daniel van den Hoek () http://orcid.org/0000-0001-8335-1620 Krzysztof Durkalec-Michalski () http://orcid.org/0000-0002-5041-2981 Michael J. Ormsbee () http://orcid.org/0000-0002-1939-7656 Hugo Sarmento () http://orcid.org/0000-0001-8681-0642

Data availability statement

The data that support the findings of this study are available from the corresponding author, DVM, upon reasonable request.

References

- 1. CrossFit[®] Growth Statistics: Global Reach, Revenue and Participation Soar. https://world metrics.org/crossfit-growth-statistics/consulted2024
- Claudino JG, Gabbett TJ, Bourgeois F, et al. CrossFit overview: systematic review and meta-analysis. Sports Med Open. 2018;4(1):11. doi: 10.1186/s40798-018-0124-5 PMID: 29484512; PMCID: PMC5826907.
- 3. Schlegel P. CrossFit[®] training strategies from the perspective of concurrent training: a systematic review. J Sports Sci Med. 2020;19(4):670–680. PMID: 33239940; PMCID: PMC7675627.
- 4. Rios M, Becker KM, Monteiro AS, et al. Effect of the fran CrossFit workout on oxygen uptake kinetics, energetics, and postexercise muscle function in trained CrossFitters. Int J Sports Physiol Perform. 2024;19(3):299–306. doi: 10.1123/ijspp.2023-0201 PMID: 38194958.

- Mangine GT, Cebulla B, Feito Y. Normative values for self-reported benchmark workout scores in CrossFit[®] practitioners. Sports Med Open. 2018;4(1):39. doi: 10.1186/s40798-018-0156-x PMID: 30128825; PMCID: PMC6102159.
- 6. Schlegel P, Režný L, Fialová D. Pilot study: performance-ranking relationship analysis in Czech crossfiters. J Hum Sport Exerc. 2021;16(1):187–198. doi: 10.14198/jhse.2021.161.17
- Fernández JF, Solana RS, Moya D, et al. Acute physiological responses during CrossFit[®] workouts. Eur J Hum Mov. 2015;35:114–124.
- Rios M, Zacca R, Azevedo R, et al. Bioenergetic analysis and fatigue assessment during the fran workout in experienced crossfitters. Int J Sports Physiol Perform. 2023;18(7):786–792. doi: 10. 1123/ijspp.2022-0411 PMID: 37225165.
- de Sousa Neto IV, de Sousa NMF, Neto FR, et al. Time course of recovery following CrossFit[®] Karen benchmark workout in trained men. Front Physiol. 2022;13:899652. doi: 10.3389/fphys. 2022.899652 PMID: 36060700; PMCID: PMC9438894.
- Forte LDM, Freire YGC, Júnior JSDS, et al. Physiological responses after two different CrossFit workouts. Biol Sport. 2022;39(2):231–236. doi: 10.5114/biolsport.2021.102928 Epub 2021 Apr 9. PMID: 35309530; PMCID: PMC8919890.
- 11. Tibana RA, De Sousa NMF, Prestes J, et al. Lactate, heart rate and rating of perceived exertion responses to shorter and longer duration CrossFit[®] training sessions. J Funct Morphol Kinesiol. 2018;3(4):60. doi: 10.3390/jfmk3040060 PMID: 33466988; PMCID: PMC7739245.
- Alsamir Tibana R, Manuel Frade de Sousa N, Prestes J, et al. Is perceived exertion a useful indicator of the metabolic and cardiovascular responses to a metabolic conditioning session of functional fitness? Sports (Basel). 2019;7(7):161. doi: 10.3390/sports7070161 PMID: 31277360; PMCID: PMC6681255.
- Thomas DT, Erdman KA, Burke LM. American college of sports medicine joint position statement. Nutrition and athletic performance. Med Sci Sports Exerc. 2016;48(3):543–568. doi: 10.1249/MSS.00000000000852 Erratum in: Med Sci Sports Exerc. 2017 Jan;49(1):222. PMID: 26891166.
- Gonzalez DE, McAllister MJ, Waldman HS, et al. International society of sports nutrition position stand: tactical athlete nutrition. J Int Soc Sports Nutr. 2022;19(1):267–315. doi: 10. 1080/15502783.2022.2086017 PMID: 35813846; PMCID: PMC9261739.
- Maxwell C, Ruth K, Friesen C. Sports nutrition knowledge, perceptions, resources, and advice given by certified CrossFit trainers. Sports (Basel). 2017;5(2):21. doi: 10.3390/sports5020021 PMID: 29910381; PMCID: PMC5968988.
- Coulston AM, Liu GC, Reaven GM. Plasma glucose, insulin and lipid responses to high-carbohydrate low-fat diets in normal humans. Metabolism. 1983;32(1):52–56. doi: 10. 1016/0026-0495(83)90155-5 PMID: 6336816.
- 17. Cordain L, Friel J. The paleo diet for athletes: the ancient nutritional formula for peak athletic performance. Harmony/Rodale/Convergent: Rodale Books; 2012.
- Burke LM, Cox GR, Culmmings NK, et al. Guidelines for daily carbohydrate intake: do athletes achieve them? Sports Med. 2001;31(4):267–299. doi: 10.2165/00007256-200131040-00003 PMID: 11310548.
- 19. Cermak NM, van Loon LJ. The use of carbohydrates during exercise as an ergogenic aid. Sports Med. 2013;43(11):1139–1155. doi: 10.1007/s40279-013-0079-0 PMID: 23846824.
- Jeukendrup A. A step towards personalized sports nutrition: carbohydrate intake during exercise. Sports Med. 2014;44(Suppl S1):S25–33. doi: 10.1007/s40279-014-0148-z PMID: 24791914; PMCID: PMC4008807.
- Santos Quaresma MvI D, Guazzelli Marques C, Nakamoto FP. Effects of diet interventions, dietary supplements, and performance-enhancing substances on the performance of CrossFit-trained individuals: a systematic review of clinical studies. Nutrition. 2021;82:110994. doi: 10.1016/j.nut.2020.110994 Epub 2020 Aug 28. PMID: 33051114.
- de Souza RAS, da Silva AG, de Souza MF, et al. A systematic review of CrossFit[®] workouts and dietary and supplementation interventions to Guide nutritional strategies and future research in CrossFit[®]. Int J Sport Nutr Exerc Metab. 2021;31(2):187–205. doi: 10.1123/ijsnem. 2020-0223 Epub 2021 Jan 29. PMID: 33513565.

- 32 🕒 D. V. MARTINHO ET AL.
 - 23. Higgins JP, Thomas J, Chandler J, et al. Cochrane handbook for systematic reviews of interventions. Chichester (UK): John Wiley & Sons; 2019.
 - Page MJ, McKenzie JE, Bossuyt PM, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. BMJ. 2021;372:n71. doi: 10.1136/bmj.n71 PMID: 33782057; PMCID: PMC8005924.
 - Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): checklist and explanation. Ann Intern Med. 2018;169(7):467–473. doi: 10.7326/M18-0850 Epub 2018 Sep 4. PMID: 30178033.
 - 26. Lefebvre C, Glanville J, Briscoe S, et al. Technical supplement to chapter 4: searching for and selecting studies. In: Higgins J; Thomas J; Chandler J; Cumpston M; Li T; Page M Welch V, editors. Cochrane handbook for systematic reviews of interventions version 6.4 (updated February 2024). Cochrane; 2024. Available from: www.training.cochrane.org/handbook
 - Wojtyniak JG, Britz H, Selzer D, et al. Data digitizing: accurate and precise data extraction for quantitative systems pharmacology and physiologically-based pharmacokinetic modeling. CPT Pharmacometrics Syst Pharmacol. 2020;9(6):322–331. doi: 10.1002/psp4.12511 Epub 2020 Jun 16. PMID: 32543786; PMCID: PMC7306621.
 - Hopkins WG, Marshall SW, Batterham AM, et al. Progressive statistics for studies in sports medicine and exercise science. Med Sci Sports Exerc. 2009;41(1):3–13. doi: 10.1249/MSS. 0b013e31818cb278 PMID: 19092709.
 - López-López JA, Page MJ, Lipsey MW, et al. Dealing with effect size multiplicity in systematic reviews and meta-analyses. Res Synth Methods. 2018;9(3):336–351. doi: 10.1002/jrsm.1310 Epub 2018 Jul 3. PMID: 29971966.
 - Higgins JP, Thompson SG. Quantifying heterogeneity in a meta-analysis. Stat Med. 2002;21 (11):1539–1558. doi: 10.1002/sim.1186 PMID: 12111919.
 - Duval S, Tweedie R. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. Biometrics. 2000;56(2):455–463. doi: 10.1111/j.0006-341x.2000.00455.x PMID: 10877304.
 - National Heart, Lung, and Blood Institute. Quality assessment tool for observational cohort and cross-sectional studies. 2019. https://www.nhlbi.nih.gov/health-topics/study-qualityassessment-tools
 - Cashin AG, McAuley JH. Clinimetrics: physiotherapy evidence database (PEDro) scale. J Physiother. 2020;66(1):59. doi: 10.1016/j.jphys.2019.08.005 Epub 2019 Sep 11. PMID: 31521549.
 - 34. de Jesus MGM, de Melo TD, Bassoli BK, et al. Nutritional intervention to improve anthropometric profile in CrossFit[®] athletes. Revista Brasileira De Nutricao Esportiva. 2024;18:23–30.
 - Brustolin MR, Souza CT, Souza DT, et al. Assessment of the pre- and post-workout dietary profile of Crossfit[®] practitioners. Rbne-Revista Brasileira De Nutricao Esportiva. 2024;18:13–22.
 - Santos Quaresma MvI D, Marques CG, Magalhães ACO, et al. Prevalence of dietary supplement consumption among CrossFit practitioners. Nutrire. 2023;48(1):9. doi: 10.1186/s41110-023-00193-z Epub 2023 Feb 13. PMID: 38625281; PMCID: PMC9924833.
 - Rezende CB, Junqueira ACP, Braga TM. Macronutrient intake, nutritional knowledge and motivation to practice physical activity in two modalities of physical activity. Rbne-Revista Brasileira De Nutricao Esportiva. 2023;17:488.
 - de Faria CHK, Yamaguchi GD. Use of nutritional ergogenic resources in CrossFit[®] practitioners. Rbne-Revista Brasileira De Nutricao Esportiva. 2023;17:101–108.
 - 39. Comerlatto V, Zanella PB, Hoefel AL. Crossfit[®] practitioners profile with regard to the prevalence of use of dietary supplements and anabolic androgenic steroids as ergogenic resources. Rbne-Revista Brasileira De Nutricao Esportiva. 2023;17(106):575–584.
 - 40. Rebouças Filho M, Mendonça LP. Perfil alimentar e utilização de suplementos nutricionais de praticantes de cross training do box Crossfit[®] sertão no município de mossoró. RBNE-Revista Brasileira de Nutrição Esportiva. 2022;16(101):466–474.
 - Pearson RC, Jenkins NT. Dietary intake of adults who participate in CrossFit[®] exercise regimens. Sports (Basel). 2022;10(3):38. doi: 10.3390/sports10030038 PMID: 35324647; PMCID: PMC8954808.

- 42. Gonçalves YT, Costa EE, Caetano M, et al. BODY COMPOSITION and DIETARY INTAKE in BRAZILIAN CROSSFIT® ATHLETES: a DESCRIPTIVE STUDY: 1979. Med Sci Sports Exerc. 2022;54(9S):590–590. doi: 10.1249/01.mss.0000882456.75459.cc
- Mesquita MHS, Cavalcanti DR. Food profile associated with the nutritional status of Crossfit[®] practitioners. Rbne-Revista Brasileira De Nutricao Esportiva. 2022;16:247–253.
- Brisebois M, Kramer S, Lindsay KG, et al. Dietary practices and supplement use among CrossFit[®] participants. J Int Soc Sports Nutr. 2022;19(1):316–335. doi: 10.1080/15502783. 2022.2086016 PMID: 35813850; PMCID: PMC9261745.
- Vieira IC, Soares Nunes AA, Iori M, et al. Dietary calories and macronutrients intake in Crossfit[®] practitioners. Med Sci Sports Exerc. 2021;53(8S):275–275. doi: 10.1249/01.mss.0000762284. 86696.ff
- 46. Higino DD, Freitas RF. Prevalência e fatores associados ao uso de suplementos alimentares e esteroides anabólicos androgênicos em praticantes de CrossFIT[®]. Rbne-Revista Brasileira De Nutricao Esportiva. 2021;15(90):9–23.
- 47. Kuch A. Recreational female CrossFit[®] athletes and low energy availability [master's dissertation]. South Dakota State University; 2021.
- Gogojewicz A, Śliwicka E, Durkalec-Michalski K. Assessment of dietary intake and nutritional status in CrossFit-trained individuals: a descriptive study. Int J Environ Res Public Health. 2020;17(13):4772. doi: 10.3390/ijerph17134772 PMID: 32630749; PMCID: PMC7369802.
- Zaykova D. Evaluation of diet of people training crossfit[®]. In: lancheva T, Djobova S, Kuleva M, editors. International scientific congress on applied sports sciences (Icass)/Balkan scientific congress on physical education. BULGARIA: Sports, Health, Sofia; 2019. p. 20–24.
- Brescansin BM, Naziazeno RFT, de Miranda TV. Análise do perfil alimentar de praticantes de CrossFit[®] na região de belém do pará. Rbne-Revista Brasileira De Nutricao Esportiva. 2019;13 (81):830–838.
- 51. Terry MS. The relationship of dietary intakes of calcium, magnesium, and zinc on Body composition in female and male athletes of various sports [master's dissertation]. Drexel Univ. 2019.
- 52. Fayad D. A influência da estratégia nutricional no rendimento de atletas competitivos de Crossfit[®] [Undergraduate thesis]. Centro Universitário de Brasília, Faculdade de Ciências da Educação E Saúde. 2019.
- Lins TCL, de Souza LPV. Dieta pré e pós treino em praticantes de crossfit[®]: um perfil qualitativo do consumo de alimentos e suplementos. Rbne-Revista Brasileira De Nutricao Esportiva. 2019;13(82):946–953.
- 54. Pacheco ASR. Nutrição e Crossfit[®]: características da realidade portuguesa [master's dissertation]. Faculdade de Medicina Universidade de Coimbra. 2018.
- 55. Bueno BA, Ribas MR, Bassan JC. Determination of micro and macronutrients intake for CrossFit[®] practitions. Rbne-Revista Brasileira De Nutricao Esportiva. 2016;10:579–586.
- 56. Freitas JCRDSO. Potencial ergogénico e uso da creatinina e da beta-alanina no contexto do crossFit[®] e da musculação [master's dissertation]. Escola Superior de Desporto E Lazer Inst Politécnico de Viana Do Castelo. 2016.
- Eroglu MN, Rodríguez-Longobardo C, Ramírez-Adrados A, et al. The effects of 24-h fasting on exercise performance and metabolic parameters in a Pilot study of female CrossFit athletes. Nutrients. 2023;15(22):4841. doi: 10.3390/nu15224841 PMID: 38004236; PMCID: PMC10674902.
- Ficarra S, Di Raimondo D, Navarra GA, et al. Effects of Mediterranean Diet combined with CrossFit training on trained Adults' performance and body composition. J Pers Med. 2022;12 (8):1238. doi: 10.3390/jpm12081238 PMID: 36013187; PMCID: PMC9409826.
- Durkalec-Michalski K, Domagalski A, Główka N, et al. Effect of a four-week vegan diet on performance, training efficiency and blood biochemical indices in CrossFit-trained participants. Nutrients. 2022;14(4):894. doi: 10.3390/nu14040894 PMID: 35215544; PMCID: PMC8878731.

34 🛭 😔 D. V. MARTINHO ET AL.

- 60. Durkalec-Michalski K, Nowaczyk PM, Główka N, et al. Is a four-week ketogenic Diet an effective nutritional strategy in CrossFit-trained female and male athletes? Nutrients. 2021;13(3):864. doi: 10.3390/nu13030864 PMID: 33800770; PMCID: PMC8001376.
- Kephart WC, Pledge CD, Roberson PA, et al. The three-month effects of a ketogenic Diet on Body composition, blood parameters, and performance metrics in CrossFit trainees: a Pilot study. Sports (Basel). 2018;6(1):1. doi: 10.3390/sports6010001 PMID: 29910305; PMCID: PMC5969192.
- Escobar KA, Morales J, Vandusseldorp TA. The effect of a moderately Low and high carbohydrate intake on crossfit performance. Int J Exerc Sci. 2016;9(3):460–470. doi: 10.70252/ IZLO1608 PMID: 27766133; PMCID: PMC5065325.
- Gregory RM. A low-carbohydrate ketogenic diet combined with six weeks of crossfit[®] training improves body composition and performance [master's dissertation]. James Madison Univ. 2016.
- Grijota FJ, Toro-Román V, Bartolomé I, et al. Acute effects of 30 g cyclodextrin intake during CrossFit[®] training on performance and fatigue. J Funct Morphol Kinesiol. 2024;9(1):27. doi: 10. 3390/jfmk9010027 PMID: 38390927; PMCID: PMC10885023.
- 65. Główka N, Malik J, Podgórski T, et al. The dose-dependent effect of caffeine supplementation on performance, reaction time and postural stability in CrossFit - a randomized placebo-controlled crossover trial. J Int Soc Sports Nutr. 2024;21(1):2301384. doi: 10.1080/ 15502783.2023.2301384 Epub 2024 Jan 16. PMID: 38226646; PMCID: PMC10795626.
- 66. Durkalec-Michalski K, Kamińska J, Saunders B, et al. Does sodium bicarbonate based extra-cellular buffering support reduce high intensity exercise-induced fatigue and enhance short-term recovery assessed by selected blood biochemical indices? Biol Sport. 2024;41 (1):17–27. doi: 10.5114/biolsport.2024.125591 Epub 2023 May 25. PMID: 38188117; PMCID: PMC10765444.
- 67. Silva de Souza RA, Barreto G, Alves Freire PA, et al. Sodium bicarbonate improved CrossFit® benchmark fran, but not subsequent 500 m rowing performance. Res Sports Med. 2024;32 (6):965–980. doi: 10.1080/15438627.2024.2324254 Epub 2024 Mar 14. PMID: 38482841.
- Ziyaiyan A, Shabkhiz F, Hofmeister M. Supplementation of caffeine and sodium bicarbonate together could not improve performance and performance-related factors in CrossFit participants: a randomized, double-blind, placebo-controlled study. J Int Soc Sports Nutr. 2023;20 (1):2206390. doi: 10.1080/15502783.2023.2206390 PMID: 37151086; PMCID: PMC10167875.
- Zawieja E, Durkalec-Michalski K, Sadowski M, et al. Betaine supplementation improves CrossFit performance and increases testosterone levels, but has no influence on Wingate power: randomized crossover trial. J Int Soc Sports Nutr. 2023;20(1):2231411. doi: 10.1080/ 15502783.2023.2231411 PMID: 37409757; PMCID: PMC10327519.
- 70. Mattos MES, Vieira Camargo MK, Santos JR, et al. Respostas bioquímicas e de percepção subjetiva de esforço pós suplementação alimentar em praticantes de Crossfit[®]: um estudo crossover. Rbne-Revista Brasileira De Nutricao Esportiva. 2023;17:705–713.
- Oliveira MDSM, Macêdo PR, Santana NL, et al. Dose-response of acute capsiate supplementation on muscle endurance performance during CrossFit[®] in trained men: a randomized, crossover and double-blind study. Nutr Metab (Lond). 2023;33:200215. doi: 10.1016/j.hnm. 2023.200215
- Martin RAXJ, Hilton NP, Sparks SA, et al. The effects of enteric-coated sodium bicarbonate supplementation on 2 km rowing performance in female CrossFit[®] athletes. Eur J Appl Physiol. 2023;123(6):1191–1198. doi: 10.1007/s00421-023-05140-4 Epub 2023 Jan 27. PMID: 36705750; PMCID: PMC10191925.
- 73. Gomes GC, Ribeiro P, Binoti ML. Effect of acute sodium bicarbonate supplementation on performance, subjective perception of effort and gastrointestinal discomfort in CrossFit[®] male practitioners. Rbne-Revista Brasileira De Nutricao Esportiva. 2023;17:439–449.
- 74. Caetano ML, Souza MLR, Loureiro LL, et al. The effects of acute caffeine supplementation on performance in trained CrossFit[®] athletes: a randomized, double-blind, placebo-controlled, and crossover trial. Sci Sports. 2023;38(7):701–707. doi: 10.1016/j.scispo.2022.04.007 PubMed PMID: 172043316.

- 75. Silvestre JC. Efeitos da suplementação crônica com β-alanina sobre o desempenho físico e a recuperação em praticantes de Crossfit[®]: um estudo clínico randomizado e duplo cego [master's dissertation]. Universidade Federal de São Paulo. 2023.
- 76. Fernández-Lázaro D, Seco-Calvo J, Pascual-Fernández J, et al. 6-week supplementation with *Tribulus terrestris* L. to trained Male CrossFit[®] athletes on muscle, inflammation, and antioxidant biomarkers: a randomized, single-blind, placebo-controlled trial. Int J Environ Res Public Health. 2022;19(23):16158. doi: 10.3390/ijerph192316158 PMID: 36498228; PMCID: PMC9736311.
- 77. Maroufi K, Razavi R, Gaeini AA, et al. The effects of acute consumption of carbohydrateprotein supplement in varied ratios on CrossFit athletes' performance in two CrossFit exercises: a randomized cross-over trial. J Sports Med Phys Fitness. 2021;61(10):1362–1368. doi: 10.23736/S0022-4707.20.11774-2 Epub 2020 Dec 14. PMID: 33314886.
- 78. Fernández-Lázaro D, Mielgo-Ayuso J, Del Valle Soto M, et al. The effects of 6 weeks of *Tribulus terrestris* L. Supplementation on body composition, hormonal response, perceived exertion, and CrossFit® performance: a randomized, single-blind, placebo-controlled study. Nutrients. 2021;13(11):3969. doi: 10.3390/nu13113969 PMID: 34836225; PMCID: PMC8623187.
- 79. Toledo LP, Vieira JG, Dias MR. Acute effect of sodium bicarbonate supplementation on the performance during CrossFit[®] training. Motriz Revista de Educacao Fis. 2020;26(4). doi: 10. 1590/S1980-6574202000040075
- Stein JA, Ramirez M, Heinrich KM. Acute caffeine supplementation does not improve performance in trained CrossFit[®] athletes. Sports (Basel). 2020;8(4):54. doi: 10.3390/sports8040054 PMID: 32340121; PMCID: PMC7240695.
- Moro T, Badiali F, Fabbri I, et al. Betaine supplementation does not improve muscle hypertrophy or strength Following 6 weeks of cross-fit training. Nutrients. 2020;12(6):1688. doi: 10. 3390/nu12061688 PMID: 32516959; PMCID: PMC7352895.
- Lima Ricordi M, da Silva Rodrigues I, Barbosa de Oliveira V, et al. Acute nitrate supplementation does not improve crossfit's physical performance, a double-blind cross-over study. Revista Brasileira de Nutrição E Esportiva. 2020;14(89):551–559. PubMed PMID: 152050795.
- Garnacho-Castaño MV, Palau-Salvà G, Serra-Payá N, et al. Understanding the effects of beetroot juice intake on CrossFit performance by assessing hormonal, metabolic and mechanical response: a randomized, double-blind, crossover design. J Int Soc Sports Nutr. 2020;17(1):56. doi: 10.1186/s12970-020-00388-z PMID: 33187518; PMCID: PMC7666517.
- 84. Fogaça LJ, Santos SL, Soares RC, et al. Effect of caffeine supplementation on exercise performance, power, markers of muscle damage, and perceived exertion in trained CrossFit men: a randomized, double-blind, placebo-controlled crossover trial. J Sports Med Phys Fitness. 2020;60(2):181–188. doi: 10.23736/S0022-4707.19.10043-6 Epub 2019 Oct 25. PMID: 31665873.
- Durkalec-Michalski K, Zawieja EE, Podgórski T, et al. The effect of chronic progressive-dose sodium bicarbonate ingestion on CrossFit-like performance: a double-blind, randomized cross-over trial. PLOS ONE. 2018;13(5):e0197480. doi: 10.1371/journal.pone.0197480 PMID: 29771966; PMCID: PMC5957406.
- Rountree JA, Krings BM, Peterson TJ, et al. Efficacy of carbohydrate ingestion on CrossFit exercise performance. Sports (Basel). 2017;5(3):61. doi: 10.3390/sports5030061 PMID: 29910421; PMCID: PMC5968949.
- Howarth TP, Bozanich R, Bridge M, et al. AD LIBITUM CONSUMPTION of SPORT DRINK DOES NOT IMPROVE PERFORMANCE in CROSSFIT ATHLETES. J Australian Strength Cond. 2017;25 (6):30–. PubMed PMID: 135834915.
- Kramer SJ, Baur DA, Spicer MT, et al. The effect of six days of dietary nitrate supplementation on performance in trained CrossFit athletes. J Int Soc Sports Nutr. 2016;13(1):39. doi: 10.1186/ s12970-016-0150-y PMID: 27822169; PMCID: PMC5093951.
- Outlaw JJ, Wilborn CD, Smith-Ryan AE, et al. Effects of a pre-and post-workout protein-carbohydrate supplement in trained CrossFit individuals. Springerplus. 2014;3 (1):369. doi: 10.1186/2193-1801-3-369 PubMed PMID: 25110627; PubMed Central PMCID: PMC4125607.

- 36 🕒 D. V. MARTINHO ET AL.
 - Jacobs PL. The acute effects of a commercial pre workout product, wodFuel[®], on performance of a CrossFit exercise series, the Cindy. J Int Soc Sports Nutr. 2014;11(sup1):1–2. doi: 10.1186/ 1550-2783-11-S1-P21 PubMed PMID: 156079320.
 - Heydenreich J, Kayser B, Schutz Y, et al. Total energy expenditure, energy intake, and body composition in endurance athletes across the training season: a systematic review. Sports Med Open. 2017;3(1):8. doi: 10.1186/s40798-017-0076-1 Epub 2017 Feb 4. PMID: 28161872; PMCID: PMC5292109.
 - 92. Magkos F, Yannakoulia M. Methodology of dietary assessment in athletes: concepts and pitfalls. Curr Opin Clin Nutr Metab Care. 2003;6(5):539–549. doi: 10.1097/00075197-200309000-00007 PMID: 12913671.
 - Capling L, Beck KL, Gifford JA, et al. Validity of dietary assessment in athletes: a systematic review. Nutrients. 2017;9(12):1313. doi: 10.3390/nu9121313 PMID: 29207495; PMCID: PMC5748763.
 - Burke LM, Hawley JA, Wong SH, et al. Carbohydrates for training and competition. J Sports Sci. 2011;29(Suppl 1):S17–27. doi: 10.1080/02640414.2011.585473 Epub 2011 Jun 9. PMID: 21660838.
 - Wardenaar F, Brinkmans N, Ceelen I, et al. Macronutrient intakes in 553 Dutch elite and sub-elite endurance, team, and strength athletes: does intake differ between sport disciplines? Nutrients. 2017;9(2):119. doi: 10.3390/nu9020119 PMID: 28208581; PMCID: PMC5331550.
 - CROSSFIT®. https://www.sportsdietitians.com.au/factsheets/food-for-your-sport/crossfit/ consulted2024
 - 97. Bytomski JR. Fueling for performance. Sports Health. 2018;10(1):47–53. doi: 10.1177/ 1941738117743913 Epub 2017 Nov 27. PMID: 29173121; PMCID: PMC5753973.
 - Timón R, Olcina G, Camacho-Cardeñosa M, et al. 48-hour recovery of biochemical parameters and physical performance after two modalities of CrossFit workouts. Biol Sport. 2019;36 (3):283–289. doi: 10.5114/biolsport.2019.85458 Epub 2019 Jul 31. PMID: 31624423; PMCID: PMC6786329.
 - Stellingwerff T, Boit MK, Res PT. International association of athletics federations. Nutritional strategies to optimize training and racing in middle-distance athletes. J Sports Sci. 2007;25 (Suppl 1):S17–28. doi: 10.1080/02640410701607213 Erratum in: J Sports Sci. 2009 Apr;27(6):667. PMID: 18049980.
- Stellingwerff T, Morton JP, Burke LM. A framework for periodized nutrition for athletics. Int J Sport Nutr Exerc Metab. 2019;29(2):141–151. doi: 10.1123/ijsnem.2018-0305 Epub 2019 Apr 3. PMID: 30632439.
- 101. Payne J, Welshans M, Seeland S, et al. Nutritional considerations for the power Athlete. Curr Sports Med Rep. 2022;21(7):224–231. doi: 10.1249/JSR.00000000000975 PMID: 35801723.
- 102. Jäger R, Kerksick CM, Campbell BI, et al. International society of sports nutrition position stand: protein and exercise. J Int Soc Sports Nutr. 2017;14(1):20. doi: 10.1186/s12970-017-0177-8 PMID: 28642676; PMCID: PMC5477153.
- 103. Morton RW, Murphy KT, McKellar SR, et al. A systematic review, meta-analysis and meta-regression of the effect of protein supplementation on resistance training-induced gains in muscle mass and strength in healthy adults. Br J Sports Med. 2018;52(6):376–384. doi: 10.1136/bjsports-2017-097608 Epub 2017 Jul 11. Erratum in: Br J Sports Med. 2020 Oct;54(19):e7.PMID: 28698222; PMCID: PMC5867436.
- 104. Carreker JD, Grosicki GJ. Physiological predictors of performance on the CrossFit "murph" challenge. Sports (Basel). 2020;8(7):92. doi: 10.3390/sports8070092 PMID: 32605265; PMCID: PMC7404702.
- 105. Mangine GT, Tankersley JE, McDougle JM, et al. Predictors of CrossFit open performance. Sports (Basel). 2020;8(7):102. doi: 10.3390/sports8070102 PMID: 32698335; PMCID: PMC7404807.
- 106. Mangine GT, McDougle JM, Feito Y. Relationships between body composition and performance in the High-intensity functional training workout "fran" are modulated by competition

class and percentile rank. Front Physiol. 2022;13:893771. doi: 10.3389/fphys.2022.893771 PMID: 35721570; PMCID: PMC9197730.

- 107. Hall M, Trojian TH. Creatine supplementation. Curr Sports Med Rep. 2013;12(4):240–244. doi: 10.1249/JSR.0b013e31829cdff2 PMID: 23851411.
- 108. Kreider RB, Kalman DS, Antonio J, et al. International society of sports nutrition position stand: safety and efficacy of creatine supplementation in exercise, sport, and medicine. J Int Soc Sports Nutr. 2017;14(1):18. doi: 10.1186/s12970-017-0173-z PMID: 28615996; PMCID: PMC5469049.
- 109. Wax B, Kerksick CM, Jagim AR, et al. Creatine for exercise and sports performance, with recovery considerations for healthy populations. Nutrients. 2021;13(6):1915. doi: 10.3390/ nu13061915 PMID: 34199588; PMCID: PMC8228369.
- 110. Devries MC, Phillips SM. Supplemental protein in support of muscle mass and health: advantage whey. J Food Sci. 2015;80(Suppl1):A8–A15. doi: 10.1111/1750-3841.12802 PMID: 25757896.
- 111. Zare R, Devrim-Lanpir A, Guazzotti S, et al. Effect of soy protein supplementation on muscle adaptations, metabolic and antioxidant status, hormonal response, and exercise performance of active individuals and athletes: a systematic review of Randomised controlled trials. Sports Med. 2023;53(12):2417–2446. doi: 10.1007/s40279-023-01899-w Epub 2023 Aug 21. PMID: 37603200; PMCID: PMC10687132.
- 112. Tipton KD, Wolfe RR. Protein and amino acids for athletes. J Sports Sci. 2004;22(1):65–79. doi: 10.1080/0264041031000140554 PMID: 14971434.
- 113. Wakolbinger-Habel R, Muschitz C. Comment on Kaufman et al. Popular dietary trends' impact on athletic performance: a critical analysis review. Nutrients. 2023;15(22):4709. doi: 10.3390/ nu15224709 PMID: 38004103; PMCID: PMC10675511.
- 114. Volek JS, Freidenreich DJ, Saenz C, et al. Metabolic characteristics of keto-adapted ultra-endurance runners. Metabolism. 2016;65(3):100–110. doi: 10.1016/j.metabol.2015.10. 028 Epub 2015 Nov 2. PMID: 26892521.
- 115. Burke LM. Ketogenic low-CHO, high-fat diet: the future of elite endurance sport? J Physiol. 2021;599(3):819–843. doi: 10.1113/JP278928 Epub 2020 Jun 10. PMID: 32358802; PMCID: PMC7891323.
- 116. Leverve X, Batandier C, Fontaine E. Choosing the right substrate. Novartis Found Symp. 2007;280:108–121; discussion 121–7, 160–4. PMID: 17380791.
- 117. Burke LM, Whitfield J, Heikura IA, et al. Adaptation to a low carbohydrate high fat diet is rapid but impairs endurance exercise metabolism and performance despite enhanced glycogen availability. J Physiol. 2021;599(3):771–790. doi: 10.1113/JP280221 Epub 2020 Aug 19. PMID: 32697366; PMCID: PMC7891450.
- 118. Joshi S, Shi R, Patel J. Risks of the ketogenic diet in CKD the con part. Clin Kidney J. 2023;17 (1):sfad274. doi: 10.1093/ckj/sfad274 PMID: 38186877; PMCID: PMC10768778.
- 119. Williams C, Rollo I. Carbohydrate nutrition and team sport performance. Sports Med. 2015;45 (Suppl 1):S13–22. doi: 10.1007/s40279-015-0399-3 PMID: 26553494; PMCID: PMC4672015.
- 120. Rollo I, Williams C. Carbohydrate nutrition and skill performance in soccer. Sports Med. 2023;53(Suppl 1):7–14. doi: 10.1007/s40279-023-01876-3 Epub 2023 Jul 8. PMID: 37421586; PMCID: PMC10721660.
- 121. Russell M, Kingsley M. The efficacy of acute nutritional interventions on soccer skill performance. Sports Med. 2014;44(7):957–970. doi: 10.1007/s40279-014-0184-8 PMID: 24728928.
- 122. Hulton AT, Malone JJ, Clarke ND, et al. Energy requirements and nutritional strategies for male soccer players: a review and suggestions for practice. Nutrients. 2022;14(3):657. doi: 10.3390/ nu14030657 PMID: 35277016; PMCID: PMC8838370.
- 123. Jeukendrup AE. Training the gut for athletes. Sports Med. 2017;47(Suppl 1):101–110. doi: 10. 1007/s40279-017-0690-6 PMID: 28332114; PMCID: PMC5371619.
- 124. King A, Helms E, Zinn C, et al. The ergogenic effects of acute carbohydrate feeding on resistance exercise performance: a systematic review and meta-analysis. Sports Med.

38 🕒 D. V. MARTINHO ET AL.

2022;52(11):2691–2712. doi: 10.1007/s40279-022-01716-w Epub 2022 Jul 9. PMID: 35809162; PMCID: PMC9584980.

- 125. Grgic J. Effects of caffeine on resistance exercise: a review of recent research. Sports Med. 2021;51(11):2281–2298. doi: 10.1007/s40279-021-01521-x Epub 2021 Jul 22. PMID: 34291426.
- 126. Southward K, Rutherfurd-Markwick KJ, Ali A. The effect of acute caffeine ingestion on endurance performance: a systematic review and meta-analysis. Sports Med. 2018;48 (8):1913–1928. doi: 10.1007/s40279-018-0939-8 Erratum in: Sports Med. 2018 Oct;48(10):2425–2441. PMID: 29876876.
- 127. Grgic J, Trexler ET, Lazinica B, et al. Effects of caffeine intake on muscle strength and power: a systematic review and meta-analysis. J Int Soc Sports Nutr. 2018;15:11. doi: 10.1186/s12970-018-0216-0 PMID: 29527137; PMCID: PMC5839013.
- 128. Grgic J, Mikulic P, Schoenfeld BJ, et al. The influence of caffeine supplementation on resistance exercise: a review. Sports Med. 2019;49(1):17–30. doi: 10.1007/s40279-018-0997-y PMID: 30298476.
- 129. Graham TE. Caffeine and exercise: metabolism, endurance and performance. Sports Med. 2001;31(11):785–807. doi: 10.2165/00007256-200131110-00002 PMID: 11583104.
- 130. Saunders B, de Oliveira LF, da Silva RP, et al. Placebo in sports nutrition: a proof-of-principle study involving caffeine supplementation. Scand J Med Sci Sports. 2017;27(11):1240–1247. doi: 10.1111/sms.12793 Epub 2016 Nov 23. PMID: 27882605.
- 131. Astorino TA, Terzi MN, Roberson DW, et al. Effect of two doses of caffeine on muscular function during isokinetic exercise. Med Sci Sports Exerc. 2010;42(12):2205–2210. doi: 10. 1249/MSS.0b013e3181e3a11d PMID: 20421833.
- 132. Warren GL, Park ND, Maresca RD, et al. Effect of caffeine ingestion on muscular strength and endurance: a meta-analysis. Med Sci Sports Exerc. 2010;42(7):1375–1387. doi: 10.1249/MSS. 0b013e3181cabbd8 PMID: 20019636.
- 133. Polito M, Souza D, Casonatto J, et al. Acute effect of caffeine consumption on isotonic muscular strength and endurance: a systematic review and meta-analysis. Sci Sports. 2016;31(3):119–128.
- 134. Pickering C, Grgic J. Caffeine and exercise: what next? Sports Med. 2019;49(7):1007–1030. doi: 10.1007/s40279-019-01101-0 PMID: 30977054; PMCID: PMC6548757.
- 135. Mitchell L, Wilson L, Duthie G, et al. Methods to assess energy expenditure of resistance exercise: a systematic scoping review. Sports Med. 2024;54(9):2357–2372. doi: 10.1007/s40279-024-02047-8 Epub 2024 Jun 19. PMID: 38896201; PMCID: PMC11393209.
- 136. Martinho DV, Rebelo A, Gouveia ÉR, et al. The physical demands and physiological responses to CrossFit®: a scoping review with evidence gap map and meta-correlation. BMC Sports Sci Med Rehabil. 2024;16(1):196. doi: 10.1186/s13102-024-00986-3 PMID: 39300545; PMCID: PMC11414238.