

# Please cite the Published Version

Onambélé-Pearson, GL, Breen, L and Stewart, CE (2010) Influence of exercise intensity in older persons with unchanged habitual nutritional intake: skeletal muscle and endocrine adaptations. Age, 32 (2). pp. 139-153. ISSN 0161-9152

DOI: https://doi.org/10.1007/s11357-010-9141-0

Publisher: Springer Verlag

Version: Accepted Version

Downloaded from: https://e-space.mmu.ac.uk/97661/

Usage rights: O In Copyright

Additional Information: The original publication is available at www.springerlink.com

### Enquiries:

If you have questions about this document, contact openresearch@mmu.ac.uk. Please include the URL of the record in e-space. If you believe that your, or a third party's rights have been compromised through this document please see our Take Down policy (available from https://www.mmu.ac.uk/library/using-the-library/policies-and-guidelines) Onambélé GL<sup>1</sup>, Breen L<sup>1,2</sup> and Stewart CE<sup>1,3</sup>

 <sup>1</sup> Department of Exercise & Sport Science, Manchester Metropolitan University, Alsager, UK.
 <sup>2</sup> Current address: School of Sport and Exercise Sciences, University of Birmingham, Birmingham, UK<sup>.</sup>
 <sup>3</sup> Current address: Institute for Biomedical Research into Human Movement and Health,

Manchester Metropolitan University, John Dalton Building, Manchester, UK.

Address for correspondence: Dr Gladys Onambélé-Pearson, Department of Exercise & Sport Sciences, Manchester Metropolitan University, Hassall Road, Alsager, ST7 2HL, UK. Email: <u>g.pearson@mmu.ac.uk</u>; Tel: +44 (0) 161 247 5594; Fax: +44 (0) 161 247 6375.

Short title: Exercise intensity for the elderly

#### <u>Abstract</u>

**Background:** Long-term adherence to training programs is difficult to attain. Yet the benefits of exercise to general health and well-being are undeniable. Any measure to demonstrate the minimum required exercise for maximal benefit to a person is a promising avenue towards increasing the uptake and adherence to physical activity for the general public. Purpose: To compare the effects of two different intensities of resistance training in healthy older adults. We hypothesize that compared to high intensity resistance exercise, relatively low training intensity could also improve in vivo markers of healthy physiologic and endocrine functions in previously sedentary older individuals. Methods and Results: Thirty (out of a possible 34 recruited) older adults were randomly assigned to low (LowR, i.e. ~40% one Repetition Maximum (1-RM) versus high resistance training (HighR, i.e. ~80% 1RM) for 12 weeks. Neither intervention significantly impacted upon body composition markers including: Body mass index (BMI), waist:hip ratio and bioelectric impedance. Muscle strength data showed an advantage for the HighR protocol with  $51 \pm$ 4% and 22.4  $\pm$  10.2% (P<0.05) improvements in 1-RM strength and bilateral knee extension torque respectively, compared with  $17 \pm 1\%$  and  $10.3 \pm 4.7\%$  (P<0.05) increments in 1-RM strength and bilateral torque in the LowR group. Unilateral torque did not change significantly in either group. Muscle thickness data also showed a significantly greater benefit of the HighR protocol ( $5.8 \pm 2.6\%$ increase) compared with the LowR protocol (no change). Functional ability tests including: Get-Upand-Go (GUG), Standing from lying and the six-minute walk, showed changes of  $-11.6 \pm 4.8\%$ , -15.6% and  $8.5 \pm 1.7\%$  (P<0.05) respectively in HighR compared with only one significant improvement in the LowR, namely a  $-10.8 \pm 3\%$  (P<0.05) improvement in the GUG test. Overnight fasting serum levels of IGFBP-3 increased, NPY decreased and TNF- $\alpha$  decreased significantly in the LowR group. Serum levels of glucose increased and NPY decreased significantly in HighR. Circulating levels of insulin, IL-6 and IGF-I did not change with either intervention. Conclusion: In vivo physiologic changes show functional advantages for older persons carrying out high resistance training. At the endocrine level, such an advantage is not clear. In fact, in terms of changes in sera levels of fasting glucose, IGFBP-3 and TNF- $\alpha$ , there appears to be an advantage to carrying out the lower intensity exercises for the aged populations where endocrine adaptations are key.

Key words: Ageing, Endocrinology, Cytokines, Physiology, Resistance Exercise Intensity.

# **Introduction**

Increased dependence in old age occurs through a number of circumstances, with decreased mobility being one of the most acutely debilitating (Doherty, 2003). In hand with decreased mobility, come notable reductions in skeletal muscle mass (sarcopenia), which occurs at ~5% per decade after the 4<sup>th</sup> decade (Roubenoff et al., 2000), and increases by a further ~1% every year after the 65<sup>th</sup> birthday (Greenlund & Nair, 2003; Klitgaard et al., 1990; Lexell, Taylor, & Sjostrom, 1988; Narici, Maganaris, Reeves, & Capodaglio, 2003; Young, Stokes, & Crowe, 1985). Sarcopenia is also associated with asthenia or decreased muscle strength whereby in the knee extensors for instance, strength losses are in the order of 20-50% between the ages of 25 to 73 years (Poulin, Vandervoort, Paterson, Kramer, & Cunningham, 1992; Vandervoort, Kramer, & Wharram, 1990; Young et al., 1985).

That lifelong (Pearson et al., 2002) and/or newly adopted (Onambele et al., 2008) high levels of physical activity slow down and/or reverse age related declines in strength is undoubtedly demonstrable. What is clear also, is that whilst endurance (Klitgaard et al., 1990) and aerobic (Sipila, Elorinne, Alen, Suominen, & Kovanen, 1997) work have their cardiovascular benefits, resistance work more specifically, can be associated with improved functional ability in daily tasks requiring strength (Coggan et al., 1992; Tracy et al., 1999). In their seminal paper, Frontera and co-workers (Frontera, Meredith, O'Reilly, Knuttgen, & Evans, 1988) used high intensity strength training in healthy older men aged 60-72. The striking increments in strength (~150%) and mid-thigh muscle mass (11.4%) after 12-weeks of resistance training, have since established the training programme used as 'normal practice'. This protocol involved 3 sets of 8 repetitions (at 80% of the one repetition maximum (1 RM)), 3 times a week, for each muscle group being trained. In the same vein, in recent work, it was concluded that 70-90% 1RM training in healthy older individuals was the most effective protocol for obtaining training-induced increments in muscle mass, with strength improvements being comparable to those seen in young individuals (Chin, van Uffelen, Riphagen, & van Mechelen, 2008; Fielding, 1995).

Another important aspect of ageing, which may in fact interfere with responsiveness to exercise, is the background of increased levels of inflammatory cytokines, a phenomenon referred to as 'inflamed ageing' (Butcher & Lord, 2004), coupled with worsened metabolic profile, particularly in terms of hormones associated with both appetite and glucose metabolism (Adamczak, Rzepka, Chudek, & Wiecek, 2005; Baranowska et al., 2006; Schutte et al., 2007). In a context where a) muscle hypertrophy can be achieved through either the decreased breakdown or the increased production of proteins (Short et al., 2004), b) ageing-related sarcopaenia is related to decreased myosin heavy chain (crucial in muscle contractile function) synthesis rate, and c) hypoactivity is linked to a greater degree with decreased synthesis than any changes in breakdown activity (Rennie et al., 2009), the synthesis rate of myosin heavy chain has been shown to correlate to circulating levels of IGF-I, DHEAS, and testosterone (Balagopal et al., 1997). Thus, whilst a causal relationship has not necessarily been shown, and in fact increased levels of GH and/or IGF-I are beneficial in deficient but not healthy individuals in terms of hypertophic response (for a review read (Velloso, 2008), these compounds may yet offer an insight into protein synthesis (and thus muscle anabolism) through implied modulated ability to synthesise myosin heavy chain. Together, the above phenomena suggest that the strategy to prescribing exercise in an older population ought to be manipulated in order to also take the altered hormonal environment into account, thus potentially maximising all of the potential positive effects of exercise intervention.

Given the worldwide age demographics, coupled with the fact that older people have the greatest barriers (mostly in terms of accessibility and social context) to participating in exercise, simple advice on the minimum requirement for maximal benefits are likely to be the key to empowering older persons in terms of their personal lifestyle strategies. There are currently no systematic studies geared towards minimising the level of exercise intensity whilst maximising muscle strength and size as well as improving the endocrine profile of older adults.

The current study therefore aimed to determine whether, in an otherwise healthy older population, a) high resistance activity would have positive muscle volume, functional and endocrine effects in older persons and b) low resistance activity could in fact elicit similar degrees of improvement (as high resistance activity) in hormonal, metabolic and functional parameters associated with physical wellbeing. Based on the principle that a previously reduced physical activity background entails a relatively greater potential for positive response to increased activity particularly in the older individual (Mroszczyk-McDonald, Savage, & Ades, 2007), we hypothesized that compared to high intensity resistance exercise, relatively low training intensity could also improve markers of healthy physiologic and endocrine functions in previously sedentary older individuals. In other words, in view of the fact that the literature has already reported a link with several endorine factors and resistance exercise (for reviews see (Crewther, Keogh, Cronin, & Cook, 2006) and (Velloso, 2008)), the purpose of the current study was in fact to a) determine the effectiveness of relatively low resistance exercise and then b) establish any link of changes in endocrine factors with exercise intensity in older people.

# **Methods**

#### 1-Participants

Thirty-nine older adults volunteered to participate in the current study having responded to advertisements posted locally. All applicants gave written informed consent to take part in the study though five of these prospective volunteers were excluded as they had a known history of cardiovascular, neurological, inflammatory or myopathic disease. Each participant's general practitioner gave medical consent for the exercise regime. Thus the current study participants were healthy, community dwelling, and habitually active individuals, with no recent history of structured resistance training. The local Human Ethics Committee approved all experimental procedures.

Of the 34 healthy older adults who started the study, 30 completed the 12-week exercise intervention. Participants were randomly assigned to either a low (LowR), or a high (HighR) resistance-training group: The gender repartition, as well as age and physical characteristics of the completing participants are presented in table 1.

# $\rightarrow$ [Table 1 near Here]

## 2-Muscle strength measurements

**2a. One Repeated maximum measurement**- During a familiarization session no more than 7-days prior to the 12 week intervention, participants' 1 Repetition Maximum (1RM) was determined for all exercises employed in the training programme. Participants first performed a standardized warmup on the leg press ( $6 \times 50\%$  perceived 1RM;  $4 \times 70\%$  perceived 1RM with three min recovery). Note here that we used 'perceived sub-maximal efforts' so as to not induce fatigue in the participants and/or injury by asking them to perform a maximal contraction on a 'cold' muscle. After warming up, the load was set at 90 % of the initially estimated 1RM, and increased after each successful lift by 5kg until failure. Each participant was given six lifting attempts in order to achieve their 1RM and a maximum of two attempts to lift the weight, once it had been established. The greatest amount of weight lifted successfully was recorded to determine the training load. Between successive attempts, 3 min rest periods were allowed. A repetition was valid if the participant used correct form and was able to complete the entire lift in a controlled manner without assistance. Participants 1RM for each exercise was reviewed every 2-weeks during training, if 1RM had increased the training load was adjusted accordingly. Additionally if any participants felt that in-between 1RM assessments the training load was not providing adequate resistance, the load was increased so they were always lifting at the desired percentage of their maximum. Participants were familiarized with the resistance exercise training protocol on a subsequent visit to the laboratory.

**2b.** Isometric Knee extensors muscle strength measurements- Participants were familiarized with the experimental procedures on a separate occasion no more than 7 days prior to the baseline test sessions. In one single testing session quadriceps isometric strength measures were taken on the right leg and isokinetic measures on the left leg, using a Cybex Dynamometer (Cybex Norm, Cybex International Inc., NY, USA). The centre of rotation of the lever arm of the dynamometer was aligned with the axis of rotation of the knee. Participants were positioned with the hip joint at 85° (supine = 0°). In order to minimise any extraneous movement of the hip joint or the trunk, participants were strapped over the shoulders, pelvis and thighs. Settings of chair height and positioning relative to the dynamometer were adjusted individually with all settings recorded and replicated at the post-intervention testing phase. Gravity corrections were then made following the manufacturers' own procedure, having adjusted the attachment of the lever arm cuff relative to the length of the participant's shank. Previous work from within our laboratory has utilized similar methods of measurement, finding them to be both valid and reliable (Pearson & Onambele, 2005, 2006).

**2b-I- Maximal Unilateral Isometric Torque:** Maximal isometric knee-extension torque was measured with the knee at 70° angle (full knee extension = 0°) on the right leg of all participants. After a series of warm-up trials consisting of 10 isokinetic contractions at  $60^{\circ}$ s<sup>-1</sup> at 50-75% maximal effort, participants were instructed to rapidly exert maximal isometric force against the Cybex lever arm over a 3-4 s period. Participants were given both verbal and visual encouragement/feedback throughout their effort. Joint torque data were displayed on the screen of a computer (Macintosh G4; Apple Computer, Cupertino, CA), which was interfaced with an A/D system (Acknowledge, Biopac Systems, Santa Barbara, CA) with a sample frequency of 500 Hz. Isometric contractions were held for ~2 s at the plateau, with a 90 s rest period between

contractions. Peak torque was averaged over a 500 ms period at the plateau phase. The mean peak torque (MVCext) of three extensions was used as the measure of strength in each participant.

**2b-II- Maximal Bilateral Isometric Torque**- Maximal bilateral isometric knee-extension torque was measured with both knees at 70° angle (full knee extension = 0°) pushing against a customized lever arm. Similar precautions to record time of day were taken. After a series of warm-up trials consisting of 3 isokinetic contractions at  $60^{\circ}$ s-<sup>1</sup> at 50% Maximal effort, participants were instructed to rapidly exert maximal isometric force against the Cybex lever arm over a 3-4 s period. Participants were given both verbal and visual encouragement/feedback throughout their effort. Joint torque data were acquired and processed as during the unilateral efforts, with the only difference that here the best of the three efforts was used as the measure of strength for each participant.

# 3-Mid-thigh muscle thickness

Real-time B mode ultrasonography with a 7.5 MHz linear-array probe (AU5, Esaote, Genoa, Italy) was used to study mid-thigh muscle thickness in the area of the Vastus Intermedius (VI) and the Vastus Lateralis (VL) muscles. Muscle thickness was taken at rest with the knee at 70° angle. Scans were acquired in the mid-sagittal plane, at approximately 50% length of the VL muscle as measured from the origin at the linea aspera & lateral femur to insertion at the tibial tuberosity via patella tendon on the anterior surface. Medio-lateral width of the VL was determined over the skin surface and the position of one half of the width was used as the measurement site. The ultrasound probe was coated with water-soluble transmission gel to provide acoustic contact and was held in place without depressing the dermal surface by the assessor. Three separate recordings of mid-thigh muscle thickness were made with 90 s rests between each recording. Ultrasound images were acquired using a digital recorder and frames exported to capture software (iMovie HD6, Apple computer Inc, USA). The VL thickness was measured as the distance from the top of the peripheral muscle aponeurosis to the deep aponeurosis. VI thickness was from the deep aponeuroses to the surface of the femur. The three points of interest were measured at 3 standardised points on each ultrasound frame to obtain an average tissue thickness using ImageJ analysis software (ImageJ 1.37, Maryland, USA). Total muscle thickness was computed as the sum of VI+VL muscle thickness measured.

# 4-Body composition

**4a. Waist:Hip ratio-** A measuring tape was used to measure the circumference of the hips at the widest part of the buttocks, and the waist at the smaller circumference of the individual's natural waist (just above the navel). The waist:hip ratio was used as the individual's score of central adiposity.

**4b. Bioelectrical impedance analysis** (BIA) (BODYSTAT, Isle of Man, British Isles) was used to estimate body composition based on the difference in electric conductive properties of various tissues. The BODYSTAT applied 500 micro Amps at a single frequency of 50 kHz through self adhesive electrodes placed on the right hand and foot, of a participant lying flat on their back with their arms away from the trunk, thighs not touching and ankles at least 20cm apart. Although BIA has acknowledged limitations when applied to non-standard or elderly populations due to the built in equation utilised to calculate body composition, it is important to state that the BIA in this study was used to assess within participant adaptations as a consequence of the intervention, and as such carries more validity. What is more, to lend greater external validity to data from this instrument, raw values were corrected for the inherent ~15% overestimation in body fat content (when comparing pilot participants BIA versus DEXA outputs – others have drawn similar conclusions regarding the differences in body fat content values depending on the methodology used (Jorgensen et al., 1996); (Lintsi, Kaarma, & Kull, 2004) and have tended to consider DEXA readings as gold standard since it yields values similar to those obtained from the hydrostatic weighing method (Prior et al., 1997).

**4c. Body mass index** (BMI) was defined as the individual's body mass (Kg) divided by the square of their height (m<sup>2</sup>).

# 5- Functional abilities measures

The tests of functional ability were all performed on one day at the outset and completion of the study and were similar to a battery of functional tests performed by participants in previous research (Chandler, Duncan, Kochersberger, & Studenski, 1998; Skelton, Young, Greig, & Malbut, 1995).

**5a. Get-Up-and-Go-** Three cones were placed 1m apart on the floor in front of a rigid chair of adjustable height. The participant's knee-to-floor height (i.e. the distance from the knee joint axis to the floor) was recorded prior to testing to determine the appropriate chair height for each test. Once knee height was determined the chair height was set at 100, 80 and 60% of each individual knee-to-floor height for the tests. The test started with the participant seated on the chair (at the appropriate height), with feet flat on the floor and arms folded across the chest. They were then asked to rise unaided as quickly as possible, walk around the furthest cone (3m away) and back to the initial seated position on the chair. The elapsed time between 'chair rise' to 'sitting back down' was recorded. The quickest of three trials was used as the participant's score.

**5b. Standing from Lying (SFL)** - Participants were asked to lie flat on the floor on a gym mat and on their preferred side, with their arm on the floor outstretched and their head flat on the outstretched arm. Participants were then instructed to rise as quickly as possible using their preferred technique. The time elapsed between the instruction to 'Go' and the participant standing upright and steadily with both feet firmly on the ground, was recorded. The fastest of 3 trials was utilised as their score.

**5c. Six-minute walk-** A 10m course was set with cones 1m apart in a straight line. Participants were instructed to walk around this course using their fastest, non-running, walking pace, with the aim of completing as many revolutions of the circuit as possible in 6 minutes. The score was calculated as the total distance covered in the allocated time.

# 6- Metabolic & Endocrine profiling

At the onset and end of the intervention and following an overnight fasting period, participants reported to the laboratory. A 21-gauge 1-inch ultra thin wall needle (Terumo medical corporation, New Jersey, USA) was inserted into the anticubital vein of the forearm. Using a vacutainer assembly and serum separator tubes (Monovette, Sarstedt, Numbrecht, Germany), 10mL blood samples were collected. Blood glucose was analysed immediately using a single drop of freshly sampled blood using the AccuChek Advantage System (Roche Diagnostics Ltd, Lewes, UK. Sensitivity of <10 mg/dL (i.e. minimum detectable concentration); Intra-assay variability of 2% (i.e. coefficient of variation). The remainder of the sample was centrifuged at 2-5°C for 5 minutes at 4000 rpm, with the supernatant being removed and stored in eppendorfs at -70° Celsius for later

analyses. Insulin (Biosource, Nivelles, Belgium. Sensitivity of 0.15 µlU/ml; Intra-assay variability of 4.2%). Insulin-like Growth Factor-I (IGF-I; Biocode-Hycel, Liege, Belgium. Sensitivity of 4.9 ng/ml; Intra-assay variability of 8.0%). Tumour Necrosis Factor-α (TNF-α; Diaclone, Besancon Cedex, France. Sensitivity <8 pg/ml; Intra-assay variability of 3.3%). NeuroPeptide Y (NPY; Phoenix Europe GmbH, Karlsruhe, Germany. Sensitivity of 0.13ng/ml; Intra-assay variability <5%). Insulin-like Growth Factor Binding Protein-3 (IGFBP-3; Biocode-Hycel, Liege, Belgium. Sensitivity of 10.5ng/ml; Intra-assay variability of 6.5%) and InterLeukin-6 (IL-6; Diaclone, Besancon Cedex, France. Sensitivity <0.8 pg/ml; Intra-assay variability of 3.3%) were analysed using standard enzyme-linked immuno-sorbent assay (ELISA) procedures.

# 7- Training programme

The training programme was 12-weeks in duration and consisted of one supervised gym-based class and two home-based sessions per week in LowR. In HighR the programme was for two supervised gym-based classes and one home-based session per week. All exercise sessions were 1 hr in duration.

Briefly, the supervised exercise classes consisted of a warm-up (stretching, aerobic and coordination work), resistance exercises (using therabands for all major muscle groups, Leg Press, Leg Extension, Calf Rotator and Glute conditioner (Technogym, Gambettola, Italy) as partly illustrated on figure 1 below, with a progression from 8-11 reps in 2-4 sets at 40 or 80% 1RM), and a cool-down (i.e. stretches, pilates, Tai Chi). The unsupervised home-based exercises were similar in design to the supervised classes with the exception that all the resistance work was carried out using therabands, and a 20-min brisk walk was also included. An exercise booklet illustrated, using photographic and/or cartoons, all the exercises in detail. Home-based exercise was not to be performed the day preceding or following the supervised class exercise.

 $\rightarrow$  [Figure 1 near Here]

## Statistical analyses

T-tests were carried out to compare data at baseline. Two-way factorial ANOVAs were carried out with group as one factor (2 levels: HighR vs. LowR) and phase as the second factor (2 levels: baseline vs. post-intervention) to determine any main effects of the group or interventions. An ANCOVA was run on the Get-Up-and-GO, IGFBP3 data sets since the LowR and HighR

populations differed in these parameters at the onset of the study. Data are expressed as mean  $\pm$  S.E.M (standard error of the mean) unless otherwise stated. Significance was set at P  $\leq$  0.05.

# <u>Results</u>

The two intervention groups did not significantly differ in age, habitual physical activity levels, height or weight at the onset of the current programme. Interestingly however, the HighR group showed a tendency towards a greater overall body mass increase ( $\sim$ 3%; from 76.2 ± 3.5 to 78.3 ± 4.2 Kg) after the 12-week intervention compared to the changes seen in the LowR group ( $\sim$ 1%; from 73.7 ± 2.50 to 74.5 ± 2.1 Kg). This effect however was not significant.

#### Body composition changes

No significant difference in BMI was observed in the two groups after the 12-week interventions. What is more, there was no change in waist:hip ratio in either the LowR population  $(0.90 \pm 0.02 \text{ to} 0.91 \pm 0.02)$  or the HighR population  $(0.88 \pm 0.03 \text{ to} 0.87 \pm 0.02)$ . Nonetheless, interestingly, LowR exhibited a marked trend for increased body-fat percentage of  $18.5 \pm 9.0\%$  (from a value of  $26.3 \pm 2.1\%$  to  $29.8 \pm 2.2\%$  post-intervention, P = N.S.), whereas such a trend for a rise in body fat percentage was attenuated in the HighR population (value of  $9.2 \pm 7.9\%$  i.e. from  $29.1 \pm 2.7\%$  to  $30.8 \pm 2.6\%$ , P = N.S.).

### Knee extensors muscle strength changes

<u>a) 1-RM</u> - Prior to training there was no significant difference in the mean 1RM strength measured by leg press, leg extension, calf rotation or gluteus conditioner between groups (P = N.S.). At 4, 8 and 12 weeks (i.e the post-intervention phase), the average strength increase for each exercise was significant for both intervention groups (P < 0.05). However the increase for LowR was significantly less pronounced in all exercises compared to HighR (see table 2). The average 12week increase in strength was  $17 \pm 1\%$  for LowR ( $8.9 \pm 1.7$ kg) and  $51 \pm 4\%$  for HighR ( $23.6 \pm 3.4$ kg).

# $\rightarrow$ [Table 2 near Here]

<u>b)</u> Isometric: Unilateral and Bilateral torque - Prior to training there was no significant difference in either the mean isometric unilateral or bilateral knee extensor strength (MVCext) between the two

intervention groups (P = N.S.). Table 2 shows the mean changes in MVCext after the 12-week interventions in each group. Briefly, trends towards increased unilateral isometric force were seen post interventions though these effects were not significant. The HighR group exhibited a non-significant change in MVCext of  $9.3 \pm 5.1\%$  ( $122.6 \pm 13.6$ Nm to  $133 \pm 13.6$ Nm, P = N.S.). LowR MVCext changed non-significantly by  $1.3 \pm 3\%$  ( $122.2 \pm 9.5$ Nm to  $122.8 \pm 9.3$ Nm, P = N.S.). However, the 12-week intervention resulted in a significant increase in bilateral torque in both HighR ( $22.4 \pm 10.2\%$ , P = 0.024) and LowR ( $10.3 \pm 4.7\%$ , P = 0.024) groups.

### Mid-thigh muscle thickness

Total muscle thickness (VL+VI) was significantly increased in the HighR participants by  $5.8 \pm 2.6\%$  ( $34.4 \pm 1.8$  mm to  $36.4 \pm 2.0$  mm; P = 0.038). LowR was without effect on total muscle thickness ( $32.6 \pm 2.0$  mm to  $34.6 \pm 2.2$  mm;  $6.8 \pm 5.4\%$  change, P = N.S.).

### Functional measures

<u>a- Get-Up-and-Go</u> - At baseline there was a significant difference in reaction time (as determined by the get-up-and-go (GUG) test) between the two groups, whereby LowR participants were taking significantly longer to complete the task (P < 0.04). Both groups completed the GUG test significantly quicker post-intervention. GUG time significantly improved by  $10.8 \pm 3\%$  (6.8 ± 0.4 sec to  $6.0 \pm 0.3$  sec) in the LowR population. Similarly, HighR also significantly improved by 11.6  $\pm 4.8\%$  (5.5 ± 0.3 sec to  $4.9 \pm 0.3$  sec). Even after accounting for the between groups differing baseline values through the ANCOVA test, the difference in response to treatments remained significant (P < 0.05).

<u>b-Standing from lying (SFL)</u> - Functional power determined by SFL time showed significant improvement in the HighR population  $(2.73 \pm 0.19 \text{ to } 2.36 \pm 0.2, \text{ i.e} -15.6\% \text{ change}, P = 0.025)$  whilst the LowR population in fact showed a trend towards increased SFL  $(3.69 \pm 0.33 \text{ to } 3.75 \pm 0.42, \text{ i.e} 5.2\% \text{ change}, P = N.S.)$ .

<u>c-Six minute walk</u> - The distance covered during the six-minute walk test was increased by  $8.5 \pm 1.7\%$  (25.7  $\pm 1.3$ m to 28  $\pm 1.7$ m) in the HighR participants (P < 0.003). The LowR participants showed no change in distance covered (24.2  $\pm 1$ m to 24.5  $\pm 0.9$ m;  $1.9 \pm 2.3\%$  change, P=N.S.).

# Metabolic & endocrine characteristics changes

Data of the endocrine changes (i.e. post-intervention values normalised for baseline values) are summarised in figure 2.

# $\rightarrow$ [Figure 2 near Here]

## Metabolic profile – Plasma levels of Glucose and Insulin

At baseline, there was no significant difference in the mean plasma glucose levels between intervention groups. The 12-week intervention resulted in a significant elevation in fasting plasma glucose of  $16.1 \pm 4.1\%$  (P = 0.001) in the HighR group ( $4.80 \pm 0.19$  mmol/L to  $5.51 \pm 0.08$  mmol/L). The LowR group exhibited a (non-significant) trend for lower post-intervention plasma glucose of  $-2.3 \pm 4.9\%$  ( $5.34 \pm 0.20$  mmol/L to  $5.11 \pm 0.19$  mmol/L).

Prior to training there was no significant difference in the mean plasma insulin levels between the two groups. The 12-week intervention resulted in no change in plasma insulin levels for both LowR  $(3 \pm 7.5\%$  change (i.e.  $10 \pm 1 \mu$ U/ml to  $10.4 \pm 1.4 \mu$ U/ml) and HighR ( $6 \pm 6.6\%$  change (i.e.  $8.06 \pm 0.69 \mu$ U/ml to  $8.51 \pm 0.92 \mu$ U/ml) groups.

# Control of energy homeostasis in relation to metabolic status- Plasma levels of NPY

Prior to training there was no significant difference in the mean plasma NPY levels between the two intervention groups. The 12-week interventions resulted in significant decreases in plasma NPY levels for both LowR (-13.7% change (i.e.  $23.4 \pm 1.66$  pg/ml to  $20.2 \pm 1.67$  pg/ml), P = 0.026) and HighR groups (-27.3% change (i.e.  $28.4 \pm 4.03$  pg/ml to  $20.7 \pm 3.53$  pg/ml), P = 0.044).

#### Plasma levels of IGF-I and IGFBP-3

Serum levels of IGF-I did not differ significantly between the two groups at baseline. Post intervention, in the LowR group, values changed from  $342.6 \pm 23.1$  ng/ml at baseline to  $313.8 \pm 20.8$  ng/ml post training, a -8.4% change with the intervention, which was not, however, significant. Similarly in the HighR group, values decreased from  $404.2 \pm 31.2$  ng/ml at baseline to  $383.5 \pm 19.0$  ng/ml post training, also a non-significant – 5.1% change in serum IGF-I levels with the intervention.

Serum levels of IGFBP-3 differed significantly between the two groups at baseline (P = 0.035). Post intervention, in the LowR group, values changed significantly from  $3827.2 \pm 214.9$  ng/ml at baseline to  $4317.8 \pm 282.0$  ng/ml post training, a 12.8% (P = 0.005) increase with the intervention. However in the HighR group, values did not change significantly with the intervention whilst decreasing from  $3050.9 \pm 259.4$  ng/ml at baseline to  $2587.8 \pm 121.1$  ng/ml post training, a -15.2% non-significant change. Even after accounting for the between groups differing baseline values through the ANCOVA test, the difference in response to treatments by group remained significant (P = 0.012).

# Cellular degradation/inflammation signal - Plasma levels of TNF-α and IL-6

Serum levels of TNF- $\alpha$  did not differ significantly between the two groups at baseline. Post intervention, values decreased in both groups. Thus, in the LowR group, values changed significantly from 35.2 ± 8.7 pg/ml at baseline to 29.6 ± 7.7 pg/ml post training, a -15.8% (P = 0.030) change with the intervention. In the HighR group, values did not change significantly (29.0 ± 18.6 pg/ml at baseline vs. 31.0 ± 20.9 pg/ml post training; an 11% change, P = N.S.).

Serum levels of IL-6 did not differ significantly between the two groups at baseline and were unaltered by training (LowR:  $3.00 \pm 0.58$  pg/ml at baseline to  $2.95 \pm 0.31$  pg/ml post training; HighR group:  $2.14 \pm 0.45$  pg/ml at baseline to  $2.56 \pm 0.71$  pg/ml post training).

#### **Discussion**

# Summary of findings in view of the study hypothesis

The current study sought to compare *in vivo* changes in skeletal muscle function, body composition and endocrine parameters in older people using two intensities of exercise, where one group of older persons performed resistance exercise once a week under supervision at 40% 1-RM using resistance machines, plus two unsupervised home-based training sessions using mainly therabands, and the second group performed resistance exercise twice a week under supervision at 80% 1-RM using resistance machines, and one unsupervised training session at home, using mainly therabands. We aimed to determine whether a) high resistance activity had positive muscle volume, functional and endocrine effects in older persons and b) low resistance activity could in fact elicit a similar

degree of improvement (as high resistance activity) in several factors (e.g. hormonal and functional) associated with physical wellbeing. We hypothesized that, owing to the impact of baseline function on responsiveness to training, compared to high intensity resistance exercise relatively low training intensity could also improve *in vivo* markers of healthy physiologic and endocrine functions in previously sedentary older individuals.

Our current findings have partially supported our hypotheses. Indeed, whilst none of the interventions significantly impacted upon body composition markers including, BMI, waist: hip ratio and bioelectric impedance, muscle strength data showed an advantage for the HighR protocol with  $51 \pm 4\%$  and  $22.4 \pm 10.2\%$  (P<0.05) improvements in 1-RM strength and bilateral knee extension torque respectively, compared with  $17 \pm 1\%$  and  $10.3 \pm 4.7\%$  1% (P<0.05) increments in the LowR group. Muscle thickness data also showed a greater benefit of a HighR protocol ( $5.8 \pm 2.6\%$  increase, P < 0.05) compared with the LowR protocol (no significant change). Functional ability tests including Get-Up-and-Go, Standing from lying and the six-minute walk all showed significant changes of  $-11.6 \pm 4.8\%$ , -15.6% and  $8.5 \pm 1.7\%$  (P<0.05) respectively in HighR. In comparison only the GUG test showed a significant decrease of  $-10.8 \pm 3\%$  (P<0.05) in LowR. Finally, overnight fasting serum levels of IGFBP-3, NPY and TNF- $\alpha$  changed significantly in LowR whereas only serum levels of glucose and NPY changed significantly in HighR groups. Circulating levels of IL-6, insulin and IGF-I did not alter significantly with either intervention.

### Muscle size and strength changes

Sixteen weeks of resistance exercise training has been shown to increase not only the strength but also type II muscle fiber size in old adults (Kosek, Kim, Petrella, Cross, & Bamman, 2006). Indeed, numerous studies have demonstrated that resistance training is an effective measure to counteract sarcopenia in both healthy and frail older adults (Carmeli, Reznick, Coleman, & Carmeli, 2000; Charette et al., 1991; Ferri et al., 2003; Frontera et al., 2000; Frontera et al., 2003; Frontera et al., 1988). It would appear that with an appropriate and effective exercise regimen older adults may expect gains of 5-10% muscle CSA, accompanied by increases of 20-100% or more in muscle strength, depending on the muscle group under investigation. Our data thus agree with previous studies of similar duration (Charette et al., 1991; Frontera et al., 1988; Hakkinen, Kraemer, Newton, & Alen, 2001; Lexell, 1995) in terms of the magnitude of muscle strength and size benefits to be had from high intensity resistance exercise. Studies other than the current one however, have not

systematically compared lower intensity resistance exercise with high intensity programmes to try to quantify physiologic and endocrine adaptations, making our data novel in the current exercise intervention literature, as far as we are aware.

In agreement also with our current data, the majority of previous studies investigating the effects of strength training on muscle strength and mass in older adults have observed greater increases in muscle strength than in muscle CSA. Similar to the events seen in younger adults, the discrepancies between gains in muscle strength and gains in muscle size have mainly been attributed to neural adaptations (Fiatarone et al., 1990). Findings indicate that fibre activation and recruitment are more rapidly modulated than muscle hypertrophy response. This is especially true of the low intensity-trained participants in the current study who exhibited significant strength increments post-intervention, without however, showing any muscle size increments.

### Molecular background of the observed in vivo muscle characteristics

Repeated bouts of resistance exercise increase protein synthesis (Hartman, Moore, & Phillips, 2006; Yarasheski, 2003), the metabolic determinant of gains in skeletal muscle mass (Kosek et al., 2006) in young and old adults, even if this increase in muscle protein synthesis with resistance exercise training in the old is lower than that seen in the young (Welle, Totterman, & Thornton, 1996). Our current findings suggest that there is no direct link between physical exercise and presence of chronically elevated, systemic IGF-I. In fact there were no discernable changes in the levels of IGF-I at end of the two training interventions. Interestingly, levels of IGFBP-3 increased significantly but only in the lower intensity-trained population, whilst the high intensity trained participants showed no change in the levels of this IGF-modulatory protein. Most IGF-I (~80%) circulates in a 150-kDa, high-affinity complex, which also contains IGFBP-3 and an acid-labile subunit. Analogous to the present findings, studies have found no change in IGF-I and IGFBP-3 concentrations following 8 (Bermon, Ferrari, Bernard, Altare, & Dolisi, 1999) and 12 weeks (Kraemer et al., 1999) of resistance training in a similarly aged population. In contrast, a significant increase in plasma IGF-I concentrations following an acute exercise bout was reported in one of these studies (Bermon et al., 1999). In endurance training studies, IGFBP-3 has been shown to be independent of IGF-I responses and potentially to have its own biological activity at the level of the cell (Jones & Clemmons, 1995). Eliakim et al. (Eliakim et al., 1997) showed that in rat, muscle IGF-I concentrations can increase with relatively short duration endurance training, despite lack of

change in muscle IGF-I mRNA or serum IGF-I. These authors proposed that the mechanisms through which muscle IGF-I concentrations post-training could alter with no concurrent changes in muscle IGF-I mRNA nor indeed serum IGF-I levels may either involve a) translational and/or posttranslational mediated events, b) exercise-induced increase in total blood volume, or c) modulated characteristics of the IGF binding proteins in the muscle tissue. Then again, a lack of change in serum IGF-I with training may suggest that IGF-I in the circulation may not be a meaningful marker of the activity of the GH-IGF-I system. Alternatively, increased IGF-I secretion from the liver may be quickly sequestered by the tissue in order to maintain an homeostatic balance of IGF-I in the systemic circulation (Kraemer et al., 1999), which may explain why we observed little change in these endocrine measures with either intervention but pronounced gains in muscle strength. Finally, it is also possible that the issue of timing of the sera sampling might have played a role in our observations and that hormonal adaptations may have occurred in a more acute fashion, which we would have missed (Kraemer & Ratamess, 2005). In addition, it is notable that IGF-I basal levels in our two populations tended to show high values at baseline in the HighR group, and for IGFBP-3 levels to be different at baseline between the two groups. In the case of IGF-I, this was possibly owing to the gender repartition in that the HighR group had more females and these tend to exhibit higher levels of the ligand (Brabant & Wallaschofski, 2007). In the case of IGFBP-3, the between groups difference might be owing to the overall higher number of female participants in LowR compared with HighR (Berrigan et al., 2008).

The increments in muscle strength found in the present study do not appear to be due to an increase in muscle mass resulting from an increase in the myofibrillar protein synthesis rate. Of course an increased synthetic rate is insufficient to stimulate muscle hypertrophy. Notably, ageing is also associated with increases in the protein breakdown mechanism (Trappe et al., 2004), and the impairment of this signalling may be an alternative pathway for maintained and/or improved skeletal muscle mass and hence performance. Our data illustrated significantly decreased TNF- $\alpha$ levels in the lower intensity exercise group only, with the high intensity trained counterparts exhibiting no significant IL-6 or TNF- $\alpha$  responses.

Physiological age-dependant hormonal changes may partly explain the impaired muscle protein synthesis in older adults. A subsequent increase in levels of catabolic cytokines, particularly TNF- $\alpha$  may suppress protein synthesis (Sakurai, Zhang, & Wolfe, 1996) and exacerbate protein

degradation in the muscle (Bales & Ritchie, 2002; Moulias, Meaume, & Raynaud-Simon, 1999). Resistance exercise may attenuate this process by suppressing skeletal muscle TNF- $\alpha$  expression (Greiwe, Cheng, Rubin, Yarasheski, & Semenkovich, 2001), an effect seen in the LowR group only in this study. However in the case of high intensity exercise, there may be a potential for greater 'micro-muscle-damage' (Nikolaidis et al., 2008) resulting in a changed inflammatory response with this exercise intensity. The strength gains observed in the present study for HighR strengthen claims that these gains may be attributed to neural factors, however, acute adaptations of anabolic hormones cannot be ruled out. An inverse relationship between local levels of TNF- $\alpha$  and skeletal muscle protein synthesis in frail older adults has been hypothesized (Greiwe et al., 2001), and this is in agreement with the effect shown through the LowR intervention which stimulated a reduction in plasma concentrations of TNF- $\alpha$ . If increased myofibrillar protein synthesis was responsible for strength gains in the present study, one would postulate that circulating levels of TNF- $\alpha$  would decrease following 12 weeks of resistance training, as Greiwe et al (Greiwe et al., 2001) reported in an octogenarian population. The preference for decreased catabolic signaling for LowR in the present study was accompanied by modest strength gains compared to the strength gains in HighR. This suggests that low intensity exercise in older populations may be sufficient to help alleviate one important modulator of inflamed ageing, while not impacting on the phenotypic expression of skeletal muscle mass. What is clear however is that the hormonal pathways used for LowR appear different to those seen with HighR exercise.

Finaly NPY, the sympathetic cotransmitter which causes vasoconstriction, decreases coronary blood flow and decreases cardiac output (Hauser et al., 1996) was seen to diminish significantly with the two levels of exercise, hence denoting positive circulatory adaptations with exercise. The fact that the chronic circulating levels of the immunoreactive NPY decreased to a similar extent with both exercise intensities reinforces the effectiveness of lower exercise intensity in the older age group.

# Body composition and metabolism at the whole body as well as endocrine level

Our findings are that body fat content did not significantly change whilst the body weight of participants was maintained from baseline to the post-intervention phase. These data on body composition changes appear to be at odds with a previous study that demonstrated a decrease in body fat content following a programme of resistance exercise in older populations (Kitamura, 2003). It should however be noted that more than resistance training, aerobic exercise is generally

linked to decreased adiposity and even then, there are aerobic studies showing no body composition changes post-intervention (Tonino, 1989). This fact, combined with the fact that the current programme was limited in terms of aerobic work (participants did not carry out the aerobic phase of their programme at 85% of maximum Heart rate, nor was this phase more than 15-20 min per session altogether), may explain the lack of changes in this parameter in the current study. What is more, ageing is linked to a decline in resting metabolic rate (RMR), presumably owing to the ageing-associated loss of muscle mass (Welle & Nair, 1990), decreased physical activity (Vaughan, Zurlo, & Ravussin, 1991) and decreased rate of protein metabolism (Welle & Nair, 1990). The mechanism by which RMR is increased with resistance training is presumed to be linked to an increase in protein turnover and associated increased a) muscle protein synthesis, b) muscle-tissue damage and c) muscle-tissue repair. Thus, the fact that these functions were generally not significantly changed with the training interventions, may explain why any impact on body composition was largely limited. What is more, our current findings are aligned with the suggestions that maintenance of fat mass may be an important coping mechanism in the elderly, even in the presence of increased physical activity. Indeed loss of adipose tissue predisposes individuals, especially the elderly, to chronic skin ulcers, disturbances of body temperature and decreased energy reserves in the face of chronic illness. Thus, there is no evidence to date to suggest that reduction of adipogenesis and/or excessive fat loss per se have any beneficial effects (Zhu et al., 2007) in older persons. In contrast, impaired adipogenesis has been reported to be associated with insulin resistance i.e. increased glucose intolerance (Heilbronn et al., 2004; Yang et al., 2004), thus potentially contributing to the paradoxical development of type II diabetes in very old lean patients.

Moreover in the current study, whilst there were no significant changes or in fact any noticeable trends in the levels of insulin in the two groups, the high intensity trained participants in fact concluded the programme with significantly higher levels of glucose, whereas glucose levels in the low intensity-trained participants remained unchanged. A study by others proposes that exercise alone may not be sufficient to induce changes in fasting plasma glucose and insulin levels (Matthews et al., 1985). In the face no changes in body composition in our high intensity-trained study population, it is therefore not surprising that the glucose profiles also bucked the trend for expected (Ryan, 2000) reductions in glucose levels as well as fat content and/or central adiposity. It cannot be ruled out however, whether this study group in fact altered their habitual nutritional

intake, hence altering glucose metabolism through a different mechanism. It should also be pointed out here that although the glucose level changes were significant in HighR (~16.1% increment), they were not clinically relevant as the status of the participants remained healthy (absolute glucose levels increased from ~  $4.80 \pm 0.19$  mmol/L to  $5.51 \pm 0.08$  mmol/L, thus remaining within the expected healthy norm, post-intervention). Whatever the case, the effects of resistance training as a single intervention for impacting on plasma glucose warrants further investigations.

# Functional abilities changes with exercise training

Finally, the two training programmes employed in the present study appear to have enhanced habitual function, more successfully than previous studies' attempts to do so, particularly where the get-up-and-go test is concerned (Earles, Judge, & Gunnarsson, 2001; Fiatarone et al., 1990; Schlicht, Camaione, & Owen, 2001; Skelton et al., 1995). Nevertheless, our findings of improved function with training do corroborate with the findings of some earlier studies (Skelton, Young, & Greig, 1997; Taaffe, Duret, Wheeler, & Marcus, 1999), even where the lower intensity group is concerned.

## **Conclusion**

The two training intensities yielded a number of positive adaptations, including for both, improvements in strength and functional abilities. It is notable however that whilst *in vivo* physiologic function changes showed a clear advantage for older persons to carrying out high resistance training, at the endocrine level, such an advantage (at least chronically) is not seen. In fact, in terms of changes in sera levels of overnight fasted resting levels of glucose, IGFBP-3 and TNF- $\alpha$ , there appears to be an advantage to carrying out the lower intensity, long term exercises for the aged populations where only endocrine factors are taken into account. Together these studies suggest that mixed exercise intensity interventions should be advocated.

#### **References**

- Adamczak, M., Rzepka, E., Chudek, J., & Wiecek, A. (2005). Ageing and plasma adiponectin concentration in apparently healthy males and females. *Clin Endocrinol (Oxf)*, *62*(1), 114-118.
- Bales, C. W., & Ritchie, C. S. (2002). Sarcopenia, weight loss, and nutritional frailty in the elderly. *Annu Rev Nutr, 22*, 309-323.

- Baranowska, B., Bik, W., Baranowska-Bik, A., Wolinska-Witort, E., Szybinska, A., Martynska, L., et al. (2006). Neuroendocrine control of metabolic homeostasis in Polish centenarians. J Physiol Pharmacol, 57 Suppl 6, 55-61.
- Bermon, S., Ferrari, P., Bernard, P., Altare, S., & Dolisi, C. (1999). Responses of total and free insulin-like growth factor-I and insulin-like growth factor binding protein-3 after resistance exercise and training in elderly subjects. *Acta Physiol Scand*, 165(1), 51-56.
- Berrigan, D., Potischman, N., Dodd, K. W., Hursting, S. D., Lavigne, J., Barrett, J. C., et al. (2008). Race/ethnic variation in serum levels of IGF-I and IGFBP-3 in US adults. *Growth Horm IGF Res.*
- Brabant, G., & Wallaschofski, H. (2007). Normal levels of serum IGF-I: determinants and validity of current reference ranges. *Pituitary*, *10*(2), 129-133.
- Butcher, S. K., & Lord, J. M. (2004). Stress responses and innate immunity: aging as a contributory factor. *Aging Cell*, *3*(4), 151-160.
- Carmeli, E., Reznick, A. Z., Coleman, R., & Carmeli, V. (2000). Muscle strength and mass of lower extremities in relation to functional abilities in elderly adults. *Gerontology*, *46*(5), 249-257.
- Chandler, J. M., Duncan, P. W., Kochersberger, G., & Studenski, S. (1998). Is lower extremity strength gain associated with improvement in physical performance and disability in frail, community-dwelling elders? *Arch Phys Med Rehabil*, *79*(1), 24-30.
- Charette, S. L., McEvoy, L., Pyka, G., Snow-Harter, C., Guido, D., Wiswell, R. A., et al. (1991). Muscle hypertrophy response to resistance training in older women. *J Appl Physiol*, 70(5), 1912-1916.
- Chin, A. P. M. J., van Uffelen, J. G., Riphagen, I., & van Mechelen, W. (2008). The functional effects of physical exercise training in frail older people : a systematic review. *Sports Med*, *38*(9), 781-793.
- Coggan, A. R., Spina, R. J., King, D. S., Rogers, M. A., Brown, M., Nemeth, P. M., et al. (1992). Skeletal muscle adaptations to endurance training in 60- to 70-yr-old men and women. J Appl Physiol, 72(5), 1780-1786.
- Crewther, B., Keogh, J., Cronin, J., & Cook, C. (2006). Possible stimuli for strength and power adaptation: acute hormonal responses. *Sports Med*, *36*(3), 215-238.
- Doherty, T. J. (2003). Invited review: Aging and sarcopenia. J Appl Physiol, 95(4), 1717-1727.
- Earles, D. R., Judge, J. O., & Gunnarsson, O. T. (2001). Velocity training induces power-specific adaptations in highly functioning older adults. *Arch Phys Med Rehabil*, 82(7), 872-878.
- Eliakim, A., Moromisato, M., Moromisato, D., Brasel, J. A., Roberts, C., Jr., & Cooper, D. M. (1997). Increase in muscle IGF-I protein but not IGF-I mRNA after 5 days of endurance training in young rats. *Am J Physiol*, 273(4 Pt 2), R1557-1561.
- Ferri, A., Scaglioni, G., Pousson, M., Capodaglio, P., Van Hoecke, J., & Narici, M. V. (2003). Strength and power changes of the human plantar flexors and knee extensors in response to resistance training in old age. *Acta Physiol Scand*, 177(1), 69-78.
- Fiatarone, M. A., Marks, E. C., Ryan, N. D., Meredith, C. N., Lipsitz, L. A., & Evans, W. J. (1990). High-intensity strength training in nonagenarians. Effects on skeletal muscle. *Jama*, 263(22), 3029-3034.
- Fielding, R. A. (1995). The role of progressive resistance training and nutrition in the preservation of lean body mass in the elderly. *J Am Coll Nutr, 14*(6), 587-594.
- Frontera, W. R., Hughes, V. A., Fielding, R. A., Fiatarone, M. A., Evans, W. J., & Roubenoff, R. (2000). Aging of skeletal muscle: a 12-yr longitudinal study. *J Appl Physiol*, 88(4), 1321-1326.

- Frontera, W. R., Hughes, V. A., Krivickas, L. S., Kim, S. K., Foldvari, M., & Roubenoff, R. (2003). Strength training in older women: early and late changes in whole muscle and single cells. *Muscle Nerve*, 28(5), 601-608.
- Frontera, W. R., Meredith, C. N., O'Reilly, K. P., Knuttgen, H. G., & Evans, W. J. (1988). Strength conditioning in older men: skeletal muscle hypertrophy and improved function. *J Appl Physiol*, 64(3), 1038-1044.
- Greenlund, L. J., & Nair, K. S. (2003). Sarcopenia--consequences, mechanisms, and potential therapies. *Mech Ageing Dev, 124*(3), 287-299.
- Greiwe, J. S., Cheng, B., Rubin, D. C., Yarasheski, K. E., & Semenkovich, C. F. (2001). Resistance exercise decreases skeletal muscle tumor necrosis factor alpha in frail elderly humans. *FASEB J*, *15*(2), 475-482.
- Hakkinen, K., Kraemer, W. J., Newton, R. U., & Alen, M. (2001). Changes in electromyographic activity, muscle fibre and force production characteristics during heavy resistance/power strength training in middle-aged and older men and women. *Acta Physiol Scand*, 171(1), 51-62.
- Hartman, J. W., Moore, D. R., & Phillips, S. M. (2006). Resistance training reduces whole-body protein turnover and improves net protein retention in untrained young males. *Appl Physiol Nutr Metab*, 31(5), 557-564.
- Hauser, G. J., Danchak, M. R., Colvin, M. P., Hopkins, R. A., Wocial, B., Myers, A. K., et al. (1996). Circulating neuropeptide Y in humans: relation to changes in catecholamine levels and changes in hemodynamics. *Neuropeptides*, 30(2), 159-165.
- Heilbronn, L. K., Rood, J., Janderova, L., Albu, J. B., Kelley, D. E., Ravussin, E., et al. (2004).
  Relationship between serum resistin concentrations and insulin resistance in nonobese, obese, and obese diabetic subjects. *J Clin Endocrinol Metab*, 89(4), 1844-1848.
- Jones, J. I., & Clemmons, D. R. (1995). Insulin-like growth factors and their binding proteins: biological actions. *Endocr Rev, 16*(1), 3-34.
- Jorgensen, J. O., Vahl, N., Hansen, T. B., Thuesen, L., Hagen, C., & Christiansen, J. S. (1996). Growth hormone versus placebo treatment for one year in growth hormone deficient adults: increase in exercise capacity and normalization of body composition. *Clin Endocrinol (Oxf)*, 45(6), 681-688.
- Kitamura, I. T., N. Tokudome, M. Yamanouchi, K. Oshida, Y. Sato, Y. (2003). Effects of aerobic and resistance training on insulin action in the elderly. *Geriatrics and Gerontology International, 3*, 50-55.
- Klitgaard, H., Mantoni, M., Schiaffino, S., Ausoni, S., Gorza, L., Laurent-Winter, C., et al. (1990). Function, morphology and protein expression of ageing skeletal muscle: a cross-sectional study of elderly men with different training backgrounds. *Acta Physiol Scand*, 140(1), 41-54.
- Kosek, D. J., Kim, J. S., Petrella, J. K., Cross, J. M., & Bamman, M. M. (2006). Efficacy of 3 days/wk resistance training on myofiber hypertrophy and myogenic mechanisms in young vs. older adults. *J Appl Physiol*, 101(2), 531-544.
- Kraemer, W. J., Hakkinen, K., Newton, R. U., Nindl, B. C., Volek, J. S., McCormick, M., et al. (1999). Effects of heavy-resistance training on hormonal response patterns in younger vs. older men. *J Appl Physiol*, 87(3), 982-992.
- Kraemer, W. J., & Ratamess, N. A. (2005). Hormonal responses and adaptations to resistance exercise and training. *Sports Med*, *35*(4), 339-361.
- Lexell, J. (1995). Human aging, muscle mass, and fiber type composition. J Gerontol A Biol Sci Med Sci, 50 Spec No, 11-16.

- Lexell, J., Taylor, C. C., & Sjostrom, M. (1988). What is the cause of the ageing atrophy? Total number, size and proportion of different fiber types studied in whole vastus lateralis muscle from 15- to 83-year-old men. *J Neurol Sci*, *84*(2-3), 275-294.
- Lintsi, M., Kaarma, H., & Kull, I. (2004). Comparison of hand-to-hand bioimpedance and anthropometry equations versus dual-energy X-ray absorptiometry for the assessment of body fat percentage in 17-18-year-old conscripts. *Clin Physiol Funct Imaging, 24*(2), 85-90.
- Matthews, D. R., Hosker, J. P., Rudenski, A. S., Naylor, B. A., Treacher, D. F., & Turner, R. C. (1985). Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*, *28*(7), 412-419.
- Moulias, R., Meaume, S., & Raynaud-Simon, A. (1999). Sarcopenia, hypermetabolism, and aging. Z Gerontol Geriatr, 32(6), 425-432.
- Mroszczyk-McDonald, A., Savage, P. D., & Ades, P. A. (2007). Handgrip strength in cardiac rehabilitation: normative values, interaction with physical function, and response to training. *J Cardiopulm Rehabil Prev*, 27(5), 298-302.
- Narici, M. V., Maganaris, C. N., Reeves, N. D., & Capodaglio, P. (2003). Effect of aging on human muscle architecture. *J Appl Physiol*, *95*(6), 2229-2234.
- Nikolaidis, M. G., Jamurtas, A. Z., Paschalis, V., Fatouros, I. G., Koutedakis, Y., & Kouretas, D. (2008). The effect of muscle-damaging exercise on blood and skeletal muscle oxidative stress: magnitude and time-course considerations. *Sports Med*, *38*(7), 579-606.
- Onambele, G. L., Maganaris, C. N., Mian, O. S., Tam, E., Rejc, E., McEwan, I. M., et al. (2008). Neuromuscular and balance responses to flywheel inertial versus weight training in older persons. *J Biomech*, 41(15), 3133-3138.
- Pearson, S. J., & Onambele, G. N. (2005). Acute changes in knee-extensors torque, fiber pennation, and tendon characteristics. *Chronobiol Int*, 22(6), 1013-1027.
- Pearson, S. J., & Onambele, G. N. (2006). Influence of time of day on tendon compliance and estimations of voluntary activation levels. *Muscle Nerve*, *33*(6), 792-800.
- Pearson, S. J., Young, A., Macaluso, A., Devito, G., Nimmo, M. A., Cobbold, M., et al. (2002). Muscle function in elite master weightlifters. *Med Sci Sports Exerc*, 34(7), 1199-1206.
- Poulin, M. J., Vandervoort, A. A., Paterson, D. H., Kramer, J. F., & Cunningham, D. A. (1992). Eccentric and concentric torques of knee and elbow extension in young and older men. *Can J Sport Sci*, 17(1), 3-7.
- Prior, B. M., Cureton, K. J., Modlesky, C. M., Evans, E. M., Sloniger, M. A., Saunders, M., et al. (1997). In vivo validation of whole body composition estimates from dual-energy X-ray absorptiometry. *J Appl Physiol*, 83(2), 623-630.
- Rennie, M. J., Selby, A., Atherton, P., Smith, K., Kumar, V., Glover, E. L., et al. (2009). Facts, noise and wishful thinking: muscle protein turnover in aging and human disuse atrophy. *Scand J Med Sci Sports*.
- Roubenoff, R., Hughes, V. A., Dallal, G. E., Nelson, M. E., Morganti, C., Kehayias, J. J., et al. (2000). The effect of gender and body composition method on the apparent decline in lean mass-adjusted resting metabolic rate with age. *J Gerontol A Biol Sci Med Sci*, 55(12), M757-760.
- Ryan, A. S. (2000). Insulin resistance with aging: effects of diet and exercise. *Sports Med*, 30(5), 327-346.
- Sakurai, Y., Zhang, X. J., & Wolfe, R. R. (1996). TNF directly stimulates glucose uptake and leucine oxidation and inhibits FFA flux in conscious dogs. *Am J Physiol*, 270(5 Pt 1), E864-872.

- Schlicht, J., Camaione, D. N., & Owen, S. V. (2001). Effect of intense strength training on standing balance, walking speed, and sit-to-stand performance in older adults. J Gerontol A Biol Sci Med Sci, 56(5), M281-286.
- Schutte, A. E., Huisman, H. W., Schutte, R., van Rooyen, J. M., Malan, L., & Malan, N. T. (2007). Aging influences the level and functions of fasting plasma ghrelin levels: the POWIRS-Study. *Regul Pept*, 139(1-3), 65-71.
- Sipila, S., Elorinne, M., Alen, M., Suominen, H., & Kovanen, V. (1997). Effects of strength and endurance training on muscle fibre characteristics in elderly women. *Clin Physiol*, 17(5), 459-474.
- Skelton, D. A., Young, A., & Greig, C. A. (1997). Muscle function of women aged 65-89 years meeting two sets of health criteria. *Aging (Milano), 9*(1-2), 106-111.
- Skelton, D. A., Young, A., Greig, C. A., & Malbut, K. E. (1995). Effects of resistance training on strength, power, and selected functional abilities of women aged 75 and older. J Am Geriatr Soc, 43(10), 1081-1087.
- Taaffe, D. R., Duret, C., Wheeler, S., & Marcus, R. (1999). Once-weekly resistance exercise improves muscle strength and neuromuscular performance in older adults. J Am Geriatr Soc, 47(10), 1208-1214.
- Tonino, R. P. (1989). Effect of physical training on the insulin resistance of aging. *Am J Physiol*, 256(3 Pt 1), E352-356.
- Tracy, B. L., Ivey, F. M., Hurlbut, D., Martel, G. F., Lemmer, J. T., Siegel, E. L., et al. (1999). Muscle quality. II. Effects Of strength training in 65- to 75-yr-old men and women. J Appl Physiol, 86(1), 195-201.
- Trappe, T., Williams, R., Carrithers, J., Raue, U., Esmarck, B., Kjaer, M., et al. (2004). Influence of age and resistance exercise on human skeletal muscle proteolysis: a microdialysis approach. *J Physiol*, 554(Pt 3), 803-813.
- Vandervoort, A. A., Kramer, J. F., & Wharram, E. R. (1990). Eccentric knee strength of elderly females. J Gerontol, 45(4), B125-128.
- Vaughan, L., Zurlo, F., & Ravussin, E. (1991). Aging and energy expenditure. *Am J Clin Nutr,* 53(4), 821-825.
- Velloso, C. P. (2008). Regulation of muscle mass by growth hormone and IGF-I. *Br J Pharmacol*, *154*(3), 557-568.
- Welle, S., & Nair, K. S. (1990). Relationship of resting metabolic rate to body composition and protein turnover. *Am J Physiol*, *258*(6 Pt 1), E990-998.
- Welle, S., Totterman, S., & Thornton, C. (1996). Effect of age on muscle hypertrophy induced by resistance training. J Gerontol A Biol Sci Med Sci, 51(6), M270-275.
- Yang, X., Jansson, P. A., Nagaev, I., Jack, M. M., Carvalho, E., Sunnerhagen, K. S., et al. (2004). Evidence of impaired adipogenesis in insulin resistance. *Biochem Biophys Res Commun*, 317(4), 1045-1051.
- Yarasheski, K. E. (2003). Exercise, aging, and muscle protein metabolism. J Gerontol A Biol Sci Med Sci, 58(10), M918-922.
- Young, A., Stokes, M., & Crowe, M. (1985). The size and strength of the quadriceps muscles of old and young men. *Clin Physiol*, *5*(2), 145-154.
- Zhu, M., Lee, G. D., Ding, L., Hu, J., Qiu, G., de Cabo, R., et al. (2007). Adipogenic signaling in rat white adipose tissue: modulation by aging and calorie restriction. *Exp Gerontol*, 42(8), 733-744.

# **Tables, Figures and Legends**

Group	Gender	Ν	Age (Yrs)	Activity (mins/wk)	Height (m)	Weight (Kg)
LowR	Male	10	76 ± 2	245 ± 214	1.7 ± 0.05	79.1 ± 9.4
	Female	8	76 ± 4	396 ± 457	1.58 ± 0.05	68.3 ± 11.7
	All	18	76.3 ± 3.4	320.6 ± 335.6	1.64 ± 0.05	73.7 ± 10.6
HighR	Male	6 (4*)	68 ± 7	273 ± 175	1.72 ± 0.03	81.7 ± 8.9
	Female	6	66 ± 4.5	313 ± 183	1.6 ± 0.04	70.6 ± 13.2
	All	12	69 ± 5.8	293.3 ± 179	1.66 ± 0.04	76.2 ± 11.1

<u>**Table 1:**</u> Baseline characteristics of completing participant. (4\*) Indicates the number of participants who dropped out of the study before completion. Means  $\pm$  S.D.

<u>Table 2:</u> Change in 1RM load lifted (Kg), and maximum isometric torque (Nm) from pre- to post 12-week intervention. \*Indicates significantly lower increment in LowR compared to HighR (P < 0.05). Values are Mean  $\pm$  S.E.M.

		LowR		HighR		
	PRE	POST	Δ (%)	PRE	POST	Δ (%)
Leg Press (Kg)	90.4 ± 6.2	105.5 ± 7.1	*17	89.7 ± 7.7	125.4 ± 7.5	40
Leg Extensor (Kg)	30.5 ± 2.7	35.5 ± 3	*16	28.3 ± 1.8	42.8 ± 1.9	51
Calf Rotator (Kg)	34.7 ± 2	42.1 ± 2.4	*21	33.4 ± 3.2	55.8 ± 3.8	67
Glute Conditioner (Kg)	55 ± 4.1	63.1 ± 4	*15	45.7 ± 4.1	67.4 ± 5	47
Unilateral MVC (Nm)	122.2 ± 9.5	122.8 ± 9.3	1.3 ± 3.0	122.6 ± 12.3	133.0 ± 13.6	9.3 ± 5.1
Bilateral MVC (Nm)	152.0 ± 13.0	165.7 ± 14.0	10.3 ± 4.7	137.5 ± 19.2	160.3 ± 18.3	22.4 ± 10.2

**Figure 1:** Illustration of types of exercises carried out during the supervised classes. The hour-long classes incorporated an aerobic warm-up component (Fig 1a), a core with resistance work (Figs 1b and 1c), as well as well as a cool down phase. It is notable that the protocol was similar for the home-based / unsupervised sessions with the exception that no resistance machines were utilised (but therabands instead) and the aerobic phase consisted of a 20mins brisk walk outdoors.



**Figure 2:** Summary of endocrine characteristics changes with the two levels of exercise (high versus low intensity). The \* indicates where post-intervention data were significantly different to baseline values, with P<0.05.

